



wwPDB EM Validation Summary Report ⓘ

Oct 3, 2022 – 11:05 pm BST

PDB ID : 7Z46
EMDB ID : EMD-14485
Title : Top part (C5) of bacteriophage SU10 capsid
Authors : Siborova, M.; Fuzik, T.; Prochazkova, M.; Novacek, J.; Plevka, P.
Deposited on : 2022-03-03
Resolution : 3.40 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	11% 98%
1	B	352	98%
1	C	352	5% 99%
1	D	352	6% 98%
1	E	352	6% 97%
1	F	352	98%
1	G	352	5% 98%
1	H	352	97%
1	I	352	5% 98%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	352	97%
1	K	352	98%
1	L	352	98%
1	M	352	98%
1	N	352	99%
1	O	352	99%
1	P	352	97%
1	Q	352	98%
1	R	352	97%
1	S	352	97%
1	T	352	98%
1	U	352	98%
1	V	352	98%
1	W	352	98%
1	X	352	98%
1	Y	352	98%
1	Z	352	99%
1	a	352	98%
1	b	352	98%
1	c	352	99%
1	d	352	99%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 80206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major head protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	345	2672	1676	467	519	10	0	0
1	B	346	2677	1680	468	519	10	0	0
1	C	347	2685	1684	470	521	10	0	0
1	D	345	2670	1675	467	518	10	0	0
1	E	346	2677	1680	468	519	10	0	0
1	F	346	2676	1678	468	520	10	0	0
1	G	347	2685	1684	470	521	10	0	0
1	H	343	2657	1668	465	514	10	0	0
1	I	346	2676	1678	468	520	10	0	0
1	J	342	2644	1656	464	514	10	0	0
1	K	347	2685	1684	470	521	10	0	0
1	L	347	2685	1684	470	521	10	0	0
1	M	346	2676	1678	468	520	10	0	0
1	N	347	2685	1684	470	521	10	0	0
1	O	347	2685	1684	470	521	10	0	0
1	P	343	2654	1665	464	515	10	0	0
1	Q	346	2676	1678	468	520	10	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	342	Total	C	N	O	S	0	0
			2644	1656	464	514	10		
1	S	341	Total	C	N	O	S	0	0
			2637	1651	463	513	10		
1	T	345	Total	C	N	O	S	0	0
			2668	1674	466	518	10		
1	U	346	Total	C	N	O	S	0	0
			2676	1678	468	520	10		
1	V	345	Total	C	N	O	S	0	0
			2670	1675	467	518	10		
1	W	347	Total	C	N	O	S	0	0
			2685	1684	470	521	10		
1	X	346	Total	C	N	O	S	0	0
			2677	1680	468	519	10		
1	Y	346	Total	C	N	O	S	0	0
			2676	1678	468	520	10		
1	Z	347	Total	C	N	O	S	0	0
			2685	1684	470	521	10		
1	a	346	Total	C	N	O	S	0	0
			2677	1680	468	519	10		
1	b	346	Total	C	N	O	S	0	0
			2676	1678	468	520	10		
1	c	347	Total	C	N	O	S	0	0
			2685	1684	470	521	10		
1	d	347	Total	C	N	O	S	0	0
			2685	1684	470	521	10		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major head protein




- Molecule 1: Major head protein



- Molecule 1: Major head protein




- Molecule 1: Major head protein



- Molecule 1: Major head protein






• Molecule 1: Major head protein




• Molecule 1: Major head protein




• Molecule 1: Major head protein




• Molecule 1: Major head protein




• Molecule 1: Major head protein




• Molecule 1: Major head protein






- Molecule 1: Major head protein

Chain L: 98%



- Molecule 1: Major head protein

Chain M: 98%





- Molecule 1: Major head protein

Chain N: 99%





- Molecule 1: Major head protein

Chain O: 10% 99%



- Molecule 1: Major head protein

Chain P: 10% 97%




- Molecule 1: Major head protein

Chain Q:  98%




- Molecule 1: Major head protein

Chain R:  12% 97%




- Molecule 1: Major head protein

Chain S:  17% 97%




- Molecule 1: Major head protein

Chain T:  11% 98%




- Molecule 1: Major head protein

Chain U:  98%



- Molecule 1: Major head protein


Chain V:  47% 98%




• Molecule 1: Major head protein




• Molecule 1: Major head protein




• Molecule 1: Major head protein



• Molecule 1: Major head protein




• Molecule 1: Major head protein




• Molecule 1: Major head protein



• Molecule 1: Major head protein



• Molecule 1: Major head protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C5	Depositor
Number of particles used	22146	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	49	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2700	Depositor
Magnification	59000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.239	Depositor
Minimum map value	-0.136	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.011	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	552.0, 552.0, 552.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.38, 1.38, 1.38	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2723	0.44	0/3690
1	B	0.24	0/2728	0.44	0/3694
1	C	0.25	0/2736	0.43	0/3706
1	D	0.25	0/2720	0.43	0/3683
1	E	0.24	0/2728	0.45	0/3694
1	F	0.25	0/2727	0.44	0/3695
1	G	0.24	0/2736	0.46	0/3706
1	H	0.25	0/2706	0.45	0/3661
1	I	0.24	0/2727	0.44	0/3695
1	J	0.25	0/2693	0.43	0/3646
1	K	0.25	0/2736	0.45	0/3706
1	L	0.24	0/2736	0.42	0/3706
1	M	0.25	0/2727	0.45	0/3695
1	N	0.25	0/2736	0.44	0/3706
1	O	0.24	0/2736	0.44	0/3706
1	P	0.25	0/2704	0.43	0/3662
1	Q	0.25	0/2727	0.45	0/3695
1	R	0.25	0/2693	0.44	0/3646
1	S	0.25	0/2686	0.45	0/3636
1	T	0.24	0/2719	0.43	0/3683
1	U	0.25	0/2727	0.44	0/3695
1	V	0.25	0/2720	0.44	0/3683
1	W	0.25	0/2736	0.43	0/3706
1	X	0.24	0/2728	0.42	0/3694
1	Y	0.25	0/2727	0.45	0/3695
1	Z	0.25	0/2736	0.43	0/3706
1	a	0.24	0/2728	0.45	0/3694
1	b	0.25	0/2727	0.43	0/3695
1	c	0.25	0/2736	0.44	0/3706
1	d	0.25	0/2736	0.46	0/3706
All	All	0.25	0/81725	0.44	0/110691

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected

by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
1	G	0	2
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	144	THR	Peptide
1	G	167	PHE	Peptide
1	G	168	GLN	Peptide

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/352 (97%)	286 (83%)	57 (17%)	0	100	100
1	B	344/352 (98%)	284 (83%)	59 (17%)	1 (0%)	41	72
1	C	345/352 (98%)	309 (90%)	36 (10%)	0	100	100
1	D	343/352 (97%)	308 (90%)	35 (10%)	0	100	100
1	E	344/352 (98%)	272 (79%)	68 (20%)	4 (1%)	13	41
1	F	344/352 (98%)	298 (87%)	46 (13%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	345/352 (98%)	288 (84%)	56 (16%)	1 (0%)	41	72
1	H	339/352 (96%)	303 (89%)	36 (11%)	0	100	100
1	I	344/352 (98%)	322 (94%)	22 (6%)	0	100	100
1	J	340/352 (97%)	308 (91%)	32 (9%)	0	100	100
1	K	345/352 (98%)	276 (80%)	68 (20%)	1 (0%)	41	72
1	L	345/352 (98%)	314 (91%)	31 (9%)	0	100	100
1	M	344/352 (98%)	318 (92%)	26 (8%)	0	100	100
1	N	345/352 (98%)	314 (91%)	31 (9%)	0	100	100
1	O	345/352 (98%)	264 (76%)	81 (24%)	0	100	100
1	P	341/352 (97%)	294 (86%)	47 (14%)	0	100	100
1	Q	344/352 (98%)	324 (94%)	20 (6%)	0	100	100
1	R	340/352 (97%)	310 (91%)	30 (9%)	0	100	100
1	S	339/352 (96%)	287 (85%)	52 (15%)	0	100	100
1	T	343/352 (97%)	306 (89%)	37 (11%)	0	100	100
1	U	344/352 (98%)	313 (91%)	31 (9%)	0	100	100
1	V	343/352 (97%)	307 (90%)	36 (10%)	0	100	100
1	W	345/352 (98%)	298 (86%)	47 (14%)	0	100	100
1	X	344/352 (98%)	312 (91%)	32 (9%)	0	100	100
1	Y	344/352 (98%)	319 (93%)	25 (7%)	0	100	100
1	Z	345/352 (98%)	323 (94%)	22 (6%)	0	100	100
1	a	344/352 (98%)	284 (83%)	60 (17%)	0	100	100
1	b	344/352 (98%)	328 (95%)	16 (5%)	0	100	100
1	c	345/352 (98%)	311 (90%)	34 (10%)	0	100	100
1	d	345/352 (98%)	286 (83%)	59 (17%)	0	100	100
All	All	10305/10560 (98%)	9066 (88%)	1232 (12%)	7 (0%)	54	82

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	44	GLN
1	G	168	GLN
1	B	20	ALA
1	E	45	THR
1	E	145	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/287 (99%)	284 (100%)	1 (0%)	91	95
1	B	285/287 (99%)	284 (100%)	1 (0%)	91	95
1	C	286/287 (100%)	286 (100%)	0	100	100
1	D	284/287 (99%)	284 (100%)	0	100	100
1	E	285/287 (99%)	285 (100%)	0	100	100
1	F	285/287 (99%)	285 (100%)	0	100	100
1	G	286/287 (100%)	286 (100%)	0	100	100
1	H	283/287 (99%)	283 (100%)	0	100	100
1	I	285/287 (99%)	285 (100%)	0	100	100
1	J	281/287 (98%)	281 (100%)	0	100	100
1	K	286/287 (100%)	286 (100%)	0	100	100
1	L	286/287 (100%)	284 (99%)	2 (1%)	84	92
1	M	285/287 (99%)	283 (99%)	2 (1%)	84	92
1	N	286/287 (100%)	286 (100%)	0	100	100
1	O	286/287 (100%)	286 (100%)	0	100	100
1	P	282/287 (98%)	282 (100%)	0	100	100
1	Q	285/287 (99%)	283 (99%)	2 (1%)	84	92
1	R	281/287 (98%)	281 (100%)	0	100	100
1	S	280/287 (98%)	279 (100%)	1 (0%)	91	95
1	T	284/287 (99%)	283 (100%)	1 (0%)	91	95
1	U	285/287 (99%)	283 (99%)	2 (1%)	84	92
1	V	284/287 (99%)	283 (100%)	1 (0%)	91	95
1	W	286/287 (100%)	285 (100%)	1 (0%)	92	97
1	X	285/287 (99%)	284 (100%)	1 (0%)	91	95
1	Y	285/287 (99%)	283 (99%)	2 (1%)	84	92
1	Z	286/287 (100%)	286 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	a	285/287 (99%)	285 (100%)	0	100	100
1	b	285/287 (99%)	285 (100%)	0	100	100
1	c	286/287 (100%)	286 (100%)	0	100	100
1	d	286/287 (100%)	286 (100%)	0	100	100
All	All	8539/8610 (99%)	8522 (100%)	17 (0%)	93	98

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	256	ARG
1	Y	298	ARG
1	Q	264	LYS
1	S	68	ARG
1	T	163	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	V	279	GLN
1	a	102	ASN
1	W	262	ASN
1	Y	141	GLN
1	a	279	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14485. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 58




Y Index: 347




Z Index: 210


6.3.2 Raw map



X Index: 66



Y Index: 335




Z Index: 211

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 3366 nm^3 ; this corresponds to an approximate mass of 3040 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.35	3.77	3.40
Unmasked-calculated*	3.76	4.53	3.84


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.76 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14485 and PDB model 7Z46. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlays

9.1.1 Map-model overlay [i](#)




9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).































































9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7553	 0.3830
A	 0.7265	 0.3210
B	 0.8091	 0.3980
C	 0.8086	 0.4110
D	 0.7957	 0.4000
E	 0.7958	 0.3920
F	 0.8216	 0.4260
G	 0.7817	 0.3770
H	 0.8291	 0.4230
I	 0.8186	 0.4010
J	 0.8260	 0.4230
K	 0.7563	 0.3360
L	 0.8415	 0.4430
M	 0.8566	 0.4390
N	 0.8412	 0.4280
O	 0.7551	 0.3370
P	 0.7787	 0.3900
Q	 0.8387	 0.4240
R	 0.7174	 0.3890
S	 0.6792	 0.3430
T	 0.7478	 0.3530
U	 0.8421	 0.4290
V	 0.4041	 0.2520
W	 0.3533	 0.2210
X	 0.8384	 0.4440
Y	 0.8440	 0.4330
Z	 0.5785	 0.3330
a	 0.5757	 0.3180
b	 0.8448	 0.4540
c	 0.7987	 0.4040
d	 0.7532	 0.3550

