

wwPDB X-ray Structure Validation Summary Report (i)

Nov 7, 2023 – 01:11 PM JST

PDB ID	:	4Z63
Title	:	The plant peptide hormone receptor in arabidopsis
Authors	:	Chai, J.; Wang, J.; Han, Z.
Deposited on	:	2015-04-03
Resolution	:	2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630(2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Q	uality of chain	
1	А	631	4% 78°	%	17% ••
2	Р	5	40%	40%	20%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5075 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Phytosulfokine receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	610	Total 4748	C 3003	N 829	O 897	S 19	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
А	649	HIS	-	expression tag	UNP Q9ZVR7
А	650	HIS	-	expression tag	UNP Q9ZVR7
А	651	HIS	-	expression tag	UNP Q9ZVR7
A	652	HIS	-	expression tag	UNP Q9ZVR7
А	653	HIS	-	expression tag	UNP Q9ZVR7
А	654	HIS	_	expression tag	UNP Q9ZVR7

• Molecule 2 is a protein called Phytosulfokine.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	Р	5	Total 57	C 33		O 16	$\begin{array}{c} \mathrm{S} \\ \mathrm{2} \end{array}$	0	0	0

• Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 14 8 1 5	0	0
3	А	1	Total C N O 13 8 1 4	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	143	Total O 143 143	0	0
4	Р	2	Total O 2 2	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Phytosulfokine receptor 1





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants	92.89Å 92.89Å 242.49Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.54 - 2.51	Depositor
Resolution (A)	33.54 - 2.51	EDS
% Data completeness	98.0 (33.54-2.51)	Depositor
(in resolution range)	97.8(33.54-2.51)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$5.62 (at 2.51 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
D D.	0.229 , 0.276	Depositor
R, R_{free}	0.229 , 0.273	DCC
R_{free} test set	1821 reflections (5.00%)	wwPDB-VP
Wilson B-factor $(Å^2)$	37.2	Xtriage
Anisotropy	0.691	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35 , 30.6	EDS
L-test for twinning ²	$ < L >=0.51, < L^2>=0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5075	wwPDB-VP
Average B, all atoms $(Å^2)$	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain B		nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.50	2/4850~(0.0%)	0.71	3/6572~(0.0%)	
2	Р	0.51	0/23	0.93	0/27	
All	All	0.50	2/4873~(0.0%)	0.71	3/6599~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	492	ARG	CA-C	6.06	1.68	1.52
1	А	358	GLU	CG-CD	5.15	1.59	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	491	SER	O-C-N	-9.73	107.13	122.70
1	А	358	GLU	OE1-CD-OE2	-6.81	115.13	123.30
1	А	492	ARG	N-CA-C	5.65	126.26	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:



4Z63

Mol	Chain	Res	Type	Group
1	А	491	SER	Mainchain,Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4748	0	4710	59	1
2	Р	57	0	44	1	0
3	А	125	0	115	5	0
4	А	143	0	0	7	0
4	Р	2	0	0	0	0
All	All	5075	0	4869	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 61 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:484:THR:HG21	1:A:542:GLU:HB3	1.42	0.99
1:A:484:THR:HG21	1:A:542:GLU:CB	2.02	0.88
1:A:140:ILE:HG13	1:A:164:PRO:HG3	1.66	0.77
1:A:331:ARG:NH1	4:A:803:HOH:O	2.17	0.77
1:A:484:THR:HG23	1:A:542:GLU:O	1.87	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:ASP:OD1	1:A:511:GLU:OE1[3_564]	2.06	0.14



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	А	608/631~(96%)	592 (97%)	15~(2%)	1 (0%)	47	67
2	Р	2/5~(40%)	2(100%)	0	0	100	100
All	All	610/636~(96%)	594 (97%)	15~(2%)	1 (0%)	47	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	615	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	548/567~(97%)	512~(93%)	36 (7%)	16 30
2	Р	3/3~(100%)	2~(67%)	1 (33%)	0 0
All	All	551/570~(97%)	514 (93%)	37~(7%)	16 29

5 of 37 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	529	GLU
1	А	633	GLU
1	А	562	SER
1	А	591	LEU
1	А	265	ASP



Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	А	242	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain Res Link		Bo	Bond lengths			Bond angles			
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	TYS	Р	28	2	$15,\!16,\!17$	2.06	2 (13%)	18,22,24	0.72	0
2	TYS	Р	30	2	15,16,17	2.01	3 (20%)	18,22,24	0.85	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYS	Р	28	2	-	0/10/11/13	0/1/1/1
2	TYS	Р	30	2	-	0/10/11/13	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Р	28	TYS	OH-CZ	-7.26	1.30	1.42
2	Р	30	TYS	OH-CZ	-6.72	1.31	1.42
2	Р	30	TYS	OH-S	2.78	1.62	1.58
2	Р	28	TYS	O2-S	2.23	1.54	1.45
2	Р	30	TYS	O2-S	2.15	1.54	1.45



All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Р	30	TYS	O3-S-OH	2.74	112.44	105.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
IVIOI	Type	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	NAG	А	701	1	$14,\!14,\!15$	0.62	0	$17,\!19,\!21$	1.56	2 (11%)
3	NAG	А	707	1	14,14,15	0.55	0	17,19,21	0.84	0
3	NAG	А	705	1	$14,\!14,\!15$	0.48	0	$17,\!19,\!21$	1.02	1 (5%)
3	NAG	А	708	1	14,14,15	0.59	0	17,19,21	1.29	2 (11%)
3	NAG	А	706	1	14,14,15	0.62	0	17,19,21	1.37	2 (11%)
3	NAG	А	703	1	$14,\!14,\!15$	0.48	0	17,19,21	1.00	0
3	NAG	А	704	1	$14,\!14,\!15$	0.50	0	$17,\!19,\!21$	1.64	4 (23%)
3	NAG	А	702	1	$14,\!14,\!15$	0.59	0	17,19,21	1.22	2 (11%)
3	NAG	А	709	-	13,13,15	0.92	1 (7%)	16,17,21	2.60	<mark>6 (37%)</mark>

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



4Z63

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	А	701	1	-	0/6/23/26	0/1/1/1
3	NAG	А	707	1	-	2/6/23/26	0/1/1/1
3	NAG	А	705	1	-	0/6/23/26	0/1/1/1
3	NAG	А	708	1	-	2/6/23/26	0/1/1/1
3	NAG	А	706	1	-	0/6/23/26	0/1/1/1
3	NAG	А	703	1	-	0/6/23/26	0/1/1/1
3	NAG	А	704	1	-	2/6/23/26	0/1/1/1
3	NAG	А	702	1	-	2/6/23/26	0/1/1/1
3	NAG	А	709	-	-	2/6/19/26	0/1/1/1

'-' means no outliers of that kind were identified.

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	А	709	NAG	C3-C2	-2.26	1.49	1.52

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	А	709	NAG	C4-C3-C2	-5.88	105.26	112.53
3	А	701	NAG	C1-O5-C5	4.90	118.83	112.19
3	А	709	NAG	C3-C4-C5	-4.37	105.10	110.77
3	А	709	NAG	C3-C2-N2	-4.33	104.00	110.57
3	А	709	NAG	O5-C5-C6	4.01	113.50	107.20

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	709	NAG	C4-C5-C6-O6
3	А	709	NAG	O5-C5-C6-O6
3	А	708	NAG	O5-C5-C6-O6
3	А	704	NAG	C4-C5-C6-O6
3	А	704	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	705	NAG	2	0

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	709	NAG	3	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	610/631~(96%)	-0.05	25 (4%) 37 41	1, 10, 28, 52	0
2	Р	3/5~(60%)	0.55	0 100 100	3, 3, 8, 20	0
All	All	613/636~(96%)	-0.05	25 (4%) 37 41	1, 10, 28, 52	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	411	SER	5.1
1	А	493	ASN	5.1
1	А	75	THR	4.1
1	А	494	ILE	3.7
1	А	72	SER	3.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	TYS	Р	30	16/17	0.94	0.13	$1,\!6,\!22,\!37$	0
2	TYS	Р	28	16/17	0.96	0.13	2,4,13,18	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
3	NAG	А	709	13/15	0.80	0.19	19,36,42,46	0
3	NAG	А	706	14/15	0.85	0.29	15,30,36,46	0
3	NAG	А	708	14/15	0.91	0.21	8,28,36,37	0
3	NAG	А	707	14/15	0.91	0.25	12,16,28,33	0
3	NAG	А	703	14/15	0.92	0.14	9,15,22,31	0
3	NAG	А	705	14/15	0.92	0.17	$9,\!17,\!27,\!28$	0
3	NAG	А	704	14/15	0.94	0.14	1,7,17,19	0
3	NAG	А	701	14/15	0.96	0.14	2,5,8,10	0
3	NAG	А	702	14/15	0.97	0.12	4,7,12,14	0

6.5 Other polymers (i)

There are no such residues in this entry.

