



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 02:59 pm BST

PDB ID : 4Z78
Title : Weak TCR binding to an unstable insulin epitope drives type 1 diabetes
Authors : Rizkallah, P.J.; Cole, D.K.
Deposited on : 2015-04-06
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	 81% 17%
1	D	277	 83% 14%
1	G	277	 83% 14%
2	B	100	 85% 13%
2	E	100	 85% 14%
2	H	100	 87% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	10	<p>70% 30% 30% 30% 10%</p>
3	F	10	<p>70% 40% 20% 20% 20%</p>
3	I	10	<p>70% 30% 40% 10% 20%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO4	A	308	-	-	-	X
6	SO4	D	304	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-D alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	277	Total	C	N	O	S	0	0	0
			2287	1449	406	424	8			
1	D	277	Total	C	N	O	S	0	0	0
			2287	1449	406	424	8			
1	G	277	Total	C	N	O	S	0	1	0
			2298	1455	410	425	8			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P01902
A	114	HIS	GLN	conflict	UNP P01902
A	276	PRO	-	expression tag	UNP P01902
D	0	MET	-	initiating methionine	UNP P01902
D	114	HIS	GLN	conflict	UNP P01902
D	276	PRO	-	expression tag	UNP P01902
G	0	MET	-	initiating methionine	UNP P01902
G	114	HIS	GLN	conflict	UNP P01902
G	276	PRO	-	expression tag	UNP P01902

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	E	100	Total	C	N	O	S	0	1	0
			844	538	142	160	4			
2	H	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	initiating methionine	UNP P61769
H	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Insulin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	S	0	0	0
			81	53	13	14	1			
3	F	10	Total	C	N	O	S	0	0	0
			81	53	13	14	1			
3	I	10	Total	C	N	O	S	0	0	0
			81	53	13	14	1			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	H	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0
5	E	1	Total C O 6 3 3	0	0
5	H	1	Total C O 6 3 3	0	0
5	H	1	Total C O 6 3 3	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	G	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	100	Total	O	0	0
			100	100		
7	B	46	Total	O	0	0
			46	46		
7	D	87	Total	O	0	0
			87	87		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	E	27	Total O 27 27	0	0
7	G	63	Total O 63 63	0	0
7	H	41	Total O 41 41	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, K-D alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, K-D alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, K-D alpha chain




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin




- Molecule 3: Insulin



- Molecule 3: Insulin



- Molecule 3: Insulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.17Å 151.57Å 182.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.09 – 2.30 78.09 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (78.09-2.30) 100.0 (78.09-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.188 , 0.233 0.193 , 0.235	Depositor DCC
R_{free} test set	2925 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10098	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.7426e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	1/2357 (0.0%)	0.91	4/3205 (0.1%)
1	D	0.83	0/2357	0.90	6/3205 (0.2%)
1	G	0.80	0/2368	0.94	7/3219 (0.2%)
2	B	0.73	0/860	0.80	0/1162
2	E	0.63	0/867	0.77	0/1172
2	H	0.72	0/860	0.79	0/1162
3	C	0.91	0/82	1.29	0/107
3	F	1.03	0/82	1.28	1/107 (0.9%)
3	I	0.73	0/82	1.32	1/107 (0.9%)
All	All	0.80	1/9915 (0.0%)	0.89	19/13446 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
3	C	0	1
3	I	0	2
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	GLU	CD-OE1	5.10	1.31	1.25

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	121	ARG	NE-CZ-NH2	-10.28	115.16	120.30
1	G	121	ARG	NE-CZ-NH1	8.94	124.77	120.30
1	G	202	ARG	NE-CZ-NH2	-7.21	116.70	120.30
1	A	121	ARG	NE-CZ-NH1	-7.06	116.77	120.30
3	I	9	GLY	N-CA-C	-6.71	96.32	113.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	HIS	Peptide
3	C	8	ARG	Peptide
3	I	4	VAL	Peptide
3	I	8	ARG	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2287	0	2153	38	0
1	D	2287	0	2153	36	0
1	G	2298	0	2165	26	0
2	B	837	0	803	9	0
2	E	844	0	811	6	0
2	H	837	0	803	8	0
3	C	81	0	81	18	0
3	F	81	0	81	24	0
3	I	81	0	81	14	0
4	A	12	0	18	1	0
4	D	4	0	6	0	0
4	H	4	0	6	0	0
5	A	12	0	16	4	0
5	D	6	0	8	0	0
5	E	6	0	8	1	0
5	H	12	0	16	2	0
6	A	20	0	0	0	0
6	D	20	0	0	0	0
6	G	5	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	100	0	0	2	0
7	B	46	0	0	0	0
7	D	87	0	0	1	1
7	E	27	0	0	0	0
7	G	63	0	0	1	1
7	H	41	0	0	2	0
All	All	10098	0	9209	147	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 147 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:5:CYS:HB3	3:C:8:ARG:HH11	1.02	1.14
3:C:5:CYS:HB3	3:C:8:ARG:NH1	1.61	1.13
3:C:5:CYS:CB	3:C:8:ARG:HH11	1.64	1.11
3:F:10:PHE:OXT	3:F:10:PHE:CD1	2.11	1.04
3:F:8:ARG:NH2	3:F:8:ARG:H	1.60	0.99

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:D:435:HOH:O	7:G:431:HOH:O[3_745]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	275/277 (99%)	271 (98%)	4 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	275/277 (99%)	270 (98%)	5 (2%)	0	100	100
1	G	276/277 (100%)	270 (98%)	6 (2%)	0	100	100
2	B	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	E	99/100 (99%)	96 (97%)	3 (3%)	0	100	100
2	H	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
3	C	8/10 (80%)	6 (75%)	1 (12%)	1 (12%)	0	0
3	F	8/10 (80%)	4 (50%)	2 (25%)	2 (25%)	0	0
3	I	8/10 (80%)	5 (62%)	2 (25%)	1 (12%)	0	0
All	All	1145/1161 (99%)	1116 (98%)	25 (2%)	4 (0%)	41	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	F	8	ARG
3	I	8	ARG
3	C	7	GLU
3	F	9	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/236 (100%)	214 (91%)	22 (9%)	9	10
1	D	236/236 (100%)	219 (93%)	17 (7%)	14	18
1	G	237/236 (100%)	217 (92%)	20 (8%)	11	13
2	B	95/95 (100%)	88 (93%)	7 (7%)	13	17
2	E	96/95 (101%)	87 (91%)	9 (9%)	8	10
2	H	95/95 (100%)	90 (95%)	5 (5%)	22	31
3	C	8/8 (100%)	4 (50%)	4 (50%)	0	0
3	F	8/8 (100%)	5 (62%)	3 (38%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	I	8/8 (100%)	6 (75%)	2 (25%)	0	0
All	All	1019/1017 (100%)	930 (91%)	89 (9%)	10	12

5 of 89 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	110	LEU
1	D	272	LEU
2	H	39	LEU
1	D	111	ARG
1	D	182	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	82	GLN
1	D	114	HIS
1	G	263	HIS
1	D	86	ASN
1	D	149	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	304	-	5,5,5	0.71	0	5,5,5	2.12	2 (40%)
4	EDO	A	303	-	3,3,3	0.33	0	2,2,2	0.43	0
6	SO4	A	306	-	4,4,4	0.45	0	6,6,6	0.90	0
5	GOL	A	305	-	5,5,5	0.79	0	5,5,5	1.64	1 (20%)
4	EDO	D	301	-	3,3,3	0.52	0	2,2,2	0.24	0
5	GOL	D	302	-	5,5,5	0.72	0	5,5,5	1.35	1 (20%)
4	EDO	H	301	-	3,3,3	0.21	0	2,2,2	1.00	0
5	GOL	E	101	-	5,5,5	0.96	0	5,5,5	1.30	1 (20%)
6	SO4	D	303	-	4,4,4	0.44	0	6,6,6	0.89	0
4	EDO	A	301	-	3,3,3	0.48	0	2,2,2	0.50	0
6	SO4	A	307	-	4,4,4	0.41	0	6,6,6	0.42	0
6	SO4	A	309	-	4,4,4	0.45	0	6,6,6	0.45	0
6	SO4	G	301	-	4,4,4	0.54	0	6,6,6	0.93	0
6	SO4	D	305	-	4,4,4	0.44	0	6,6,6	0.50	0
5	GOL	H	303	-	5,5,5	1.32	0	5,5,5	1.42	1 (20%)
4	EDO	A	302	-	3,3,3	1.27	0	2,2,2	1.15	0
6	SO4	D	304	-	4,4,4	0.51	0	6,6,6	0.46	0
6	SO4	A	308	-	4,4,4	0.36	0	6,6,6	0.53	0
6	SO4	D	306	-	4,4,4	0.56	0	6,6,6	0.59	0
5	GOL	H	302	-	5,5,5	0.81	0	5,5,5	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	304	-	-	2/4/4/4	-
4	EDO	A	303	-	-	0/1/1/1	-
5	GOL	A	305	-	-	4/4/4/4	-
4	EDO	D	301	-	-	0/1/1/1	-
5	GOL	D	302	-	-	0/4/4/4	-
4	EDO	H	301	-	-	1/1/1/1	-
5	GOL	E	101	-	-	3/4/4/4	-
4	EDO	A	301	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	H	303	-	-	1/4/4/4	-
4	EDO	A	302	-	-	0/1/1/1	-
5	GOL	H	302	-	-	4/4/4/4	-

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	304	GOL	O3-C3-C2	2.98	124.48	110.20
5	A	304	GOL	C3-C2-C1	2.55	121.64	111.70
5	E	101	GOL	O3-C3-C2	2.53	122.34	110.20
5	H	303	GOL	O1-C1-C2	2.36	121.51	110.20
5	A	305	GOL	O1-C1-C2	2.31	121.27	110.20

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	304	GOL	O1-C1-C2-C3
5	A	305	GOL	O1-C1-C2-C3
5	E	101	GOL	O1-C1-C2-C3
5	A	305	GOL	C1-C2-C3-O3
5	H	302	GOL	O1-C1-C2-C3

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	304	GOL	2	0
5	A	305	GOL	2	0
5	E	101	GOL	1	0
6	G	301	SO4	1	0
4	A	302	EDO	1	0
5	H	302	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/277 (100%)	-0.21	3 (1%) 80 85	17, 32, 55, 83	0
1	D	277/277 (100%)	-0.27	3 (1%) 80 85	18, 36, 62, 93	0
1	G	277/277 (100%)	-0.13	7 (2%) 57 64	23, 41, 72, 97	0
2	B	100/100 (100%)	-0.36	0 100 100	19, 35, 58, 68	0
2	E	100/100 (100%)	-0.16	1 (1%) 82 86	21, 44, 78, 95	0
2	H	100/100 (100%)	-0.38	0 100 100	23, 37, 56, 70	0
3	C	10/10 (100%)	4.90	7 (70%) 0 0	32, 45, 62, 64	7 (70%)
3	F	10/10 (100%)	5.93	7 (70%) 0 0	33, 50, 63, 69	7 (70%)
3	I	10/10 (100%)	4.41	7 (70%) 0 0	44, 58, 66, 70	7 (70%)
All	All	1161/1161 (100%)	-0.09	35 (3%) 50 57	17, 38, 68, 97	21 (1%)

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	6	GLY	14.3
3	F	5	CYS	11.6
3	F	4	VAL	9.7
3	C	6	GLY	9.7
3	F	10	PHE	8.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	H	303	6/6	0.26	0.32	53,58,69,69	0
6	SO4	A	308	5/5	0.71	0.63	84,94,108,113	0
4	EDO	A	302	4/4	0.76	0.23	39,51,52,53	0
6	SO4	D	306	5/5	0.76	0.38	71,88,103,110	0
6	SO4	D	304	5/5	0.79	0.42	70,85,91,94	0
5	GOL	D	302	6/6	0.81	0.25	46,51,57,59	0
5	GOL	H	302	6/6	0.81	0.40	32,43,48,60	0
5	GOL	E	101	6/6	0.82	0.16	44,52,56,59	0
6	SO4	G	301	5/5	0.85	0.37	65,74,99,112	0
6	SO4	A	309	5/5	0.86	0.39	87,90,97,107	0
5	GOL	A	305	6/6	0.87	0.34	38,47,50,50	0
6	SO4	A	307	5/5	0.87	0.42	78,90,101,104	0
5	GOL	A	304	6/6	0.88	0.19	34,38,40,47	0
4	EDO	D	301	4/4	0.89	0.15	52,55,55,58	0
6	SO4	A	306	5/5	0.91	0.29	51,57,79,83	0
6	SO4	D	305	5/5	0.93	0.32	75,82,93,101	0
4	EDO	H	301	4/4	0.95	0.18	35,36,38,47	0
4	EDO	A	301	4/4	0.95	0.14	46,46,48,48	0
6	SO4	D	303	5/5	0.96	0.19	48,57,72,75	0
4	EDO	A	303	4/4	0.96	0.16	39,39,41,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.