



# Full wwPDB X-ray Structure Validation Report i

May 26, 2020 – 04:06 pm BST

PDB ID : 4ZA8  
Title : Crystal structure of A niger Fdc1 in complex with penta-fluorocinnamic acid  
Authors : Payne, K.A.P.; Leys, D.  
Deposited on : 2015-04-13  
Resolution : 1.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 1.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1202 (1.10-1.02)
Clashscore	141614	1252 (1.10-1.02)
Ramachandran outliers	138981	1204 (1.10-1.02)
Sidechain outliers	138945	1202 (1.10-1.02)
RSRZ outliers	127900	1178 (1.10-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	508	87%	10% ..

## 2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 4665 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative uncharacterized protein An03g06590.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	500	4097	2593	706	768	30	0	33	1

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	499B	LEU	-	expression tag	UNP A2QHE5
A	499C	GLU	-	expression tag	UNP A2QHE5
A	502	HIS	-	expression tag	UNP A2QHE5
A	503	HIS	-	expression tag	UNP A2QHE5
A	504	HIS	-	expression tag	UNP A2QHE5
A	505	HIS	-	expression tag	UNP A2QHE5
A	506	HIS	-	expression tag	UNP A2QHE5
A	507	HIS	-	expression tag	UNP A2QHE5

- Molecule 2 is 1-deoxy-5-O-phosphono-1-(3,3,4,5-tetramethyl-9,11-dioxo-2,3,8,9,10,11-hexahydro-7H-quinolino[1,8-fg]pteridin-12-iium-7-yl)-D-ribitol (three-letter code: 4LU) (formula: C<sub>22</sub>H<sub>30</sub>N<sub>4</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	36	22	4	9	1	0	1

- Molecule 3 is 1-deoxy-5-O-phosphono-1-[(10aR)-2,2,3,4-tetramethyl-8,10-dioxo-1,2,8,9,10,10-a-hexahydro-6H-indeno[1,7-ef]pyrimido[4,5-b][1,4]diazepin-6-yl]-D-ribitol (three-letter code: FZZ) (formula: C<sub>22</sub>H<sub>29</sub>N<sub>4</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	36	22	4	9	1	0	1


- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mn 1 1	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total K 2 2	0	0

- Molecule 6 is penta-fluorocinnamic acid (three-letter code: F5C) (formula: C<sub>9</sub>H<sub>3</sub>F<sub>5</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C F O 16 9 5 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	477	Total O 477 477	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative uncharacterized protein An03g06590



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.01Å    63.93Å    87.84Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	64.81 – 1.06 64.81 – 1.06	Depositor EDS
% Data completeness (in resolution range)	100.0 (64.81-1.06) 99.8 (64.81-1.06)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.92 (at 1.06Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
$R$ , $R_{free}$	0.147 , 0.166 0.144 , 0.161	Depositor DCC
$R_{free}$ test set	12068 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	6.8	Xtriage
Anisotropy	1.889	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 59.3	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.48$ , $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4665	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FZZ, K, MN, F5C, 4LU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.38	32/4199 (0.8%)	1.24	37/5707 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	410	GLU	CD-OE1	11.11	1.37	1.25
1	A	377[A]	GLU	CD-OE1	10.04	1.36	1.25
1	A	377[B]	GLU	CD-OE1	10.04	1.36	1.25
1	A	410	GLU	CD-OE2	9.93	1.36	1.25
1	A	480[A]	SER	CB-OG	8.36	1.53	1.42
1	A	480[B]	SER	CB-OG	8.36	1.53	1.42
1	A	380	ARG	CA-CB	8.18	1.72	1.53
1	A	65	ARG	NE-CZ	8.00	1.43	1.33
1	A	389	ASN	CG-ND2	-7.39	1.14	1.32
1	A	202	GLU	CD-OE2	6.94	1.33	1.25
1	A	65	ARG	CZ-NH1	-6.71	1.24	1.33
1	A	337	LYS	CE-NZ	-6.69	1.32	1.49
1	A	5	PRO	CB-CG	-6.64	1.16	1.50
1	A	282	GLU	CD-OE2	6.44	1.32	1.25
1	A	486	LYS	CD-CE	-6.36	1.35	1.51
1	A	84[A]	LEU	CA-CB	-6.31	1.39	1.53
1	A	84[B]	LEU	CA-CB	-6.31	1.39	1.53
1	A	18[A]	LYS	CD-CE	6.13	1.66	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	18[B]	LYS	CD-CE	6.13	1.66	1.51
1	A	202	GLU	CD-OE1	6.03	1.32	1.25
1	A	336	ARG	CZ-NH1	-5.82	1.25	1.33
1	A	370	ARG	CZ-NH2	5.79	1.40	1.33
1	A	473	GLU	CA-CB	5.73	1.66	1.53
1	A	282	GLU	CD-OE1	5.63	1.31	1.25
1	A	481	TYR	CE2-CZ	5.50	1.45	1.38
1	A	398	ARG	CZ-NH1	-5.36	1.26	1.33
1	A	243	SER	CA-CB	5.35	1.60	1.52
1	A	18[A]	LYS	CE-NZ	5.29	1.62	1.49
1	A	18[B]	LYS	CE-NZ	5.29	1.62	1.49
1	A	83	ARG	CB-CG	-5.20	1.38	1.52
1	A	15[A]	GLU	CB-CG	5.01	1.61	1.52
1	A	15[B]	GLU	CB-CG	5.01	1.61	1.52

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	398	ARG	NE-CZ-NH2	-13.27	113.67	120.30
1	A	398	ARG	NE-CZ-NH1	12.82	126.71	120.30
1	A	404	ASP	CB-CG-OD2	11.39	128.55	118.30
1	A	65	ARG	NE-CZ-NH2	-9.58	115.51	120.30
1	A	365	ASP	CB-CG-OD1	9.30	126.67	118.30
1	A	306	ARG	NE-CZ-NH1	8.66	124.63	120.30
1	A	380	ARG	CG-CD-NE	8.54	129.75	111.80
1	A	435	ARG	NE-CZ-NH2	-7.70	116.45	120.30
1	A	306	ARG	NE-CZ-NH2	-7.41	116.60	120.30
1	A	380	ARG	NE-CZ-NH1	6.90	123.75	120.30
1	A	382	ARG	NE-CZ-NH2	-6.87	116.86	120.30
1	A	404	ASP	CB-CG-OD1	-6.74	112.23	118.30
1	A	254	ASP	CB-CG-OD1	6.45	124.10	118.30
1	A	286	TYR	CB-CG-CD2	-6.34	117.20	121.00
1	A	83	ARG	CB-CA-C	6.20	122.81	110.40
1	A	97	ARG	NE-CZ-NH2	-6.01	117.29	120.30
1	A	337	LYS	CD-CE-NZ	6.01	125.52	111.70
1	A	298	TYR	CG-CD2-CE2	6.00	126.10	121.30
1	A	382	ARG	CG-CD-NE	-5.99	99.23	111.80
1	A	302	ARG	NE-CZ-NH2	-5.83	117.38	120.30
1	A	86	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	65	ARG	NH1-CZ-NH2	5.82	125.80	119.40
1	A	481	TYR	CD1-CE1-CZ	5.72	124.95	119.80
1	A	156	MET	N-CA-CB	-5.57	100.58	110.60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	GLU	OE1-CD-OE2	5.44	129.83	123.30
1	A	128	ASP	CB-CG-OD2	-5.32	113.52	118.30
1	A	286	TYR	CB-CG-CD1	5.32	124.19	121.00
1	A	377[A]	GLU	CG-CD-OE2	-5.27	107.76	118.30
1	A	377[B]	GLU	CG-CD-OE2	-5.27	107.76	118.30
1	A	473	GLU	CG-CD-OE1	5.26	128.82	118.30
1	A	86	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	A	407	ASP	CB-CG-OD1	5.17	122.95	118.30
1	A	185[A]	LEU	CB-CG-CD2	5.17	119.78	111.00
1	A	185[B]	LEU	CB-CG-CD2	5.17	119.78	111.00
1	A	41	ARG	NE-CZ-NH1	-5.14	117.73	120.30
1	A	404	ASP	CB-CA-C	-5.09	100.22	110.40
1	A	452	ARG	CG-CD-NE	-5.03	101.24	111.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	380	ARG	Sidechain
1	A	83	ARG	Sidechain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4097	0	3971	21	1
2	A	36	0	22	4	0
3	A	36	0	19	3	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
6	A	16	0	2	2	0
7	A	477	0	0	10	3
All	All	4665	0	4014	26	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:601[A]:4LU:H14	3:A:602[A]:FZZ:H5	1.51	0.93
2:A:601[A]:4LU:H13	2:A:601[A]:4LU:H14	1.54	0.89
2:A:601[A]:4LU:H14	3:A:602[A]:FZZ:C7	2.08	0.82
1:A:59[A]:MET:SD	7:A:881:HOH:O	2.42	0.77
1:A:59[A]:MET:CE	7:A:881:HOH:O	2.35	0.75
1:A:484[A]:ASP:OD1	7:A:701:HOH:O	2.13	0.66
1:A:59[A]:MET:HE3	7:A:881:HOH:O	1.96	0.63
1:A:196[B]:HIS:HE1	1:A:270:SER:O	1.84	0.61
1:A:76[B]:SER:HB3	7:A:930:HOH:O	2.02	0.58
6:A:606:F5C:FD2	6:A:606:F5C:H1	1.93	0.57
1:A:282:GLU:CD	1:A:439[B]:LEU:HD13	2.26	0.56
1:A:5:PRO:HG2	1:A:8:LEU:HD12	1.88	0.56
1:A:117:VAL:HG23	1:A:248[B]:VAL:HG22	1.90	0.54
1:A:280:PHE:O	1:A:282:GLU:HG3	2.08	0.53
1:A:445:HIS:HE1	7:A:807:HOH:O	1.92	0.53
1:A:439[A]:LEU:CD2	6:A:606:F5C:CB	2.87	0.51
1:A:129[A]:ASP:O	1:A:129[A]:ASP:CG	2.49	0.49
1:A:496:MET:O	7:A:702:HOH:O	2.20	0.48
1:A:196[B]:HIS:CE1	1:A:270:SER:O	2.65	0.48
1:A:363:ARG:NH1	7:A:708:HOH:O	2.47	0.47
1:A:377[B]:GLU:HG3	7:A:713:HOH:O	2.13	0.46
1:A:196[A]:HIS:HE1	1:A:270:SER:O	1.99	0.46
1:A:59[A]:MET:HG2	7:A:826:HOH:O	2.16	0.45
2:A:601[A]:4LU:H7	3:A:602[A]:FZZ:H20	1.84	0.42
1:A:108[A]:ASP:OD1	1:A:108[A]:ASP:C	2.58	0.41
1:A:319:LEU:HA	1:A:320:THR:HA	1.91	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:795:HOH:O	7:A:795:HOH:O[2_655]	1.11	1.09
7:A:753:HOH:O	7:A:920:HOH:O[2_655]	1.61	0.59
1:A:502:HIS:ND1	7:A:948:HOH:O[3_545]	1.99	0.21

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	529/508 (104%)	514 (97%)	15 (3%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	441/430 (103%)	439 (100%)	2 (0%)	88 66

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	480[A]	SER
1	A	480[B]	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	445	HIS

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	4LU	A	601[A]	5,4	32,39,39	2.44	11 (34%)	41,62,62	4.19	16 (39%)
3	FZZ	A	602[A]	5,4	31,39,39	4.48	12 (38%)	32,62,62	1.76	5 (15%)
6	F5C	A	606	-	13,16,16	3.05	6 (46%)	20,23,23	2.58	11 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4LU	A	601[A]	5,4	-	3/18/30/30	0/3/4/4
3	FZZ	A	602[A]	5,4	-	2/16/29/29	0/3/4/4
6	F5C	A	606	-	-	0/3/5/5	0/1/1/1

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602[A]	FZZ	C3-C17	14.64	1.53	1.38
3	A	602[A]	FZZ	C5-C4	13.20	1.60	1.50
3	A	602[A]	FZZ	C15-C16	-9.09	1.34	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602[A]	FZZ	C18-N4	-7.49	1.40	1.48
2	A	601[A]	4LU	C3-C5	-6.13	1.44	1.54
6	A	606	F5C	CG-CD2	-5.56	1.31	1.39
2	A	601[A]	4LU	C9A-N10	4.84	1.45	1.38
6	A	606	F5C	FE1-CE1	4.84	1.42	1.35
2	A	601[A]	4LU	C4-N3	4.82	1.41	1.33
2	A	601[A]	4LU	C1'-N10	-4.70	1.43	1.48
3	A	602[A]	FZZ	C5-C6	-4.03	1.51	1.56
6	A	606	F5C	FD1-CD1	-4.01	1.28	1.35
6	A	606	F5C	CB-CA	4.01	1.51	1.31
6	A	606	F5C	CE1-CD1	-3.50	1.31	1.37
3	A	602[A]	FZZ	P1-O7	-3.39	1.41	1.54
2	A	601[A]	4LU	C6-C5A	3.28	1.48	1.43
3	A	602[A]	FZZ	O3-C19	-3.21	1.36	1.43
2	A	601[A]	4LU	C13-C5	3.18	1.62	1.53
3	A	602[A]	FZZ	C22-C21	-3.07	1.47	1.51
3	A	602[A]	FZZ	O1-C1	-3.05	1.16	1.24
2	A	601[A]	4LU	C4-C4A	-3.05	1.36	1.41
2	A	601[A]	4LU	C9-C8	3.02	1.45	1.37
2	A	601[A]	4LU	O2'-C2'	3.00	1.49	1.43
6	A	606	F5C	CG-CD1	2.78	1.43	1.39
3	A	602[A]	FZZ	C14-C12	2.71	1.44	1.37
2	A	601[A]	4LU	C6-C7	2.62	1.49	1.40
3	A	602[A]	FZZ	O6-C22	2.41	1.54	1.44
2	A	601[A]	4LU	C8M-C8	-2.37	1.46	1.51
3	A	602[A]	FZZ	C9-C15	-2.20	1.37	1.41

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601[A]	4LU	C4-C4A-C10	14.76	129.72	119.95
2	A	601[A]	4LU	C4-N3-C2	11.75	125.06	115.14
2	A	601[A]	4LU	C4A-C4-N3	-10.08	109.64	123.43
2	A	601[A]	4LU	C12-C5-C6	8.56	120.10	111.72
2	A	601[A]	4LU	C13-C5-C6	-6.58	105.28	111.72
2	A	601[A]	4LU	C4A-C10-N10	5.59	126.05	120.30
2	A	601[A]	4LU	C7-C6-C5A	-5.57	113.52	119.06
6	A	606	F5C	CG-CB-CA	-4.91	111.56	127.03
3	A	602[A]	FZZ	C3-C1-N1	-4.87	116.77	123.43
6	A	606	F5C	FE2-CE2-CD2	4.72	129.12	119.27
3	A	602[A]	FZZ	C1-N1-C2	3.84	123.97	115.56
6	A	606	F5C	FE2-CE2-CZ	-3.82	111.31	119.27

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601[A]	4LU	C1'-N10-C10	3.69	121.72	118.41
6	A	606	F5C	C-CA-CB	-3.61	115.98	123.69
3	A	602[A]	FZZ	O3-C19-C20	3.60	117.84	109.10
6	A	606	F5C	CE2-CZ-CE1	-3.56	115.58	119.53
6	A	606	F5C	CD2-CG-CD1	-3.03	111.81	115.15
2	A	601[A]	4LU	C12-C5-C13	-2.84	102.30	108.67
6	A	606	F5C	CZ-CE1-CD1	2.68	122.51	119.53
2	A	601[A]	4LU	C3-C5-C6	2.67	113.48	107.38
6	A	606	F5C	CG-CD2-CE2	2.58	125.56	122.64
3	A	602[A]	FZZ	O3-C19-C18	2.50	115.61	109.59
2	A	601[A]	4LU	C4A-N5-C5A	-2.44	119.22	120.99
3	A	602[A]	FZZ	C18-C19-C20	2.41	116.53	109.79
6	A	606	F5C	FD2-CD2-CE2	-2.41	114.25	119.27
2	A	601[A]	4LU	C9-C8-C7	2.39	122.29	119.87
6	A	606	F5C	FZ-CZ-CE1	2.29	124.06	119.27
2	A	601[A]	4LU	O2'-C2'-C1'	-2.24	104.19	109.59
2	A	601[A]	4LU	C9A-N10-C10	-2.11	119.14	121.91
2	A	601[A]	4LU	C8M-C8-C9	-2.10	115.31	120.34
2	A	601[A]	4LU	C5A-C9A-N10	-2.10	118.75	120.43
6	A	606	F5C	CD2-CG-CB	2.06	130.32	122.57

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601[A]	4LU	C2'-C1'-N10-C9A
2	A	601[A]	4LU	C2'-C3'-C4'-C5'
3	A	602[A]	FZZ	C22-O6-P1-O7
3	A	602[A]	FZZ	C22-O6-P1-O9
2	A	601[A]	4LU	C4'-C5'-O5'-P


There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601[A]	4LU	4	0
3	A	602[A]	FZZ	3	0
6	A	606	F5C	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	500/508 (98%)	-0.36	5 (1%) 82 77	8, 11, 21, 30	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	503	HIS	5.2
1	A	502	HIS	4.4
1	A	505	HIS	3.5
1	A	504	HIS	2.9
1	A	293	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no carbohydrates in this entry.

### 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	F5C	A	606	16/16	0.87	0.19	11,15,19,19	16
2	4LU	A	601[A]	36/36	0.98	0.08	7,8,13,16	36


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	FZZ	A	602[A]	36/36	0.98	0.08	7,9,14,15	36
5	K	A	604	1/1	1.00	0.05	8,8,8,8	0
4	MN	A	603	1/1	1.00	0.05	7,7,7,7	0
5	K	A	605	1/1	1.00	0.05	8,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.