



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2024 – 11:27 AM JST

PDB ID : 5ZE0
Title : Hairpin Forming Complex, RAG1/2-Nicked(with Dideoxy) 12RSS/23RSS complex in Mg²⁺
Authors : Kim, M.S.; Chuenchor, W.; Chen, X.; Gellert, M.; Yang, W.
Deposited on : 2018-02-25
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	627	 0% (upper red bar), 75%, 22%, ..
1	C	627	 2%, 78%, 21%, ..
2	B	389	 0%, 63%, 24%, •, 13%
2	D	389	 4%, 55%, 30%, •, 13%
3	N	163	 12%, 62%, 9%, •, 28%
4	F	45	 36%, 58%, 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	I	16	
6	J	16	
7	G	54	
8	L	30	
9	M	39	

2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 20394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mouse RAG1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	618	Total	C	N	O	S	0	0	0
			4964	3124	883	923	34			
1	C	623	Total	C	N	O	S	0	0	0
			5010	3152	897	927	34			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	382	GLY	-	cloning artifact	UNP P15919
A	383	PRO	-	cloning artifact	UNP P15919
C	382	GLY	-	cloning artifact	UNP P15919
C	383	PRO	-	cloning artifact	UNP P15919

- Molecule 2 is a protein called mouse RAG2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	340	Total	C	N	O	S	0	0	0
			2650	1694	448	490	18			
2	D	340	Total	C	N	O	S	0	1	0
			2659	1700	451	490	18			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	cloning artifact	UNP P21784
B	0	PRO	-	cloning artifact	UNP P21784
B	1	VAL	MET	engineered mutation	UNP P21784
D	-1	GLY	-	cloning artifact	UNP P21784
D	0	PRO	-	cloning artifact	UNP P21784
D	1	VAL	MET	engineered mutation	UNP P21784

- Molecule 3 is a protein called HMGB1 A-B box.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	N	118	833	529	142	155	7	0	0	0

- Molecule 4 is a DNA chain called DNA (45-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	F	45	928	443	169	272	44	0	0	0

- Molecule 5 is a DNA chain called DNA (5'-D(*AP*AP*TP*CP*TP*GP*GP*CP*CP*TP*GP*TP*CP*TP*TP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	I	16	321	156	54	96	15	0	0	0

- Molecule 6 is a DNA chain called DNA (5'-D(*TP*AP*TP*CP*TP*GP*GP*CP*CP*TP*GP*TP*CP*TP*TP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
6	J	16	320	156	51	98	15	0	0	0

- Molecule 7 is a DNA chain called DNA (54-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
7	G	54	1106	529	191	332	54	0	0	0

- Molecule 8 is a DNA chain called DNA (30-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
8	L	30	611	290	118	173	30	0	0	0

- Molecule 9 is a DNA chain called DNA (39-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
9	M	39	805	381	162	223	39	0	0	0


- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total Mg 1 1	0	0
10	C	1	Total Mg 1 1	0	0

- Molecule 11 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	1	Total Zn 1 1	0	0
11	C	1	Total Zn 1 1	0	0

- Molecule 12 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	1	Total C O 4 2 2	0	0
12	A	1	Total C O 4 2 2	0	0
12	C	1	Total C O 4 2 2	0	0

- Molecule 13 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	A	1	Total K 1 1	0	0
13	C	1	Total K 1 1	0	0


- Molecule 14 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	A	50	Total O 50 50	0	0
14	B	8	Total O 8 8	0	0
14	C	47	Total O 47 47	0	0
14	D	9	Total O 9 9	0	0
14	F	16	Total O 16 16	0	0
14	I	4	Total O 4 4	0	0
14	J	4	Total O 4 4	0	0
14	G	14	Total O 14 14	0	0
14	L	10	Total O 10 10	0	0
14	M	7	Total O 7 7	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: mouse RAG1





- Molecule 1: mouse RAG1










- Molecule 2: mouse RAG2




- Molecule 2: mouse RAG2



- Molecule 3: HMGB1 A-B box






- Molecule 4: DNA (45-MER)




- Molecule 5: DNA (5'-D(*AP*AP*TP*CP*TP*GP*GP*CP*CP*TP*GP*TP*CP*TP*TP*A)-3')




- Molecule 6: DNA (5'-D(*TP*AP*TP*CP*TP*GP*GP*CP*CP*TP*GP*TP*CP*TP*TP*A)-3')




- Molecule 7: DNA (54-MER)



- Molecule 8: DNA (30-MER)



- Molecule 9: DNA (39-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	132.44Å 109.03Å 156.85Å 90.00° 114.43° 90.00°	Depositor
Resolution (Å)	34.80 – 2.75 86.66 – 2.75	Depositor EDS
% Data completeness (in resolution range)	97.4 (34.80-2.75) 89.3 (86.66-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 2.73Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.199 , 0.234 0.199 , 0.231	Depositor DCC
R_{free} test set	5086 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtrriage
Anisotropy	0.463	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	20394	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, K, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	1/5067 (0.0%)	0.68	0/6826
1	C	0.56	1/5114 (0.0%)	0.68	1/6887 (0.0%)
2	B	0.47	0/2717	0.62	0/3686
2	D	0.45	0/2727	0.62	0/3700
3	N	0.37	0/847	0.45	0/1135
4	F	1.29	5/1041 (0.5%)	1.34	14/1608 (0.9%)
5	I	1.07	1/358 (0.3%)	1.17	2/550 (0.4%)
6	J	1.22	1/356 (0.3%)	1.20	3/547 (0.5%)
7	G	1.06	0/1237	1.20	5/1908 (0.3%)
8	L	1.25	3/686 (0.4%)	1.17	4/1052 (0.4%)
9	M	1.03	2/907 (0.2%)	1.04	3/1395 (0.2%)
All	All	0.71	14/21057 (0.1%)	0.82	32/29294 (0.1%)

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	L	17	DC	OP3-P	-11.24	1.47	1.61
9	M	17	DC	OP3-P	-10.17	1.49	1.61
9	M	21	DG	C3'-O3'	-7.34	1.34	1.44
4	F	27	DT	C3'-O3'	-7.28	1.34	1.44
4	F	36	DC	O5'-C5'	-7.13	1.24	1.42

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	36	DC	O5'-P-OP1	-13.32	93.71	105.70
4	F	28	DG	O5'-P-OP2	-8.98	97.62	105.70
7	G	41	DG	O5'-P-OP1	-8.50	98.05	105.70
8	L	32	DC	O4'-C4'-C3'	-7.82	101.31	106.00
9	M	17	DC	O5'-P-OP1	-7.69	98.78	105.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4964	0	4933	103	0
1	C	5010	0	4989	96	0
2	B	2650	0	2592	68	0
2	D	2659	0	2595	125	0
3	N	833	0	703	9	0
4	F	928	0	511	21	0
5	I	321	0	182	7	0
6	J	320	0	183	6	0
7	G	1106	0	614	19	0
8	L	611	0	335	11	0
9	M	805	0	435	14	0
10	A	1	0	0	0	0
10	C	1	0	0	0	0
11	A	1	0	0	0	0
11	C	1	0	0	0	0
12	A	8	0	12	1	0
12	C	4	0	6	0	0
13	A	1	0	0	0	0
13	C	1	0	0	0	0
14	A	50	0	0	5	0
14	B	8	0	0	1	0
14	C	47	0	0	1	0
14	D	9	0	0	0	0
14	F	16	0	0	0	0
14	G	14	0	0	0	0
14	I	4	0	0	0	0
14	J	4	0	0	0	0
14	L	10	0	0	0	0
14	M	7	0	0	0	0
All	All	20394	0	18090	443	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 443 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:741:LEU:HG	1:A:917:ALA:HB1	1.53	0.91
1:C:848:ARG:HG2	1:C:848:ARG:HH11	1.34	0.90
2:D:263:LEU:HD21	2:D:271:PHE:HD2	1.38	0.88
2:D:95:GLY:HA2	2:D:104:SER:O	1.73	0.87
2:D:233:LEU:HD21	2:D:297:ILE:HD11	1.58	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	616/627 (98%)	596 (97%)	20 (3%)	0	100	100
1	C	621/627 (99%)	598 (96%)	23 (4%)	0	100	100
2	B	334/389 (86%)	322 (96%)	12 (4%)	0	100	100
2	D	335/389 (86%)	323 (96%)	12 (4%)	0	100	100
3	N	108/163 (66%)	107 (99%)	1 (1%)	0	100	100
All	All	2014/2195 (92%)	1946 (97%)	68 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	542/550 (98%)	526 (97%)	16 (3%)	41	61
1	C	547/550 (100%)	530 (97%)	17 (3%)	40	60
2	B	295/344 (86%)	286 (97%)	9 (3%)	40	60
2	D	295/344 (86%)	276 (94%)	19 (6%)	17	31
3	N	68/139 (49%)	63 (93%)	5 (7%)	13	24
All	All	1747/1927 (91%)	1681 (96%)	66 (4%)	33	53

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	294	ASP
2	D	310	ASP
3	N	106	CYS
2	B	317	TRP
2	B	268	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	278	GLN
2	B	282	GLN
2	D	53	ASN
2	D	94	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	EDO	C	2004	-	3,3,3	0.48	0	2,2,2	0.23	0
12	EDO	A	2004	-	3,3,3	0.47	0	2,2,2	0.12	0
12	EDO	A	2003	-	3,3,3	0.50	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	EDO	C	2004	-	-	0/1/1/1	-
12	EDO	A	2004	-	-	0/1/1/1	-
12	EDO	A	2003	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	A	2003	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	618/627 (98%)	0.31	6 (0%) 82 87	37, 56, 90, 143	0
1	C	623/627 (99%)	0.30	11 (1%) 68 76	34, 57, 89, 121	0
2	B	340/389 (87%)	0.27	5 (1%) 73 81	46, 77, 109, 124	0
2	D	340/389 (87%)	0.49	15 (4%) 34 41	40, 70, 110, 130	0
3	N	118/163 (72%)	0.73	20 (16%) 1 1	104, 148, 162, 167	0
4	F	45/45 (100%)	-0.19	0 100 100	37, 65, 112, 129	0
5	I	16/16 (100%)	-0.03	1 (6%) 20 24	42, 92, 125, 125	0
6	J	16/16 (100%)	0.32	2 (12%) 3 4	43, 89, 130, 130	0
7	G	54/54 (100%)	0.26	3 (5%) 24 29	40, 107, 167, 192	0
8	L	30/30 (100%)	-0.12	0 100 100	40, 66, 111, 115	0
9	M	39/39 (100%)	0.15	1 (2%) 56 65	47, 104, 162, 174	0
All	All	2239/2395 (93%)	0.33	64 (2%) 51 61	34, 64, 134, 192	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	N	113	ILE	6.5
3	N	13	MET	4.8
1	A	408	LEU	4.6
3	N	11	GLY	4.4
3	N	144	TYR	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
10	MG	C	2001	1/1	0.80	0.17	54,54,54,54	0
10	MG	A	2001	1/1	0.82	0.44	61,61,61,61	0
12	EDO	A	2003	4/4	0.85	0.21	60,61,72,74	0
12	EDO	C	2004	4/4	0.86	0.35	68,73,74,81	0
12	EDO	A	2004	4/4	0.92	0.26	51,52,54,56	0
13	K	A	2005	1/1	0.97	0.10	49,49,49,49	0
13	K	C	2003	1/1	0.97	0.13	42,42,42,42	0
11	ZN	C	2002	1/1	0.99	0.21	54,54,54,54	0
11	ZN	A	2002	1/1	0.99	0.21	49,49,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.