

wwPDB NMR Structure Validation Summary Report (i)

Jun 6, 2023 – 05:48 pm BST

PDB ID : 3ZGK BMRB ID : 18910

Title: NMR solution structure of the RXLR effector AVR3a11 from Phytophthora

Capsici

Authors: Tolchard, J.; Chambers, V.S.; Boutemy, L.S.; Gathercole, R.L.; Banfield, M.J.;

Blumenschein, T.M.

Deposited on : 2012-12-18

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 77%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{array}{c} { m NMR \ archive} \ (\#{ m Entries}) \end{array}$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	72	62%	15%	22%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model			
1	A:71-A:111, A:118-A:132	0.96	3			
	(56)					

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models		
1	2, 3, 6, 8, 9, 12, 13, 14, 17, 20		
2	1, 4, 7, 11, 15, 16, 18, 19		
3	5, 10		



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1166 atoms, of which 585 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called AVR3A11.

Mol	Chain	Residues	Atoms			Trace			
1	Λ	79	Total	С	Н	N	О	S	0
1	А	12	1166	372	585	96	110	3	U

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	GLY	-	expression tag	UNP G1K3S4
A	62	PRO	-	expression tag	UNP G1K3S4
A	63	GLY	-	SEE REMARK 999	UNP G1K3S4
A	64	LEU	-	SEE REMARK 999	UNP G1K3S4
A	65	THR	-	SEE REMARK 999	UNP G1K3S4
A	66	ASP	-	SEE REMARK 999	UNP G1K3S4
A	67	LEU	-	SEE REMARK 999	UNP G1K3S4
A	68	PHE	-	SEE REMARK 999	UNP G1K3S4
A	69	LYS	_	SEE REMARK 999	UNP G1K3S4




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: AVR3A11



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

• Molecule 1: AVR3A11





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: DISTANCE GEOMETRY, SIMULATED ANNEALING.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: LOWEST ENERGY.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.3
NMRPipe	structure solution	
CcpNmr Analysis	structure solution	ANALYSIS
UNIO	structure solution	
CYANA	structure solution	
CNS USING RECOORD SCRIPTS	structure solution	SCRIPTS

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	696
Number of shifts mapped to atoms	696
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	77%



6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	В	Sond lengths	Bond angles		
MIOI	Chain	RMSZ	#Z>5	RMSZ	#Z>5	
1	A	0.69 ± 0.04	$0\pm0/472~(~0.0\pm~0.0\%)$	0.70 ± 0.04	$0\pm0/633~(~0.0\pm~0.0\%)$	
All	All	0.69	0/9440~(~0.0%)	0.70	1/12660 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	0.5 ± 0.6
All	All	0	10

There are no bond-length outliers.

All unique angle outliers are listed below.

	Mol	Chain	Ros	Type	Atoms	$egin{array}{c c} \mathbf{Z} & \mathbf{Observed}(^o) & \mathbf{Ideal}(^o) \end{array}$	$f{Z} { m Observed}(^o)$	Mod		
	WIOI	Chain	rtes	Type	Atoms			ruear()	Worst	Total
ĺ	1	A	129	ARG	NE-CZ-NH1	5.05	122.82	120.30	15	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	129	ARG	Sidechain	7
1	A	120	ARG	Sidechain	3

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	462	454	454	5±2
All	All	9240	9080	9080	108

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

5 of 66 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	$oxed{ ext{Models}}$	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:127:THR:HA	1:A:130:ASP:OD2	0.64	1.93	17	1
1:A:73:ALA:O	1:A:77:LYS:HG3	0.61	1.96	3	1
1:A:128:TYR:HA	1:A:131:TYR:CE2	0.59	2.32	6	1
1:A:95:LYS:HA	1:A:98:ASP:OD2	0.59	1.98	16	3
1:A:75:VAL:HB	1:A:96:TRP:CH2	0.58	2.33	17	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perc	entiles
1	A	55/72 (76%)	51±2 (93±4%)	3±2 (6±3%)	0±1 (1±1%)	24	71
All	All	1100/1440 (76%)	1025 (93%)	66 (6%)	9 (1%)	24	71

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	103	LEU	3
1	A	104	THR	3
1	A	118	TYR	1
1	A	111	LYS	1
1	A	71	GLU	1



6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Outliers	Pe	erce	entiles
1	A	47/60 (78%)	40±2 (84±3%)	7±2 (16±3%)		5	43
All	All	940/1200 (78%)	794 (84%)	146 (16%)		5	43

5 of 21 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	88	LYS	17
1	A	98	ASP	17
1	A	94	GLN	15
1	A	128	TYR	15
1	A	80	LYS	14

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.



6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 77% for the well-defined parts and 65% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	696
Number of shifts mapped to atoms	696
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

The following errors were found when reading this chemical shift list.

• Chemical shift has been reported more than once. First 5 (of 0) occurrences are reported below.

T:4 ID	Cl :	D	T	A 4		Shift Dat	a
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	64	LEU	HD22	0.95		1
1	A	64	LEU	HD23	0.95		1
1	A	64	LEU	HD12	0.95		1
1	A	64	LEU	HD13	0.95	•	1
1	A	65	THR	HG22	1.14		1
1	A	65	THR	HG23	1.14	•	1
1	A	73	ALA	HB2	1.48	0	1
1	A	73	ALA	HB3	1.48	0	1
1	A	74	ALA	HB2	1.52	0	1
1	A	74	ALA	HB3	1.52	0	1
1	A	75	VAL	HG22	0.28	0	2
1	A	75	VAL	HG23	0.28	0	2
1	A	75	VAL	HG12	0.29	0	2
1	A	75	VAL	HG13	0.29	0	2

Continued on next page...



 $Continued\ from\ previous\ page...$

Continue				A .		Shift Dat	a
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	79	ALA	HB2	1.61	0	1
1	A	79	ALA	HB3	1.61	0	1
1	A	81	ALA	HB2	1.55	0	1
1	A	81	ALA	HB3	1.55	0	1
1	A	82	ILE	HD12	0.49	0.04	1
1	A	82	ILE	HD13	0.49	0.04	1
1	A	82	ILE	HG22	0.78	0	1
1	A	82	ILE	HG23	0.78	0	1
1	A	84	ALA	HB2	1.48	0	1
1	A	84	ALA	HB3	1.48	0	1
1	A	89	ALA	HB2	1.2	0	1
1	A	89	ALA	HB3	1.2	0	1
1	A	92	VAL	HG12	0.69	0.01	2
1	A	92	VAL	HG13	0.69	0.01	2
1	A	92	VAL	HG22	-0.5	0	2
1	A	92	VAL	HG23	-0.5	0	2
1	A	97	ALA	HB2	1.81	0	1
1	A	97	ALA	HB3	1.81	0	1
1	A	103	LEU	HD12	0.72	0	2
1	A	103	LEU	HD13	0.72	0	2
1	A	103	LEU	HD22	0.86	0	2
1	A	103	LEU	HD23	0.86	0	2
1	A	104	THR	HG22	1.19	0	1
1	A	104	THR	HG23	1.19	0	1
1	A	106	LEU	HD22	0.37	0	2
1	A	106	LEU	HD23	0.37	0	2
1	A	106	LEU	HD12	-0.05	0	2
1	A	106	LEU	HD13	-0.05	0	2
1	A	110	LEU	HD12	0.81	0	1
1	A	110	LEU	HD13	0.81	0	1
1	A	110	LEU	HD22	0.81	0	1
1	A	110	LEU	HD23	0.81	0	1
1	A	121	VAL	HG22	1.38	0	2
1	A	121	VAL	HG23	1.38	0	2
1	A	121	VAL	HG12	1.51	0	2
1	A	121	VAL	HG13	1.51	0	2
1	A	132	VAL	HG22	0.83	•	1
1	A	132	VAL	HG23	0.83	•	1
1	A	132	VAL	HG12	0.83	•	1
1	A	132	VAL	HG13	0.83	•	1



7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	61	-0.49 ± 0.18	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	58	0.27 ± 0.11	None needed (< 0.5 ppm)
¹³ C′	0		None (insufficient data)
^{15}N	54	0.69 ± 0.28	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 77%, i.e. 597 atoms were assigned a chemical shift out of a possible 777. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}{ m H}$	$^{13}{ m C}$	$^{15}{ m N}$
Backbone	211/280 (75%)	105/113~(93%)	55/112~(49%)	51/55 (93%)
Sidechain	314/412 (76%)	$210/266 \ (79\%)$	101/129 (78%)	3/17 (18%)
Aromatic	72/85 (85%)	$36/39 \ (92\%)$	35/45~(78%)	1/1 (100%)
Overall	597/777 (77%)	351/418 (84%)	191/286 (67%)	55/73 (75%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	117	LYS	HG2	0.10	0.13 - 2.61	-5.1

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:







8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	842
Intra-residue (i-j =0)	140
Sequential (i-j =1)	202
Medium range ($ i-j >1$ and $ i-j <5$)	275
Long range (i-j ≥5)	225
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	11.7
Number of long range restraints per residue ¹	3.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	5.0	0.2
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Donton into topo o	Commit	07 1	Vic	${f Violated^3}$			Consistently Violated ⁴		
Restraints type	Count	$\%^1$	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$	
Intra-residue (i-j =0)	140	16.6	4	2.9	0.5	0	0.0	0.0	
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	120	14.3	4	3.3	0.5	0	0.0	0.0	
Sidechain-Sidechain	20	2.4	0	0.0	0.0	0	0.0	0.0	
Sequential (i-j =1)	202	24.0	11	5.4	1.3	0	0.0	0.0	
Backbone-Backbone	41	4.9	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	121	14.4	4	3.3	0.5	0	0.0	0.0	
Sidechain-Sidechain	40	4.8	7	17.5	0.8	0	0.0	0.0	
Medium range ($ i-j >1 \& i-j <5$)	275	32.7	20	7.3	2.4	0	0.0	0.0	
Backbone-Backbone	83	9.9	3	3.6	0.4	0	0.0	0.0	
Backbone-Sidechain	128	15.2	11	8.6	1.3	0	0.0	0.0	
Sidechain-Sidechain	64	7.6	6	9.4	0.7	0	0.0	0.0	
Long range ($ i-j \ge 5$)	225	26.7	12	5.3	1.4	0	0.0	0.0	
Backbone-Backbone	5	0.6	1	20.0	0.1	0	0.0	0.0	
Backbone-Sidechain	82	9.7	7	8.5	0.8	0	0.0	0.0	
Sidechain-Sidechain	138	16.4	4	2.9	0.5	0	0.0	0.0	
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0	
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0	
Total	842	100.0	47	5.6	5.6	0	0.0	0.0	
Backbone-Backbone	129	15.3	4	3.1	0.5	0	0.0	0.0	
Backbone-Sidechain	451	53.6	26	5.8	3.1	0	0.0	0.0	
Sidechain-Sidechain	262	31.1	17	6.5	2.0	0	0.0	0.0	

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	ations	5	M (8)	N/ (Å)	\mathbf{SD}^6 (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	$_{ m al} \mid { m Mean} ({ m \AA}) \mid$	Max (Å)	SD'(A)	Median (Å)
1	0	1	3	2	0	6	0.14	0.15	0.02	0.14
2	0	1	2	3	0	6	0.15	0.2	0.03	0.14
3	1	0	2	2	0	5	0.14	0.17	0.02	0.13
4	0	0	5	1	0	6	0.13	0.15	0.02	0.12
5	0	1	1	0	0	2	0.12	0.14	0.02	0.12
6	0	0	2	1	0	3	0.16	0.19	0.02	0.14
7	0	1	3	0	0	4	0.16	0.2	0.04	0.16
8	1	3	3	1	0	8	0.14	0.19	0.02	0.14
9	1	1	3	0	0	5	0.14	0.16	0.02	0.14
10	0	0	1	2	0	3	0.12	0.13	0.01	0.12
11	0	0	2	0	0	2	0.11	0.11	0.0	0.11

Continued on next page...




Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	SD(A)	Median (A)
12	0	1	1	0	0	2	0.16	0.2	0.04	0.16
13	0	3	3	1	0	7	0.14	0.19	0.02	0.13
14	0	0	1	0	0	1	0.11	0.11	0.0	0.11
15	1	3	4	1	0	9	0.15	0.19	0.03	0.16
16	1	1	4	1	0	7	0.14	0.17	0.02	0.12
17	2	3	2	1	0	8	0.14	0.2	0.03	0.13
18	2	3	3	0	0	8	0.14	0.17	0.02	0.15
19	0	1	1	1	0	3	0.13	0.18	0.03	0.11
20	0	1	2	2	0	5	0.15	0.19	0.02	0.14

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 795(IR:136, SQ:191, MR:255, LR:213, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	restra	aints	Fraction	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
2	4	11	9	0	26	1	5.0
1	4	3	0	0	8	2	10.0
0	1	0	2	0	3	3	15.0
0	1	3	1	0	5	4	20.0
1	1	1	0	0	3	5	25.0
0	0	0	0	0	0	6	30.0
0	0	2	0	0	2	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	\mathbf{SD}^1 (Å)	Median (Å)
(1,509)	1:A:94:GLN:HG2	1:A:98:ASP:H	7	0.16	0.03	0.16
(1,509)	1:A:94:GLN:HG3	1:A:98:ASP:H	7	0.16	0.03	0.16
(1,118)	1:A:99:LYS:HG2	1:A:101:TYR:H	7	0.15	0.03	0.15
(1,118)	1:A:99:LYS:HG3	1:A:101:TYR:H	7	0.15	0.03	0.15
(1,17)	1:A:95:LYS:HD2	1:A:97:ALA:H	5	0.17	0.02	0.17
(1,17)	1:A:95:LYS:HD3	1:A:97:ALA:H	5	0.17	0.02	0.17
(1,232)	1:A:76:LYS:HA	1:A:76:LYS:HD2	5	0.16	0.01	0.16
(1,232)	1:A:76:LYS:HA	1:A:76:LYS:HD3	5	0.16	0.01	0.16
(1,255)	1:A:71:GLU:HA	1:A:72:LYS:HG2	5	0.14	0.01	0.14
(1,255)	1:A:71:GLU:HA	1:A:72:LYS:HG3	5	0.14	0.01	0.14
(1,38)	1:A:94:GLN:HE21	1:A:96:TRP:H	4	0.15	0.03	0.16
(1,38)	1:A:94:GLN:HE22	1:A:96:TRP:H	4	0.15	0.03	0.16
(1,634)	1:A:95:LYS:HD2	1:A:99:LYS:HG2	4	0.14	0.02	0.15
(1,634)	1:A:95:LYS:HD2	1:A:99:LYS:HG3	4	0.14	0.02	0.15
(1,634)	1:A:95:LYS:HD3	1:A:99:LYS:HG2	4	0.14	0.02	0.15
(1,634)	1:A:95:LYS:HD3	1:A:99:LYS:HG3	4	0.14	0.02	0.15

Continued on next page...



Continued from previous page...


Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,538)	1:A:80:LYS:HB2	1:A:83:MET:HB2	4	0.14	0.02	0.15
(1,538)	1:A:80:LYS:HB2	1:A:83:MET:HB3	4	0.14	0.02	0.15
(1,538)	1:A:80:LYS:HB3	1:A:83:MET:HB2	4	0.14	0.02	0.15
(1,538)	1:A:80:LYS:HB3	1:A:83:MET:HB3	4	0.14	0.02	0.15
(1,538)	1:A:83:MET:HB2	1:A:120:ARG:HB2	4	0.14	0.02	0.15
(1,538)	1:A:83:MET:HB2	1:A:120:ARG:HB3	4	0.14	0.02	0.15
(1,538)	1:A:83:MET:HB3	1:A:120:ARG:HB2	4	0.14	0.02	0.15
(1,538)	1:A:83:MET:HB3	1:A:120:ARG:HB3	4	0.14	0.02	0.15
(1,43)	1:A:82:ILE:H	1:A:88:LYS:H	4	0.13	0.0	0.13
(1,43)	1:A:85:ASP:H	1:A:88:LYS:H	4	0.13	0.0	0.13
(1,552)	1:A:76:LYS:HB2	1:A:77:LYS:HD2	4	0.12	0.01	0.12

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table: All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,730)	1:A:110:LEU:HG	1:A:118:TYR:HD1	2	0.2
(1,730)	1:A:110:LEU:HG	1:A:118:TYR:HD2	2	0.2
(1,509)	1:A:94:GLN:HG2	1:A:98:ASP:H	7	0.2
(1,509)	1:A:94:GLN:HG3	1:A:98:ASP:H	7	0.2
(1,289)	1:A:96:TRP:HA	1:A:96:TRP:HD1	17	0.2
(1,286)	1:A:109:PHE:HA	1:A:111:LYS:H	7	0.2
(1,194)	1:A:92:VAL:HA	1:A:95:LYS:HD2	12	0.2
(1,194)	1:A:92:VAL:HA	1:A:95:LYS:HD3	12	0.2
(1,509)	1:A:94:GLN:HG2	1:A:98:ASP:H	13	0.19
(1,509)	1:A:94:GLN:HG3	1:A:98:ASP:H	13	0.19
(1,38)	1:A:94:GLN:HE21	1:A:96:TRP:H	6	0.19
(1,38)	1:A:94:GLN:HE22	1:A:96:TRP:H	6	0.19
(1,238)	1:A:80:LYS:HA	1:A:83:MET:HB2	20	0.19
(1,238)	1:A:80:LYS:HA	1:A:83:MET:HB3	20	0.19
(1,17)	1:A:95:LYS:HD2	1:A:97:ALA:H	8	0.19
(1,17)	1:A:95:LYS:HD3	1:A:97:ALA:H	8	0.19
(1,17)	1:A:95:LYS:HD2	1:A:97:ALA:H	15	0.19



10 Dihedral-angle violation analysis (i)

No dihedral-angle restraints found

