



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2024 – 08:23 AM EST

PDB ID : 2ZHL
Title : Crystal structure of human galectin-9 N-terminal CRD in complex with N-acetyllactosamine dimer (crystal 2)
Authors : Nagae, M.; Nishi, N.; Murata, T.; Usui, T.; Nakamura, T.; Wakatsuki, S.; Kato, R.
Deposited on : 2008-02-06
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	 90% 6%
1	B	148	 83% 12%
1	C	148	 90% 6%
1	D	148	 88% 8%
2	E	4	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	4	 100%
2	G	4	 100%
2	H	4	 50% 50%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galectin-9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	142	1115	714	191	205	5	0	0	0
1	B	142	1115	714	191	205	5	0	0	0
1	C	142	1115	714	191	205	5	0	0	0
1	D	142	1115	714	191	205	5	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	4	51	28	2	21	0	0	0
2	F	4	51	28	2	21	0	0	0
2	G	4	51	28	2	21	0	0	0
2	H	4	51	28	2	21	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	118	Total	O	0	0
			118	118		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	118	Total 118	O 118	0	0
3	C	122	Total 122	O 122	0	0
3	D	132	Total 132	O 132	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Galectin-9





- Molecule 1: Galectin-9



- Molecule 1: Galectin-9



- Molecule 1: Galectin-9



- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
GAL2
MAG3
GAL4

- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1
GAL2
MAG3
GAL4

- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  50% 50%

MAG1
GAL2
MAG3
GAL4

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	35.38Å 67.71Å 220.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.86 – 1.75 33.86 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.8 (33.86-1.75) 98.8 (33.86-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.185 , 0.230 0.185 , 0.230	Depositor DCC
R_{free} test set	2744 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtrriage
Anisotropy	0.552	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5154	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.3461e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1147	0.72	0/1554
1	B	0.69	0/1147	0.75	1/1554 (0.1%)
1	C	0.67	0/1147	0.73	1/1554 (0.1%)
1	D	0.71	0/1147	0.72	0/1554
All	All	0.68	0/4588	0.73	2/6216 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	18	SER	N-CA-C	5.57	126.04	111.00
1	C	101	LEU	CA-CB-CG	5.50	127.95	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	17	PHE	Peptide
1	B	17	PHE	Peptide
1	C	17	PHE	Peptide
1	D	17	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1115	0	1062	16	0
1	B	1115	0	1062	16	0
1	C	1115	0	1062	7	0
1	D	1115	0	1062	9	0
2	E	51	0	45	0	0
2	F	51	0	45	0	0
2	G	51	0	45	0	0
2	H	51	0	44	0	0
3	A	118	0	0	2	0
3	B	118	0	0	2	0
3	C	122	0	0	2	0
3	D	132	0	0	3	0
All	All	5154	0	4427	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:121:GLN:OE1	3:B:1053:HOH:O	1.62	1.15
1:A:66:PHE:CZ	1:A:91:MET:HE1	1.82	1.15
1:A:91:MET:HE3	1:A:93:PHE:HB2	1.10	1.06
1:A:91:MET:CE	1:A:93:PHE:HB2	1.86	1.05
1:A:91:MET:SD	3:A:1436:HOH:O	2.17	0.99
1:A:66:PHE:CZ	1:A:91:MET:CE	2.47	0.97
1:A:91:MET:HE3	1:A:93:PHE:CB	2.02	0.87
1:A:66:PHE:CE1	1:A:91:MET:CE	2.60	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:PHE:CE1	1:A:91:MET:HE2	2.17	0.80
1:D:106:GLN:OE1	3:D:1130:HOH:O	1.99	0.80
1:D:121:GLN:OE1	3:D:1213:HOH:O	2.05	0.75
1:D:121:GLN:NE2	3:D:1433:HOH:O	2.20	0.74
1:A:66:PHE:CE1	1:A:91:MET:HE1	2.22	0.72
1:A:90:HIS:HB2	3:C:1057:HOH:O	1.91	0.69
1:A:102:CYS:SG	3:A:1454:HOH:O	2.11	0.66
1:B:121:GLN:NE2	3:B:1434:HOH:O	2.27	0.66
1:B:63:ASN:ND2	1:B:65:ARG:HE	1.95	0.64
1:B:121:GLN:OE1	1:B:121:GLN:N	2.33	0.62
1:D:104:LEU:HD23	1:D:111:LYS:HE2	1.80	0.61
1:B:106:GLN:OE1	1:B:111:LYS:NZ	2.34	0.61
1:D:100:ASP:H	1:D:115:ASN:ND2	2.02	0.57
1:C:63:ASN:ND2	1:C:65:ARG:HE	2.01	0.57
1:D:7:GLN:NE2	1:D:22:GLN:H	2.04	0.56
1:D:63:ASN:ND2	1:D:65:ARG:HE	2.04	0.55
1:A:90:HIS:CD2	1:C:132:ASP:OD2	2.60	0.53
1:B:88:LYS:HD2	1:B:90:HIS:H	1.73	0.53
1:B:63:ASN:HD21	1:B:65:ARG:HE	1.55	0.53
1:B:88:LYS:HD2	1:B:89:THR:N	2.24	0.52
1:D:99:PHE:HA	1:D:115:ASN:HD21	1.74	0.51
1:A:63:ASN:ND2	1:A:65:ARG:HE	2.08	0.51
1:C:63:ASN:HD21	1:C:65:ARG:HE	1.58	0.51
1:B:7:GLN:NE2	1:B:22:GLN:H	2.10	0.48
1:B:88:LYS:HD2	1:B:89:THR:H	1.79	0.48
1:C:7:GLN:NE2	1:C:22:GLN:H	2.13	0.47
1:A:66:PHE:CZ	1:A:91:MET:HE2	2.41	0.46
1:C:7:GLN:HE22	1:C:22:GLN:H	1.63	0.46
1:A:63:ASN:HD21	1:A:65:ARG:HE	1.64	0.45
1:D:7:GLN:HE22	1:D:22:GLN:H	1.63	0.45
1:B:30:GLN:HG3	1:B:102:CYS:SG	2.58	0.44
1:B:88:LYS:HE2	1:B:90:HIS:HB2	2.00	0.42
1:B:48:ASN:HD22	1:B:61:HIS:HA	1.84	0.42
1:B:101:LEU:HG	1:B:103:PHE:CE2	2.54	0.42
1:A:69:GLY:HA2	1:C:54:SER:HB3	2.00	0.42
1:B:30:GLN:CG	1:B:102:CYS:SG	3.08	0.42
1:B:48:ASN:ND2	1:B:61:HIS:ND1	2.64	0.41
1:C:129:HIS:HD2	3:C:1195:HOH:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/148 (95%)	138 (99%)	2 (1%)	0	100	100
1	B	140/148 (95%)	138 (99%)	2 (1%)	0	100	100
1	C	140/148 (95%)	137 (98%)	3 (2%)	0	100	100
1	D	140/148 (95%)	136 (97%)	4 (3%)	0	100	100
All	All	560/592 (95%)	549 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/127 (97%)	123 (100%)	0	100	100
1	B	123/127 (97%)	121 (98%)	2 (2%)	62	45
1	C	123/127 (97%)	123 (100%)	0	100	100
1	D	123/127 (97%)	123 (100%)	0	100	100
All	All	492/508 (97%)	490 (100%)	2 (0%)	91	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	40	SER
1	B	88	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	48	ASN
1	A	56	ASN
1	A	63	ASN
1	B	7	GLN
1	B	48	ASN
1	B	63	ASN
1	B	79	ASN
1	B	94	GLN
1	B	129	HIS
1	C	7	GLN
1	C	48	ASN
1	C	56	ASN
1	C	63	ASN
1	C	129	HIS
1	D	7	GLN
1	D	48	ASN
1	D	63	ASN
1	D	106	GLN
1	D	115	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	E	1	2	15,15,15	1.08	1 (6%)	21,21,21	1.49	3 (14%)
2	GAL	E	2	2	11,11,12	1.31	1 (9%)	15,15,17	1.11	2 (13%)
2	NAG	E	3	2	14,14,15	0.57	0	17,19,21	1.22	2 (11%)
2	GAL	E	4	2	11,11,12	0.58	0	15,15,17	1.15	2 (13%)
2	NAG	F	1	2	15,15,15	0.87	1 (6%)	21,21,21	0.84	1 (4%)
2	GAL	F	2	2	11,11,12	1.21	1 (9%)	15,15,17	0.99	1 (6%)
2	NAG	F	3	2	14,14,15	0.42	0	17,19,21	1.32	2 (11%)
2	GAL	F	4	2	11,11,12	0.69	0	15,15,17	1.24	2 (13%)
2	NAG	G	1	2	15,15,15	1.15	1 (6%)	21,21,21	2.22	6 (28%)
2	GAL	G	2	2	11,11,12	1.17	1 (9%)	15,15,17	1.34	2 (13%)
2	NAG	G	3	2	14,14,15	0.52	0	17,19,21	1.26	1 (5%)
2	GAL	G	4	2	11,11,12	0.38	0	15,15,17	1.46	2 (13%)
2	NAG	H	1	2	15,15,15	1.05	1 (6%)	21,21,21	3.36	3 (14%)
2	GAL	H	2	2	11,11,12	1.30	1 (9%)	15,15,17	1.99	3 (20%)
2	NAG	H	3	2	14,14,15	0.62	0	17,19,21	0.81	0
2	GAL	H	4	2	11,11,12	0.58	0	15,15,17	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2	-	2/6/26/26	0/1/1/1
2	GAL	E	2	2	-	0/2/19/22	0/1/1/1
2	NAG	E	3	2	-	0/6/23/26	0/1/1/1
2	GAL	E	4	2	-	0/2/19/22	0/1/1/1
2	NAG	F	1	2	-	0/6/26/26	0/1/1/1
2	GAL	F	2	2	-	0/2/19/22	0/1/1/1
2	NAG	F	3	2	-	0/6/23/26	0/1/1/1
2	GAL	F	4	2	-	2/2/19/22	0/1/1/1
2	NAG	G	1	2	-	0/6/26/26	0/1/1/1
2	GAL	G	2	2	-	0/2/19/22	0/1/1/1
2	NAG	G	3	2	-	0/6/23/26	0/1/1/1
2	GAL	G	4	2	-	0/2/19/22	0/1/1/1
2	NAG	H	1	2	-	4/6/26/26	0/1/1/1
2	GAL	H	2	2	-	2/2/19/22	0/1/1/1
2	NAG	H	3	2	-	0/6/23/26	0/1/1/1
2	GAL	H	4	2	-	0/2/19/22	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	NAG	O4-C4	-3.70	1.34	1.43
2	H	2	GAL	O3-C3	-3.67	1.34	1.43
2	G	2	GAL	O3-C3	-3.45	1.34	1.43
2	G	1	NAG	O4-C4	-3.44	1.34	1.43
2	E	2	GAL	O3-C3	-3.38	1.35	1.43
2	H	1	NAG	O4-C4	-3.32	1.35	1.43
2	F	2	GAL	O3-C3	-3.19	1.35	1.43
2	F	1	NAG	O4-C4	-2.76	1.36	1.43

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	NAG	O4-C4-C5	12.27	139.76	109.30
2	H	1	NAG	O4-C4-C3	-7.50	93.00	110.35
2	G	1	NAG	C1-C2-C3	-7.13	100.82	110.54
2	H	2	GAL	O3-C3-C2	4.90	119.37	109.99
2	E	1	NAG	O5-C1-C2	-4.82	104.67	109.52
2	H	2	GAL	O5-C5-C6	4.28	113.91	107.20
2	G	4	GAL	O5-C5-C6	4.14	113.70	107.20
2	E	3	NAG	O5-C1-C2	-3.36	105.99	111.29
2	H	1	NAG	C4-C3-C2	3.28	115.15	110.34
2	G	3	NAG	C1-O5-C5	3.16	116.47	112.19
2	G	1	NAG	O5-C1-C2	-3.09	106.41	109.52
2	G	1	NAG	C1-O5-C5	-2.96	108.08	113.66
2	F	3	NAG	C1-O5-C5	2.93	116.16	112.19
2	G	2	GAL	O3-C3-C4	2.88	117.02	110.35
2	E	4	GAL	O5-C1-C2	2.82	115.13	110.77
2	G	2	GAL	O3-C3-C2	-2.76	104.70	109.99
2	E	2	GAL	O3-C3-C2	-2.56	105.08	109.99
2	F	3	NAG	O5-C1-C2	-2.51	107.32	111.29
2	G	1	NAG	O4-C4-C5	-2.51	103.07	109.30
2	F	1	NAG	O5-C5-C6	2.45	112.52	106.44
2	H	2	GAL	C1-O5-C5	-2.44	108.89	112.19
2	F	4	GAL	C1-C2-C3	2.41	112.62	109.67
2	G	1	NAG	O7-C7-C8	-2.39	117.61	122.06
2	G	4	GAL	C1-C2-C3	2.35	112.56	109.67
2	E	1	NAG	C1-C2-C3	-2.33	107.36	110.54
2	F	4	GAL	O5-C1-C2	2.28	114.29	110.77
2	E	4	GAL	O5-C5-C6	2.24	110.71	107.20
2	F	2	GAL	C1-C2-C3	2.13	112.28	109.67
2	G	1	NAG	C3-C2-N2	2.11	114.59	110.62
2	E	1	NAG	O5-C5-C6	2.09	111.64	106.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	3	NAG	C1-O5-C5	2.09	115.02	112.19
2	E	2	GAL	C2-C3-C4	-2.02	107.40	110.89

There are no chirality outliers.


All (10) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	H	2	GAL	O5-C5-C6-O6
2	H	2	GAL	C4-C5-C6-O6
2	H	1	NAG	C8-C7-N2-C2
2	H	1	NAG	C4-C5-C6-O6
2	H	1	NAG	O7-C7-N2-C2
2	F	4	GAL	C4-C5-C6-O6
2	H	1	NAG	O5-C5-C6-O6
2	E	1	NAG	C8-C7-N2-C2
2	F	4	GAL	O5-C5-C6-O6
2	E	1	NAG	O7-C7-N2-C2


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	142/148 (95%)	-0.13	2 (1%) 75 82	12, 20, 28, 36	0
1	B	142/148 (95%)	-0.05	3 (2%) 63 71	13, 20, 28, 37	0
1	C	142/148 (95%)	-0.17	4 (2%) 53 58	12, 18, 29, 33	0
1	D	142/148 (95%)	-0.21	3 (2%) 63 71	12, 18, 26, 36	0
All	All	568/592 (95%)	-0.14	12 (2%) 63 71	12, 19, 29, 37	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	148	GLN	3.7
1	A	90	HIS	3.4
1	C	90	HIS	3.2
1	B	148	GLN	2.9
1	B	121	GLN	2.8
1	C	113	MET	2.7
1	A	113	MET	2.6
1	D	41	SER	2.4
1	D	113	MET	2.3
1	B	130	ARG	2.2
1	C	53	PHE	2.0
1	D	40	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	H	1	15/15	0.55	0.29	53,64,66,67	0
2	GAL	H	2	11/12	0.85	0.18	31,47,50,51	0
2	NAG	E	1	15/15	0.86	0.11	20,28,33,37	0
2	GAL	E	4	11/12	0.87	0.11	27,31,34,34	0
2	GAL	F	4	11/12	0.88	0.18	31,35,37,38	0
2	NAG	F	1	15/15	0.89	0.10	19,24,30,32	0
2	NAG	G	1	15/15	0.92	0.10	18,26,30,35	0
2	GAL	G	4	11/12	0.92	0.10	26,30,34,35	0
2	NAG	F	3	14/15	0.93	0.12	22,26,30,30	0
2	NAG	E	3	14/15	0.93	0.07	19,23,27,27	0
2	NAG	H	3	14/15	0.94	0.08	20,23,30,30	0
2	GAL	H	4	11/12	0.95	0.07	15,18,19,20	0
2	NAG	G	3	14/15	0.96	0.06	20,22,25,26	0
2	GAL	F	2	11/12	0.96	0.08	16,17,19,22	0
2	GAL	E	2	11/12	0.97	0.06	18,19,20,22	0
2	GAL	G	2	11/12	0.97	0.06	16,17,19,24	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.