



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2020 – 07:29 PM BST

PDB ID : 6ZJR
Title : Cold-adapted beta-D-galactosidase from *Arthrobacter* sp. 32cB mutant E517Q
in complex with lactulose
Authors : Rutkiewicz, M.; Bujacz, A.; Bujacz, G.
Deposited on : 2020-06-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1012	
2	B	2	
2	C	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FRU	C	1	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8309 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	989	7702	4846	1390	1447	19	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLU	-	expression tag	UNP A0A023UGN9
A	0	GLN	-	expression tag	UNP A0A023UGN9
A	517	GLN	GLU	engineered mutation	UNP A0A023UGN9

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-fructofuranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0
2	C	2	23	12	11	0	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		

- Molecule 4 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	528	Total	O	0	4
			528	528		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-galactosidase



- Molecule 2: beta-D-galactopyranose-(1-4)-beta-D-fructofuranose



- Molecule 2: beta-D-galactopyranose-(1-4)-beta-D-fructofuranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	137.43Å 137.43Å 127.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.69 – 2.00 46.69 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.69-2.00) 99.9 (46.69-2.00)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.183 , 0.210 0.188 , 0.213	Depositor DCC
R_{free} test set	2098 reflections (2.23%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtrriage
Anisotropy	0.513	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8309	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MLI, GAL, FRU, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/7928	0.55	0/10796

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7702	0	7441	27	0
2	B	23	0	21	1	0
2	C	23	0	21	4	0
3	A	2	0	0	0	0
4	A	7	0	2	0	0
5	A	24	0	18	1	0
6	A	528	0	0	2	0
All	All	8309	0	7503	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:458:ARG:HH12	2:C:2:GAL:H3	1.03	1.18
1:A:458:ARG:NH1	2:C:2:GAL:H3	1.78	0.98
1:A:458:ARG:HH12	2:C:2:GAL:C3	1.91	0.82
1:A:441:GLU:OE2	2:B:1:FRU:H62	1.98	0.64
1:A:376:LEU:HD21	1:A:388:LEU:CD1	2.28	0.62
1:A:172:ARG:HD2	1:A:391:ASP:OD2	2.07	0.54
1:A:721:VAL:HG22	1:A:836:ASP:HA	1.90	0.53
1:A:748:ILE:O	1:A:751:VAL:HG22	2.11	0.51
1:A:838:GLU:OE2	1:A:1007:LYS:NZ	2.44	0.51
1:A:906:LYS:NZ	6:A:1212:HOH:O	2.44	0.51
1:A:785:TRP:CD1	1:A:790:LEU:HD12	2.46	0.50
1:A:715:PRO:HG3	1:A:939:GLY:HA2	1.93	0.50
1:A:723:PRO:HA	1:A:810:GLU:HG2	1.93	0.49
1:A:83:GLY:HA2	1:A:86:LEU:HD11	1.96	0.47
1:A:153:GLU:HA	1:A:154:SER:HA	1.74	0.45
1:A:615:TRP:CE3	1:A:673:ALA:HB2	2.54	0.43
1:A:175:HIS:HE1	6:A:1266:HOH:O	2.02	0.43
1:A:493:GLN:HG2	1:A:539:TYR:CE1	2.55	0.42
1:A:754:GLU:OE1	1:A:968:ARG:NE	2.37	0.42
1:A:80:TRP:HA	1:A:86:LEU:HD21	2.01	0.42
1:A:80:TRP:HB2	1:A:86:LEU:HD22	2.02	0.41
1:A:108:TYR:CZ	1:A:582:VAL:HB	2.56	0.41
1:A:232:PHE:HB2	2:C:2:GAL:O3	2.20	0.41
1:A:975:HIS:NE2	5:A:1111:ACT:O	2.41	0.40
1:A:112:GLN:CD	1:A:576:VAL:CG1	2.89	0.40
1:A:553:HIS:HB2	1:A:582:VAL:HG22	2.04	0.40
1:A:786:LYS:HE3	1:A:786:LYS:HB2	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	999/1012 (99%)	969 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	788/795 (99%)	782 (99%)	6 (1%)	81	86

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	172	ARG
1	A	232	PHE
1	A	310	ARG
1	A	388	LEU
1	A	518	TYR
1	A	762	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FRU	B	1	2	11,12,12	0.93	0	10,18,18	1.64	3 (30%)
2	GAL	B	2	3,2	11,11,12	0.71	0	15,15,17	2.16	2 (13%)
2	FRU	C	1	2	11,12,12	0.71	0	10,18,18	1.16	1 (10%)
2	GAL	C	2	2	11,11,12	1.17	2 (18%)	15,15,17	1.89	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	B	1	2	-	0/5/24/24	0/1/1/1
2	GAL	B	2	3,2	-	1/2/19/22	0/1/1/1
2	FRU	C	1	2	-	2/5/24/24	0/1/1/1
2	GAL	C	2	2	-	2/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	GAL	C4-C3	-2.04	1.47	1.52
2	C	2	GAL	C2-C3	-2.03	1.49	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GAL	C1-O5-C5	7.90	122.89	112.19
2	C	2	GAL	C1-O5-C5	-4.46	106.14	112.19
2	C	1	FRU	O4-C4-C3	-3.25	102.41	112.15
2	B	1	FRU	O4-C4-C5	2.49	118.26	111.05
2	B	1	FRU	C5-C4-C3	-2.49	93.92	101.91
2	B	1	FRU	O5-C5-C4	-2.37	99.60	105.49
2	C	2	GAL	O5-C1-C2	-2.34	107.16	110.77
2	B	2	GAL	C1-C2-C3	2.29	112.47	109.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	GAL	C2-C3-C4	-2.10	107.26	110.89
2	C	2	GAL	C3-C4-C5	-2.09	106.52	110.24

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	GAL	O5-C5-C6-O6
2	C	1	FRU	C4-C5-C6-O6
2	C	1	FRU	O5-C5-C6-O6
2	B	2	GAL	O5-C5-C6-O6
2	C	2	GAL	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	GAL	4	0
2	B	1	FRU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [\(i\)](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	A	1106	-	1,3,3	6.29	1 (100%)	0,3,3	0.00	-
4	MLI	A	1103	-	0,6,6	0.00	-	0,7,7	0.00	-
5	ACT	A	1107	-	1,3,3	7.31	1 (100%)	0,3,3	0.00	-
5	ACT	A	1111	-	1,3,3	6.81	1 (100%)	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	A	1108	-	1,3,3	6.33	1 (100%)	0,3,3	0.00	-
5	ACT	A	1110	-	1,3,3	6.61	1 (100%)	0,3,3	0.00	-
5	ACT	A	1109	-	1,3,3	6.30	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLI	A	1103	-	-	0/0/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1107	ACT	CH3-C	7.31	1.58	1.48
5	A	1111	ACT	CH3-C	6.81	1.57	1.48
5	A	1110	ACT	CH3-C	6.61	1.57	1.48
5	A	1108	ACT	CH3-C	6.33	1.56	1.48
5	A	1109	ACT	CH3-C	6.30	1.56	1.48
5	A	1106	ACT	CH3-C	6.29	1.56	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1111	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	989/1012 (97%)	0.32	85 (8%) 10 9	34, 48, 77, 106	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	562	GLY	8.9
1	A	561	ASP	7.2
1	A	560	ALA	7.1
1	A	85	ALA	5.7
1	A	563	VAL	5.5
1	A	559	THR	5.4
1	A	808	GLY	5.3
1	A	86	LEU	4.7
1	A	678	ASP	4.6
1	A	82	LEU	4.5
1	A	87	ASN	4.4
1	A	299	ALA	4.1
1	A	964[A]	LYS	4.0
1	A	258	ASP	3.8
1	A	677	SER	3.6
1	A	547	VAL	3.5
1	A	679	GLY	3.4
1	A	22	GLY	3.3
1	A	516	CYS	3.3
1	A	298	ALA	3.3
1	A	716	VAL	3.3
1	A	613[A]	ARG	3.2
1	A	367	SER	3.2
1	A	388	LEU	3.2
1	A	369	TYR	3.2
1	A	227	GLY	3.1
1	A	676	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	23	ARG	3.0
1	A	260	VAL	3.0
1	A	713	GLY	3.0
1	A	331	VAL	3.0
1	A	721	VAL	3.0
1	A	387	VAL	2.9
1	A	84	GLU	2.9
1	A	714	ALA	2.9
1	A	390	CYS	2.9
1	A	1009	ARG	2.9
1	A	83	GLY	2.9
1	A	366	THR	2.8
1	A	173	LEU	2.8
1	A	642	GLY	2.8
1	A	334	HIS	2.7
1	A	519	VAL	2.6
1	A	672	LEU	2.6
1	A	300	GLY	2.6
1	A	185	GLN	2.6
1	A	371	PRO	2.6
1	A	365	ARG	2.6
1	A	558	SER	2.6
1	A	368	HIS	2.6
1	A	806	GLN	2.6
1	A	297	SER	2.5
1	A	802	ALA	2.5
1	A	807	ASP	2.5
1	A	370	PRO	2.5
1	A	719	ALA	2.5
1	A	548	TRP	2.5
1	A	586	LEU	2.4
1	A	255	GLN	2.4
1	A	330	GLY	2.4
1	A	546	PHE	2.4
1	A	392	LEU	2.4
1	A	203	VAL	2.4
1	A	804	PRO	2.4
1	A	723	PRO	2.4
1	A	725	GLN	2.4
1	A	551	LEU	2.3
1	A	333	ARG	2.3
1	A	364	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	ALA	2.3
1	A	515	LEU	2.2
1	A	838	GLU	2.2
1	A	811	THR	2.2
1	A	171	SER	2.2
1	A	228	ILE	2.2
1	A	210	TRP	2.1
1	A	271	ALA	2.1
1	A	715	PRO	2.1
1	A	717	PRO	2.1
1	A	152	ILE	2.1
1	A	645	LEU	2.1
1	A	391	ASP	2.1
1	A	433	VAL	2.1
1	A	186	GLY	2.1
1	A	809	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FRU	C	1	12/12	0.44	0.41	99,104,108,108	0
2	GAL	C	2	11/12	0.78	0.24	95,98,101,102	0
2	FRU	B	1	12/12	0.90	0.16	43,45,49,51	0
2	GAL	B	2	11/12	0.97	0.17	37,38,40,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACT	A	1109	4/4	0.81	0.17	76,79,79,80	0
4	MLI	A	1103	7/7	0.82	0.20	70,71,73,74	0
5	ACT	A	1110	4/4	0.83	0.45	76,78,79,80	0
5	ACT	A	1111	4/4	0.84	0.10	89,90,91,92	0
5	ACT	A	1107	4/4	0.87	0.37	73,74,74,75	0
5	ACT	A	1106	4/4	0.89	0.17	76,77,77,77	0
5	ACT	A	1108	4/4	0.90	0.24	64,66,67,67	0
3	NA	A	1102	1/1	0.95	0.05	68,68,68,68	0
3	NA	A	1101	1/1	0.98	0.15	38,38,38,38	0

6.5 Other polymers

There are no such residues in this entry.