



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2023 – 07:15 PM EDT

PDB ID : 2ZKG  
Title : Crystal structure of unliganded SRA domain of mouse Np95  
Authors : Arita, K.; Ariyoshi, M.; Tochio, H.; Nakamura, Y.; Shirakawa, M.  
Deposited on : 2008-03-19  
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	 7% 84% 6% 10%
1	B	210	 7% 84% 6% 10%
1	C	210	 10% 78% 9% 13%
1	D	210	 8% 78% 8% 12%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6486 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase UHRF1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	190	1514	942	283	284	5	1	1	0
1	B	190	1507	939	282	281	5	0	0	0
1	C	183	1470	915	282	268	5	11	2	0
1	D	184	1465	912	279	269	5	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	404	SER	LYS	engineered mutation	UNP Q8VDF2
B	404	SER	LYS	engineered mutation	UNP Q8VDF2
C	404	SER	LYS	engineered mutation	UNP Q8VDF2
D	404	SER	LYS	engineered mutation	UNP Q8VDF2

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	154	Total O 154 154	0	0
3	B	116	Total O 116 116	0	0
3	C	128	Total O 128 128	0	0
3	D	116	Total O 116 116	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.33Å 67.01Å 69.57Å 91.11° 65.01° 84.48°	Depositor
Resolution (Å)	29.31 – 1.77 29.31 – 1.77	Depositor EDS
% Data completeness (in resolution range)	95.2 (29.31-1.77) 95.2 (29.31-1.77)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.18 (at 1.77Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.186 , 0.217 0.195 , 0.223	Depositor DCC
$R_{free}$ test set	4845 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.3	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6486	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.66	0/1554	0.70	1/2102 (0.0%)
1	B	0.61	0/1547	0.72	1/2094 (0.0%)
1	C	0.70	0/1508	0.80	3/2038 (0.1%)
1	D	0.60	0/1503	0.73	1/2032 (0.0%)
All	All	0.64	0/6112	0.74	6/8266 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	C	436[A]	ARG	NE-CZ-NH2	-7.06	116.77	120.30
1	C	436[B]	ARG	NE-CZ-NH2	-7.06	116.77	120.30
1	D	578	LEU	CA-CB-CG	6.51	130.26	115.30
1	C	585	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	500	GLN	N-CA-CB	5.39	120.31	110.60
1	B	558	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	612	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1514	0	1444	8	0
1	B	1507	0	1440	9	0
1	C	1470	0	1416	19	0
1	D	1465	0	1409	14	0
2	A	8	0	12	0	0
2	C	4	0	6	0	0
2	D	4	0	6	0	0
3	A	154	0	0	2	0
3	B	116	0	0	4	0
3	C	128	0	0	1	0
3	D	116	0	0	0	0
All	All	6486	0	5733	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:547:LYS:H	1:B:547:LYS:HD2	1.48	0.79
1:C:484:THR:CG2	1:C:579:VAL:HG21	2.21	0.70
1:C:523:GLU:O	1:C:523:GLU:HG3	1.94	0.67
1:C:533:GLN:HG2	3:C:715:HOH:O	1.95	0.65
1:C:484:THR:HG23	1:C:579:VAL:CG2	2.27	0.64
1:C:417:ILE:HD11	1:C:426:ILE:HG22	1.80	0.64
1:D:522:ASN:ND2	1:D:524:LYS:H	1.99	0.61
1:C:484:THR:HG23	1:C:579:VAL:HG21	1.82	0.59
1:D:459:ASN:HA	1:D:515:LEU:HD11	1.85	0.59
1:C:522:ASN:ND2	1:C:524:LYS:H	2.03	0.57
1:D:522:ASN:HD22	1:D:524:LYS:H	1.52	0.56
1:A:522:ASN:ND2	1:A:524:LYS:H	2.03	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:590:GLU:HG3	3:B:619:HOH:O	2.06	0.55
1:C:484:THR:CG2	1:C:579:VAL:CG2	2.84	0.54
1:C:469:GLY:HA2	1:C:548:HIS:HD2	1.73	0.54
1:C:522:ASN:HD22	1:C:523:GLU:N	2.05	0.53
1:D:602:THR:HG23	1:D:607:LEU:HB2	1.89	0.53
1:C:522:ASN:HD22	1:C:522:ASN:C	2.13	0.52
1:C:439:VAL:HG13	1:C:440:GLN:OE1	2.09	0.52
1:A:522:ASN:HD22	1:A:523:GLU:N	2.08	0.51
1:A:414:GLU:HG2	3:A:733:HOH:O	2.12	0.50
1:D:522:ASN:HD22	1:D:523:GLU:N	2.10	0.50
1:D:601:ARG:HA	1:D:604:GLN:HG2	1.94	0.49
1:C:504:GLN:CD	1:C:579:VAL:HG23	2.33	0.49
1:B:414:GLU:CG	3:B:691:HOH:O	2.63	0.47
1:D:522:ASN:HD22	1:D:522:ASN:C	2.18	0.47
1:A:439:VAL:O	1:A:443:GLU:HG3	2.14	0.47
1:C:469:GLY:HA2	1:C:548:HIS:CD2	2.50	0.46
1:B:547:LYS:HD2	1:B:547:LYS:N	2.25	0.45
1:A:414:GLU:CG	3:A:733:HOH:O	2.64	0.45
1:D:459:ASN:HA	1:D:515:LEU:CD1	2.47	0.44
1:D:474:ASP:O	1:D:548:HIS:HE1	2.00	0.44
1:C:417:ILE:HD11	1:C:426:ILE:CG2	2.46	0.44
1:C:484:THR:HG23	1:C:579:VAL:HG22	1.96	0.44
1:B:413:THR:HG23	1:B:413:THR:O	2.18	0.44
1:C:596:ARG:O	1:C:596:ARG:HD3	2.18	0.44
1:B:414:GLU:HG2	3:B:691:HOH:O	2.17	0.43
1:B:474:ASP:O	1:B:548:HIS:HE1	2.01	0.43
1:A:439:VAL:HG23	1:A:450:HIS:NE2	2.35	0.42
1:B:613:GLU:N	3:B:712:HOH:O	2.52	0.42
1:A:603:ARG:NE	1:C:532:ARG:HD3	2.35	0.41
1:B:602:THR:HG23	1:B:607:LEU:HB2	2.01	0.41
1:A:602:THR:HG23	1:A:607:LEU:HB2	2.03	0.41
1:D:459:ASN:O	1:D:459:ASN:CG	2.58	0.41
1:D:474:ASP:O	1:D:548:HIS:CE1	2.74	0.41
1:C:484:THR:HG21	1:C:579:VAL:HG21	2.00	0.41
1:D:416:THR:HG22	1:D:416:THR:O	2.21	0.41
1:D:417:ILE:HG22	1:D:418:VAL:HG13	2.03	0.40
1:D:435:TRP:CE3	1:D:441:VAL:HG22	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	187/210 (89%)	180 (96%)	7 (4%)	0	100	100
1	B	186/210 (89%)	179 (96%)	6 (3%)	1 (0%)	29	14
1	C	181/210 (86%)	174 (96%)	6 (3%)	1 (1%)	25	11
1	D	181/210 (86%)	178 (98%)	3 (2%)	0	100	100
All	All	735/840 (88%)	711 (97%)	22 (3%)	2 (0%)	41	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	470	GLY
1	B	470	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/171 (92%)	155 (98%)	3 (2%)	57	43
1	B	157/171 (92%)	154 (98%)	3 (2%)	57	43
1	C	152/171 (89%)	148 (97%)	4 (3%)	46	29
1	D	152/171 (89%)	146 (96%)	6 (4%)	32	15
All	All	619/684 (90%)	603 (97%)	16 (3%)	46	29

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	413	THR
1	A	522	ASN
1	A	574	LYS
1	B	472	GLU
1	B	522	ASN
1	B	547	LYS
1	C	505	LYS
1	C	522	ASN
1	C	547	LYS
1	C	548	HIS
1	D	459	ASN
1	D	501	SER
1	D	522	ASN
1	D	547	LYS
1	D	578	LEU
1	D	596	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	522	ASN
1	B	440	GLN
1	B	479	ASN
1	B	509	ASN
1	B	522	ASN
1	B	548	HIS
1	C	509	ASN
1	C	522	ASN
1	C	533	GLN
1	D	508	ASN
1	D	509	ASN
1	D	522	ASN
1	D	548	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EDO	D	4	-	3,3,3	0.47	0	2,2,2	0.38	0
2	EDO	C	3	-	3,3,3	0.45	0	2,2,2	0.30	0
2	EDO	A	1	-	3,3,3	0.41	0	2,2,2	0.42	0
2	EDO	A	2	-	3,3,3	0.44	0	2,2,2	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	D	4	-	-	0/1/1/1	-
2	EDO	C	3	-	-	0/1/1/1	-
2	EDO	A	1	-	-	1/1/1/1	-
2	EDO	A	2	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	EDO	O1-C1-C2-O2
2	A	2	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	190/210 (90%)	0.50	15 (7%) 12 12	23, 28, 41, 51	1 (0%)
1	B	190/210 (90%)	0.55	14 (7%) 14 14	25, 31, 41, 54	0
1	C	183/210 (87%)	0.78	22 (12%) 4 4	24, 28, 43, 50	0
1	D	184/210 (87%)	0.61	16 (8%) 10 9	26, 31, 42, 48	0
All	All	747/840 (88%)	0.61	67 (8%) 9 9	23, 30, 42, 54	1 (0%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	470	GLY	8.3
1	C	577	PHE	8.2
1	D	417	ILE	8.1
1	C	417	ILE	7.9
1	C	574	LYS	7.1
1	C	471	TYR	6.7
1	A	470	GLY	6.1
1	B	613	GLU	6.0
1	B	413	THR	5.8
1	A	548	HIS	5.1
1	C	470	GLY	5.0
1	D	416	THR	5.0
1	D	415	CYS	5.0
1	A	413	THR	4.7
1	C	529	GLU	4.6
1	C	472	GLU	4.5
1	C	416	THR	4.3
1	C	576	GLY	4.3
1	B	500	GLN	4.3
1	D	604	GLN	4.0
1	C	575	SER	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	C	488	GLY	3.9
1	A	547	LYS	3.8
1	A	500	GLN	3.6
1	A	472	GLU	3.6
1	B	471	TYR	3.6
1	B	472	GLU	3.5
1	D	472	GLU	3.2
1	A	524	LYS	3.2
1	B	524	LYS	3.2
1	D	459	ASN	3.2
1	A	414	GLU	3.1
1	B	603	ARG	3.1
1	D	537	VAL	3.1
1	C	547	LYS	3.0
1	B	414	GLU	2.9
1	C	531	TRP	2.9
1	C	548	HIS	2.9
1	B	539	VAL	2.9
1	A	539	VAL	2.8
1	A	469	GLY	2.8
1	A	613	GLU	2.7
1	C	415	CYS	2.7
1	A	596	ARG	2.6
1	A	471	TYR	2.6
1	C	469	GLY	2.6
1	A	540	VAL	2.6
1	B	467	LEU	2.6
1	C	562	ILE	2.5
1	A	473	ASP	2.5
1	C	578	LEU	2.5
1	C	584	LEU	2.5
1	C	533	GLN	2.5
1	D	577	PHE	2.4
1	D	529	GLU	2.4
1	D	524	LYS	2.4
1	D	470	GLY	2.4
1	D	574	LYS	2.4
1	D	500	GLN	2.4
1	B	488	GLY	2.3
1	C	537	VAL	2.3
1	B	513	LEU	2.2
1	B	537	VAL	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	608	THR	2.1
1	D	603	ARG	2.1
1	C	520	PRO	2.0
1	D	596	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	D	4	4/4	0.82	0.15	50,50,51,53	0
2	EDO	A	1	4/4	0.84	0.33	45,46,47,48	0
2	EDO	A	2	4/4	0.90	0.18	34,40,41,44	0
2	EDO	C	3	4/4	0.91	0.13	44,44,46,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.