

wwPDB X-ray Structure Validation Summary Report (i)

Nov 14, 2023 – 04:03 PM JST

PDB ID	:	5ZL1
Title	:	Hexameric structure of copper-containing nitrite reductase of an anammox
		organism KSU-1
Authors	:	Hira, D.; Matsumura, M.; Kitamura, R.; Fujii, T.
Deposited on	:	2018-03-26
Resolution	:	3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	326	69%	23%	• 6%			
1	В	326	% 69%	23%	• 6%			
1	С	326	65%	28%	• 6%			
1	D	326	^{2%} 66%	25%	• 6%			



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 9615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	206	Total	С	Ν	Ο	\mathbf{S}	0	0	0
	A	300	2387	1537	403	437	10	0	0	0
1	р	206	Total	С	Ν	0	S	0	0	0
1	Б	300	2387	1537	403	437	10			
1	С	C 306	Total	С	Ν	0	S	0	0	0
1			2387	1537	403	437	10			0
1 D	П	306	Total	С	Ν	Ο	S	0	0	0
	D	300	2387	1537	403	437	10		0	0

• Molecule 1 is a protein called Putative copper-type nitrite reductase.

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference			
А	1	MET	-	expression tag	UNP I3IR93			
А	2	GLY	-	expression tag	UNP I3IR93			
А	3	SER	-	expression tag	UNP I3IR93			
А	4	SER	-	expression tag	UNP I3IR93			
А	5	HIS	-	expression tag	UNP I3IR93			
А	6	HIS	-	expression tag	UNP I3IR93			
А	7	HIS	-	expression tag	UNP I3IR93			
А	8	HIS	-	expression tag	UNP I3IR93			
А	9	HIS	-	expression tag	UNP I3IR93			
А	10	HIS	-	expression tag	UNP I3IR93			
А	11	SER	-	expression tag	UNP I3IR93			
А	12	SER	-	expression tag	UNP I3IR93			
А	13	GLY	-	expression tag	UNP I3IR93			
А	14	LEU	-	expression tag	UNP I3IR93			
А	15	VAL	-	expression tag	UNP I3IR93			
А	16	PRO	-	expression tag	UNP I3IR93			
А	17	ARG	-	expression tag	UNP I3IR93			
А	18	GLY	-	expression tag	UNP I3IR93			
A	19	SER	-	expression tag	UNP I3IR93			
А	20	HIS	-	expression tag	UNP I3IR93			
А	21	MET	-	expression tag	UNP I3IR93			
	Continued on next page							



Chain	Residue	Modelled	Actual	Comment	Reference
В	1	MET	-	expression tag	UNP I3IR93
В	2	GLY	-	expression tag	UNP I3IR93
В	3	SER	-	expression tag	UNP I3IR93
В	4	SER	-	expression tag	UNP I3IR93
В	5	HIS	-	expression tag	UNP I3IR93
В	6	HIS	-	expression tag	UNP I3IR93
В	7	HIS	-	expression tag	UNP I3IR93
В	8	HIS	-	expression tag	UNP I3IR93
В	9	HIS	-	expression tag	UNP I3IR93
В	10	HIS	-	expression tag	UNP I3IR93
В	11	SER	-	expression tag	UNP I3IR93
В	12	SER	-	expression tag	UNP I3IR93
В	13	GLY	-	expression tag	UNP I3IR93
В	14	LEU	-	expression tag	UNP I3IR93
В	15	VAL	-	expression tag	UNP I3IR93
В	16	PRO	-	expression tag	UNP I3IR93
В	17	ARG	-	expression tag	UNP I3IR93
В	18	GLY	-	expression tag	UNP I3IR93
В	19	SER	-	expression tag	UNP I3IR93
В	20	HIS	-	expression tag	UNP I3IR93
В	21	MET	-	expression tag	UNP I3IR93
С	1	MET	-	expression tag	UNP I3IR93
С	2	GLY	-	expression tag	UNP I3IR93
С	3	SER	-	expression tag	UNP I3IR93
С	4	SER	-	expression tag	UNP I3IR93
С	5	HIS	-	expression tag	UNP I3IR93
С	6	HIS	-	expression tag	UNP I3IR93
С	7	HIS	-	expression tag	UNP I3IR93
С	8	HIS	-	expression tag	UNP I3IR93
С	9	HIS	-	expression tag	UNP I3IR93
С	10	HIS	-	expression tag	UNP I3IR93
С	11	SER	-	expression tag	UNP I3IR93
C	12	SER	-	expression tag	UNP I3IR93
С	13	GLY	-	expression tag	UNP I3IR93
С	14	LEU	-	expression tag	UNP I3IR93
С	15	VAL	-	expression tag	UNP I3IR93
С	16	PRO	-	expression tag	UNP I3IR93
C	17	ARG	-	expression tag	UNP I3IR93
С	18	GLY	-	expression tag	UNP I3IR93
С	19	SER	-	expression tag	UNP I3IR93
С	20	HIS	-	expression tag	UNP I3IR93
С	21	MET	-	expression tag	UNP I3IR93



Chain	Residue	Modelled	Actual	Comment	Reference
D	1	MET	-	expression tag	UNP I3IR93
D	2	GLY	-	expression tag	UNP I3IR93
D	3	SER	-	expression tag	UNP I3IR93
D	4	SER	-	expression tag	UNP I3IR93
D	5	HIS	-	expression tag	UNP I3IR93
D	6	HIS	-	expression tag	UNP I3IR93
D	7	HIS	-	expression tag	UNP I3IR93
D	8	HIS	-	expression tag	UNP I3IR93
D	9	HIS	-	expression tag	UNP I3IR93
D	10	HIS	-	expression tag	UNP I3IR93
D	11	SER	-	expression tag	UNP I3IR93
D	12	SER	-	expression tag	UNP I3IR93
D	13	GLY	-	expression tag	UNP I3IR93
D	14	LEU	-	expression tag	UNP I3IR93
D	15	VAL	-	expression tag	UNP I3IR93
D	16	PRO	-	expression tag	UNP I3IR93
D	17	ARG	-	expression tag	UNP I3IR93
D	18	GLY	-	expression tag	UNP I3IR93
D	19	SER	-	expression tag	UNP I3IR93
D	20	HIS	-	expression tag	UNP I3IR93
D	21	MET	-	expression tag	UNP I3IR93

• Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Cu 2 2	0	0
2	В	3	Total Cu 3 3	0	0
2	С	2	Total Cu 2 2	0	0
2	D	2	Total Cu 2 2	0	0

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Ca 1 1	0	0
3	В	1	Total Ca 1 1	0	0



Continued from previous page...

ЪЛ 1				7 0	
Mol	Chain	Residues	Atoms	ZeroUcc	AltConf
3	С	1	Total Ca 1 1	0	0
3	D	1	Total Ca 1 1	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	D	1	$\begin{array}{c cc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	14	Total O 14 14	0	0
5	В	6	Total O 6 6	0	0
5	С	9	Total O 9 9	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	5	Total O 5 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Putative copper-type nitrite reductase



22.27 22.28 22.28 22.28 22.28 22.28 22.39 22.39 22.39 22.44 22.44 22.44 22.44 22.44 22.45 22.55 22.55 22.55 22.55 22.55 22.55 22.55 23.05 25.55

• Molecule 1: Putative copper-type nitrite reductase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	270.22Å 270.22Å 128.91Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Bosolution (Å)	40.00 - 3.20	Depositor
Resolution (A)	39.96 - 3.20	EDS
% Data completeness	99.9 (40.00-3.20)	Depositor
(in resolution range)	100.0 (39.96-3.20)	EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$5.20 (at 3.18 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
P. P.	0.240 , 0.254	Depositor
n, n_{free}	0.229 , 0.236	DCC
R_{free} test set	1507 reflections (5.07%)	wwPDB-VP
Wilson B-factor $(Å^2)$	60.0	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.32 , 35.1	EDS
L-test for $twinning^2$	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	9615	wwPDB-VP
Average B, all atoms $(Å^2)$	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles		
IVIOI	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.42	0/2457	0.67	0/3335	
1	В	0.41	0/2457	0.66	0/3335	
1	С	0.46	0/2457	0.71	1/3335~(0.0%)	
1	D	0.43	0/2457	0.68	1/3335~(0.0%)	
All	All	0.43	0/9828	0.68	2/13340~(0.0%)	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	С	275	ARG	NE-CZ-NH1	5.11	122.85	120.30
1	D	234	GLY	N-CA-C	-5.06	100.46	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2387	0	2312	55	0
1	В	2387	0	2312	58	0
1	С	2387	0	2312	86	0
1	D	2387	0	2312	81	0
2	А	2	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	3	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	А	5	0	0	1	0
4	В	5	0	0	1	0
4	С	5	0	0	1	0
4	D	5	0	0	0	0
5	А	14	0	0	0	0
5	В	6	0	0	0	0
5	C	9	0	0	0	0
5	D	5	0	0	0	0
All	All	9615	0	9248	273	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 273 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:235:ASN:HB3	1:C:324:HIS:NE2	1.66	1.10
1:C:234:GLY:O	1:C:235:ASN:ND2	1.95	1.00
1:A:269:LYS:HE2	1:A:278:PHE:CZ	2.00	0.97
1:B:269:LYS:HE2	1:B:278:PHE:CZ	2.04	0.91
1:C:235:ASN:CB	1:C:324:HIS:NE2	2.36	0.89

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	304/326~(93%)	292~(96%)	12~(4%)	0	100	100
1	В	304/326~(93%)	290~(95%)	14~(5%)	0	100	100
1	С	304/326~(93%)	291 (96%)	13~(4%)	0	100	100
1	D	304/326~(93%)	293~(96%)	11 (4%)	0	100	100
All	All	1216/1304~(93%)	1166~(96%)	50~(4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	248/265~(94%)	236~(95%)	12 (5%)	25	61
1	В	248/265~(94%)	240~(97%)	8(3%)	39	71
1	С	248/265~(94%)	235~(95%)	13 (5%)	23	59
1	D	248/265~(94%)	235~(95%)	13 (5%)	23	59
All	All	992/1060~(94%)	946~(95%)	46 (5%)	27	63

5 of 46 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	С	235	ASN
1	D	78	THR
1	С	241	GLN
1	С	302	THR
1	D	134	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such side chains are listed below:

Mol	Chain	Res	Type
1	D	27	ASN
1	D	235	ASN
1	D	324	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	D	265	GLN
1	С	265	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 13 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Turne	Chain	Dec	Tinle	Bond lengths			Bond angles		
	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
4	SO4	А	404	-	4,4,4	0.34	0	$6,\!6,\!6$	0.09	0
4	SO4	D	404	-	4,4,4	0.34	0	6,6,6	0.06	0
4	SO4	В	405	-	4,4,4	0.33	0	6,6,6	0.06	0
4	SO4	С	404	-	4,4,4	0.33	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	404	SO4	1	0
4	В	405	SO4	1	0
4	С	404	SO4	1	0

3 monomers are involved in 3 short contacts:

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	306/326~(93%)	-0.17	1 (0%) 94 92	30, 42, 61, 77	0
1	В	306/326~(93%)	-0.03	2 (0%) 87 81	37, 50, 72, 90	0
1	С	306/326~(93%)	0.14	5 (1%) 72 59	42, 62, 90, 107	0
1	D	306/326~(93%)	0.14	5 (1%) 72 59	45, 63, 90, 119	0
All	All	1224/1304~(93%)	0.02	13 (1%) 80 69	30, 54, 83, 119	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	22	VAL	4.0
1	D	199	HIS	3.9
1	С	105	LEU	3.0
1	D	202	GLY	2.8
1	С	198	PRO	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



5ZL1

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
2	CU	А	401	1/1	0.62	0.11	49,49,49,49	0
3	CA	С	403	1/1	0.71	0.20	56, 56, 56, 56	0
2	CU	С	401	1/1	0.78	0.17	72,72,72,72	0
4	SO4	D	404	5/5	0.79	0.27	90,90,92,93	0
2	CU	С	402	1/1	0.83	0.10	72,72,72,72	0
4	SO4	С	404	5/5	0.84	0.37	81,81,82,85	0
4	SO4	В	405	5/5	0.87	0.28	74,75,76,77	0
2	CU	D	402	1/1	0.88	0.11	62,62,62,62	0
4	SO4	А	404	5/5	0.90	0.29	68,69,69,70	0
3	CA	А	403	1/1	0.90	0.14	42,42,42,42	0
2	CU	D	401	1/1	0.92	0.21	71,71,71,71	0
2	CU	В	402	1/1	0.92	0.10	59, 59, 59, 59, 59	0
2	CU	В	403	1/1	0.93	0.17	60,60,60,60	0
2	CU	А	402	1/1	0.94	0.08	57,57,57,57	0
3	CA	D	403	1/1	0.94	0.08	53,53,53,53	0
2	CU	В	401	1/1	0.96	0.16	60,60,60,60	0
3	CA	В	404	1/1	0.96	0.14	49,49,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





































6.5 Other polymers (i)

There are no such residues in this entry.

