



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 23, 2024 – 12:21 PM JST

PDB ID : 8ZQA  
Title : Crystal structure of 1,4-alpha-glucan branching protein from *Rhodothermus profundus*  
Authors : Li, Q.; Zong, Z.Y.; Liu, W.D.  
Deposited on : 2024-06-01  
Resolution : 2.91 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2797 (2.94-2.90)
Clashscore	180529	3049 (2.94-2.90)
Ramachandran outliers	177936	2981 (2.94-2.90)
Sidechain outliers	177891	2983 (2.94-2.90)
RSRZ outliers	164620	2799 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	620	 2% 65% 31% ..
1	B	620	 2% 72% 24% ..
1	C	620	 2% 66% 30% ..
1	D	620	 % 68% 26% ..
1	E	620	 2% 73% 24% ..
1	F	620	 2% 64% 30% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	620	
1	H	620	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 39395 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1,4-alpha-glucan branching enzyme GlgB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	603	Total 4886	C 3190	N 817	O 865	S 14	0	0	0
1	B	603	Total 4925	C 3212	N 832	O 867	S 14	0	0	0
1	C	606	Total 4925	C 3209	N 826	O 875	S 15	0	0	0
1	D	599	Total 4888	C 3186	N 822	O 866	S 14	0	0	0
1	E	604	Total 4917	C 3202	N 830	O 871	S 14	0	0	0
1	F	593	Total 4859	C 3166	N 818	O 861	S 14	0	0	0
1	G	606	Total 4923	C 3213	N 826	O 869	S 15	0	0	0
1	H	603	Total 4800	C 3138	N 791	O 857	S 14	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	45	Total 45	O 45	0	0
2	B	34	Total 34	O 34	0	0
2	C	31	Total 31	O 31	0	0
2	D	27	Total 27	O 27	0	0
2	E	26	Total 26	O 26	0	0
2	F	51	Total 51	O 51	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	G	34	Total	O	0	0
			34	34		
2	H	24	Total	O	0	0
			24	24		


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1,4-alpha-glucan branching enzyme GlgB




- Molecule 1: 1,4-alpha-glucan branching enzyme GlgB







• Molecule 1: 1,4-alpha-glucan branching enzyme GlgB




• Molecule 1: 1,4-alpha-glucan branching enzyme GlgB







• Molecule 1: 1,4-alpha-glucan branching enzyme GlgB




• Molecule 1: 1,4-alpha-glucan branching enzyme GlgB








● Molecule 1: 1,4-alpha-glucan branching enzyme GlgB





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.30Å 112.27Å 166.26Å 73.59° 73.77° 65.25°	Depositor
Resolution (Å)	100.10 – 2.91 100.10 – 2.91	Depositor EDS
% Data completeness (in resolution range)	98.9 (100.10-2.91) 98.9 (100.10-2.91)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 2.91Å)	Xtrriage
Refinement program	PHENIX (1.21_5207: ???)	Depositor
R, $R_{free}$	0.197 , 0.256 0.208 , 0.250	Depositor DCC
$R_{free}$ test set	129530 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.6	Xtrriage
Anisotropy	0.400	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 61.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.012 for h,h-k,h-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	39395	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/5080	0.63	0/6952
1	B	0.35	0/5123	0.60	0/7011
1	C	0.37	0/5121	0.62	0/7009
1	D	0.38	0/5082	0.62	0/6952
1	E	0.34	0/5112	0.61	0/6993
1	F	0.35	0/5049	0.58	0/6902
1	G	0.38	0/5118	0.64	0/7000
1	H	0.43	0/4993	0.67	0/6851
All	All	0.37	0/40678	0.62	0/55670

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4886	0	4484	119	0
1	B	4925	0	4535	110	0
1	C	4925	0	4517	120	0
1	D	4888	0	4499	112	0
1	E	4917	0	4523	110	0
1	F	4859	0	4500	127	0
1	G	4923	0	4527	113	0
1	H	4800	0	4320	156	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	45	0	0	3	0
2	B	34	0	0	2	0
2	C	31	0	0	0	0
2	D	27	0	0	0	0
2	E	26	0	0	1	0
2	F	51	0	0	1	0
2	G	34	0	0	1	0
2	H	24	0	0	1	0
All	All	39395	0	35905	956	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 956 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:5:THR:HG22	1:D:7:GLU:H	1.16	1.05
1:F:15:GLY:HA2	1:F:284:ASN:HD21	1.33	0.94
1:H:154:TRP:HD1	1:H:466:MET:HE2	1.34	0.92
1:H:145:SER:H	1:H:182:THR:HG22	1.38	0.87
1:C:426:VAL:HG13	1:C:432:SER:HA	1.57	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	597/620 (96%)	577 (97%)	19 (3%)	1 (0%)	44 72
1	B	599/620 (97%)	580 (97%)	18 (3%)	1 (0%)	44 72
1	C	602/620 (97%)	582 (97%)	19 (3%)	1 (0%)	44 72

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	593/620 (96%)	571 (96%)	21 (4%)	1 (0%)	44	72
1	E	598/620 (96%)	576 (96%)	21 (4%)	1 (0%)	44	72
1	F	587/620 (95%)	558 (95%)	28 (5%)	1 (0%)	44	72
1	G	600/620 (97%)	576 (96%)	23 (4%)	1 (0%)	44	72
1	H	597/620 (96%)	582 (98%)	14 (2%)	1 (0%)	44	72
All	All	4773/4960 (96%)	4602 (96%)	163 (3%)	8 (0%)	44	72

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	420	PRO
1	B	420	PRO
1	E	420	PRO
1	F	420	PRO
1	H	420	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	490/528 (93%)	464 (95%)	26 (5%)	19	48
1	B	497/528 (94%)	476 (96%)	21 (4%)	25	57
1	C	496/528 (94%)	468 (94%)	28 (6%)	17	46
1	D	494/528 (94%)	464 (94%)	30 (6%)	15	42
1	E	496/528 (94%)	477 (96%)	19 (4%)	28	61
1	F	494/528 (94%)	465 (94%)	29 (6%)	16	43
1	G	494/528 (94%)	467 (94%)	27 (6%)	18	46
1	H	472/528 (89%)	443 (94%)	29 (6%)	15	42
All	All	3933/4224 (93%)	3724 (95%)	209 (5%)	19	48

5 of 209 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	314	ASP
1	F	462	LYS
1	H	299	VAL
1	E	409	LEU
1	F	214	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	496	GLN
1	H	60	GLN
1	H	530	GLN
1	H	472	GLN
1	G	492	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	603/620 (97%)	-0.15	13 (2%) 62 56	19, 34, 56, 109	0
1	B	603/620 (97%)	-0.20	11 (1%) 67 62	17, 33, 54, 103	0
1	C	606/620 (97%)	-0.23	14 (2%) 61 55	18, 33, 60, 100	0
1	D	599/620 (96%)	-0.13	9 (1%) 71 66	22, 36, 57, 93	0
1	E	604/620 (97%)	-0.15	12 (1%) 64 58	21, 38, 60, 95	0
1	F	593/620 (95%)	-0.18	11 (1%) 66 60	20, 35, 54, 79	0
1	G	606/620 (97%)	-0.01	10 (1%) 69 63	22, 39, 64, 100	0
1	H	603/620 (97%)	0.36	31 (5%) 34 29	25, 48, 73, 88	0
All	All	4817/4960 (97%)	-0.09	111 (2%) 61 55	17, 37, 63, 109	0

The worst 5 of 111 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	420	PRO	5.3
1	G	326	GLY	5.0
1	A	272	THR	4.5
1	C	266	HIS	4.3
1	H	585	GLY	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.