



# Full wwPDB X-ray Structure Validation Report i

Nov 14, 2023 – 04:29 PM JST

PDB ID : 5ZS0  
Title : Structure of glycoprotein B Domain IV of pseudorabies virus with 7B11 antibody  
Authors : Hu, X.L.; Yang, F.L.  
Deposited on : 2018-04-26  
Resolution : 3.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see references i) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 3.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4628 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 7B11 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C 1631	N 1012	O 279	S 334	6	0	0

- Molecule 2 is a protein called 7B11 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	205	Total	C 1559	N 988	O 257	S 308	6	0	0

- Molecule 3 is a protein called Envelope glycoprotein B,Envelope glycoprotein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	187	Total	C 1438	N 891	O 261	S 274	12	0	0


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	542	GLY	-	linker	UNP A0A0U3FH21
C	543	GLY	-	linker	UNP A0A0U3FH21
C	544	SER	-	linker	UNP A0A0U3FH21
C	545	GLY	-	linker	UNP A0A0U3FH21
C	701	HIS	-	expression tag	UNP A0A1Q0AKY5
C	702	HIS	-	expression tag	UNP A0A1Q0AKY5
C	703	HIS	-	expression tag	UNP A0A1Q0AKY5
C	704	HIS	-	expression tag	UNP A0A1Q0AKY5
C	705	HIS	-	expression tag	UNP A0A1Q0AKY5
C	706	HIS	-	expression tag	UNP A0A1Q0AKY5
C	707	HIS	-	expression tag	UNP A0A1Q0AKY5
C	708	HIS	-	expression tag	UNP A0A1Q0AKY5


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 7B11 light chain



- Molecule 2: 7B11 heavy chain



- Molecule 3: Envelope glycoprotein B, Envelope glycoprotein B



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	128.91Å 128.91Å 194.91Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 3.29 37.21 – 3.29	Depositor EDS
% Data completeness (in resolution range)	80.9 (50.00-3.29) 81.1 (37.21-3.29)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.12	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.78 (at 3.32Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
$R$ , $R_{free}$	0.241 , 0.310 0.249 , 0.312	Depositor DCC
$R_{free}$ test set	559 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.0	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 14.3	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.48$ , $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	4628	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	0/1666	0.71	0/2262
2	B	0.63	0/1598	0.88	2/2179 (0.1%)
3	C	0.54	0/1463	0.89	3/1983 (0.2%)
All	All	0.56	0/4727	0.83	5/6424 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	4

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	183	LEU	CA-CB-CG	6.59	130.45	115.30
3	C	665	ARG	NE-CZ-NH2	-5.26	117.67	120.30
2	B	116	THR	N-CA-C	5.08	124.72	111.00
3	C	586	ARG	NE-CZ-NH1	5.04	122.82	120.30
3	C	665	ARG	NE-CZ-NH1	5.03	122.81	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	115	LEU	Peptide
2	B	123	THR	Peptide
2	B	124	PRO	Peptide
2	B	3	GLN	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1631	0	1557	32	0
2	B	1559	0	1522	43	0
3	C	1438	0	1390	23	1
All	All	4628	0	4469	91	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:ALA:HB3	1:A:108:ARG:NH2	1.47	1.26
1:A:15:ALA:CB	1:A:108:ARG:HH22	1.60	1.15
1:A:110:ASP:HA	1:A:140:TYR:HB3	1.18	1.09
1:A:110:ASP:CA	1:A:140:TYR:HB3	1.86	1.04
1:A:15:ALA:CB	1:A:108:ARG:NH2	2.22	0.96
1:A:110:ASP:HA	1:A:140:TYR:CB	1.99	0.92
2:B:116:THR:HG23	2:B:117:VAL:HG13	1.53	0.89
1:A:110:ASP:O	1:A:111:ALA:CB	2.28	0.81
2:B:22:CYS:SG	2:B:108:TYR:CE2	2.75	0.80
1:A:15:ALA:HB3	1:A:108:ARG:HH22	0.69	0.74
1:A:110:ASP:C	1:A:140:TYR:HB3	2.13	0.69
1:A:110:ASP:O	1:A:111:ALA:HB3	1.92	0.69
2:B:115:LEU:O	2:B:116:THR:HG22	1.93	0.68
2:B:4:LEU:HD22	2:B:108:TYR:CZ	2.28	0.67
2:B:39:GLN:HE21	2:B:43:LYS:HA	1.62	0.65
1:A:110:ASP:OD1	1:A:140:TYR:HD1	1.80	0.63
3:C:662:ASN:O	3:C:662:ASN:ND2	2.33	0.62
1:A:50:ARG:NH2	1:A:91:TYR:OH	2.34	0.61
3:C:576:SER:O	3:C:580:THR:HG23	2.01	0.61
3:C:116:ALA:HB2	3:C:609:TYR:CE2	2.37	0.59
1:A:124:GLN:HE22	1:A:131:SER:N	2.01	0.58
2:B:4:LEU:HB3	2:B:108:TYR:CG	2.39	0.58
2:B:4:LEU:HD22	2:B:108:TYR:CE1	2.39	0.58
3:C:641:GLY:HA3	3:C:650:LEU:O	2.04	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ASP:HA	1:A:140:TYR:CD1	2.40	0.57
2:B:4:LEU:HD13	2:B:108:TYR:CD1	2.39	0.57
1:A:110:ASP:HA	1:A:140:TYR:CG	2.41	0.55
2:B:96:CYS:HB3	2:B:108:TYR:CG	2.42	0.55
2:B:57:TYR:HB2	3:C:622:THR:HG21	1.89	0.55
3:C:580:THR:HG22	3:C:587:VAL:H	1.72	0.53
2:B:34:MET:HB3	2:B:79:LEU:HD22	1.91	0.53
2:B:143:THR:HG23	2:B:188:THR:HG22	1.92	0.51
1:A:89:GLN:NE2	2:B:103:PHE:HB2	2.25	0.51
1:A:29:VAL:CG1	1:A:32:ALA:HB3	2.41	0.51
2:B:22:CYS:SG	2:B:108:TYR:CZ	3.03	0.51
1:A:110:ASP:OD1	1:A:140:TYR:CD1	2.63	0.51
3:C:651:ILE:HD12	3:C:651:ILE:H	1.77	0.49
3:C:598:ILE:HD13	3:C:598:ILE:N	2.28	0.49
2:B:117:VAL:HG11	2:B:179:ASP:OD2	2.11	0.49
2:B:4:LEU:HB3	2:B:108:TYR:CD2	2.48	0.49
2:B:36:TRP:CG	2:B:81:LEU:HD22	2.47	0.49
2:B:53:GLY:HA2	2:B:72:ARG:NH1	2.27	0.49
2:B:160:TRP:CZ3	2:B:201:CYS:HB2	2.48	0.49
1:A:67:SER:HA	1:A:71:PHE:CE2	2.48	0.48
2:B:4:LEU:HD13	2:B:108:TYR:CE1	2.48	0.48
2:B:176:LEU:HD21	2:B:179:ASP:HA	1.94	0.48
1:A:110:ASP:O	1:A:111:ALA:HB2	2.10	0.48
2:B:3:GLN:O	2:B:4:LEU:HD23	2.15	0.47
3:C:136:VAL:HG13	3:C:598:ILE:HD11	1.97	0.47
3:C:616:VAL:HG12	3:C:617:PRO:HD2	1.97	0.47
3:C:618:GLY:O	3:C:620:ARG:NE	2.45	0.47
2:B:6:GLU:OE2	2:B:108:TYR:HB3	2.15	0.47
1:A:95:ASP:N	1:A:95:ASP:OD1	2.48	0.47
3:C:127:CYS:HB3	3:C:588:SER:HB2	1.97	0.47
1:A:13:THR:O	1:A:107:LYS:HB2	2.15	0.46
2:B:64:VAL:HG22	2:B:68:PHE:CG	2.51	0.46
3:C:120:GLU:N	3:C:120:GLU:OE2	2.49	0.46
3:C:126:VAL:HG22	3:C:649:LEU:HB2	1.98	0.46
1:A:174:SER:HG	2:B:170:HIS:CE1	2.34	0.46
2:B:4:LEU:HB3	2:B:108:TYR:CD1	2.51	0.46
1:A:195:GLU:HG3	1:A:206:VAL:HG22	1.98	0.45
2:B:124:PRO:HA	2:B:147:LEU:O	2.16	0.45
2:B:45:LEU:HD22	2:B:45:LEU:N	2.31	0.45
2:B:79:LEU:HD23	2:B:96:CYS:SG	2.56	0.45
3:C:610:VAL:HG22	3:C:629:VAL:HG12	1.99	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:ALA:HB2	1:A:108:ARG:NH2	2.24	0.45
2:B:14:PRO:HD3	2:B:116:THR:HA	1.99	0.44
3:C:626:ARG:NH1	3:C:643:LEU:O	2.50	0.44
1:A:29:VAL:HG12	1:A:32:ALA:HB3	2.00	0.44
3:C:116:ALA:HB2	3:C:609:TYR:CD2	2.53	0.44
2:B:105:LEU:HD23	2:B:106:PHE:N	2.33	0.44
3:C:698:ASN:O	3:C:699:LEU:HD23	2.18	0.44
2:B:59:TYR:CE2	3:C:658:PRO:HB3	2.52	0.44
2:B:42:GLU:HG2	2:B:43:LYS:CG	2.48	0.43
3:C:628:LEU:HD22	3:C:653:ARG:CD	2.49	0.42
1:A:201:SER:OG	1:A:203:SER:O	2.25	0.42
2:B:4:LEU:HD22	2:B:108:TYR:OH	2.19	0.42
2:B:27:PHE:CE1	2:B:29:PHE:HA	2.54	0.42
2:B:33:ALA:HB2	3:C:620:ARG:O	2.20	0.42
1:A:164:THR:HG23	2:B:172:PHE:CD1	2.55	0.42
3:C:591:MET:O	3:C:592:LEU:HD23	2.20	0.42
1:A:15:ALA:CB	1:A:108:ARG:CZ	2.95	0.41
1:A:50:ARG:HA	1:A:50:ARG:HD3	1.95	0.41
2:B:163:GLY:O	2:B:165:LEU:N	2.54	0.41
1:A:94:TYR:CD1	1:A:96:LEU:HD13	2.55	0.41
2:B:42:GLU:HG2	2:B:43:LYS:HG2	2.03	0.41
3:C:111:ARG:HD2	3:C:111:ARG:HA	1.91	0.41
1:A:135:PHE:CD2	2:B:186:SER:HB2	2.56	0.41
2:B:64:VAL:HG21	2:B:68:PHE:CD2	2.56	0.41
2:B:6:GLU:N	2:B:6:GLU:OE1	2.54	0.40
2:B:44:ARG:O	2:B:45:LEU:C	2.59	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:594:ASP:OD1	3:C:626:ARG:NH2[2_745]	2.17	0.03

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/211 (99%)	184 (88%)	21 (10%)	4 (2%)	8	34
2	B	201/219 (92%)	163 (81%)	28 (14%)	10 (5%)	2	14
3	C	183/254 (72%)	160 (87%)	22 (12%)	1 (0%)	29	61
All	All	593/684 (87%)	507 (86%)	71 (12%)	15 (2%)	5	27

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	111	ALA
2	B	87	ARG
2	B	123	THR
2	B	124	PRO
2	B	164	SER
1	A	32	ALA
1	A	51	ALA
2	B	45	LEU
2	B	116	THR
2	B	100	GLU
1	A	56	THR
2	B	40	SER
3	C	114	VAL
2	B	117	VAL
2	B	155	PRO

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/186 (100%)	154 (83%)	31 (17%)	2	9
2	B	178/188 (95%)	149 (84%)	29 (16%)	2	10
3	C	152/205 (74%)	130 (86%)	22 (14%)	3	14
All	All	515/579 (89%)	433 (84%)	82 (16%)	2	11

All (82) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	MET
1	A	13	THR
1	A	14	SER
1	A	18	ARG
1	A	23	CYS
1	A	43	SER
1	A	45	LYS
1	A	46	LEU
1	A	50	ARG
1	A	70	ASP
1	A	74	THR
1	A	88	CYS
1	A	89	GLN
1	A	93	ASN
1	A	95	ASP
1	A	107	LYS
1	A	110	ASP
1	A	114	THR
1	A	116	SER
1	A	123	GLU
1	A	131	SER
1	A	134	CYS
1	A	143	ASP
1	A	147	LYS
1	A	155	ARG
1	A	176	SER
1	A	177	SER
1	A	179	LEU
1	A	182	THR
1	A	194	CYS
1	A	205	ILE
2	B	3	GLN
2	B	6	GLU
2	B	11	LEU
2	B	35	SER
2	B	45	LEU
2	B	64	VAL
2	B	69	THR
2	B	72	ARG
2	B	74	ASN
2	B	76	LYS
2	B	88	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	89	ASP
2	B	97	THR
2	B	99	GLN
2	B	105	LEU
2	B	109	TRP
2	B	113	THR
2	B	114	ILE
2	B	117	VAL
2	B	122	THR
2	B	125	PRO
2	B	126	SER
2	B	127	VAL
2	B	146	CYS
2	B	166	SER
2	B	182	THR
2	B	183	LEU
2	B	186	SER
2	B	210	THR
3	C	120	GLU
3	C	133	SER
3	C	547	ASP
3	C	564	ARG
3	C	572	ARG
3	C	573	LEU
3	C	586	ARG
3	C	595	VAL
3	C	598	ILE
3	C	613	SER
3	C	615	ARG
3	C	622	THR
3	C	630	THR
3	C	632	GLU
3	C	633	HIS
3	C	636	THR
3	C	652	SER
3	C	660	THR
3	C	662	ASN
3	C	669	LEU
3	C	685	VAL
3	C	698	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	GLN
1	A	156	GLN
1	A	157	ASN
1	A	210	ASN
2	B	77	ASN
2	B	205	HIS
3	C	633	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/211 (100%)	0.12	3 (1%) 75 75	30, 80, 108, 120	0
2	B	205/219 (93%)	-0.18	0 100 100	28, 48, 97, 127	0
3	C	187/254 (73%)	-0.14	2 (1%) 80 81	27, 56, 100, 117	0
All	All	603/684 (88%)	-0.06	5 (0%) 86 86	27, 63, 106, 127	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	110	ASP	4.0
1	A	108	ARG	3.2
3	C	547	ASP	2.4
3	C	550	SER	2.4
1	A	190	ASN	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.