

wwPDB NMR Structure Validation Summary Report (i)

Jun 6, 2023 – 07:55 pm BST

PDB ID : 6ZV1 BMRB ID : 34538

Title: TFIIS N-terminal domain (TND) from human IWS1

Authors : Veverka, V. Deposited on : 2020-07-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-ShiftChecker} &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 90%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive
Metric	$(\# ext{Entries})$	$(\# ext{Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain						
1	A	150	81%	5%	14%				



2 Ensemble composition and analysis (i)

This entry contains 40 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues							
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model				
1	A:556-A:684 (129)	0.37	9				

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 8 clusters and 10 single-model clusters were found.

Cluster number	Models
1	7, 9, 20, 23, 30, 31
2	6, 10, 12, 13, 17, 35
3	14, 15, 19, 39
4	2, 4, 8, 21
5	5, 26, 29, 33
6	11, 16
7	18, 24
8	25, 38
Single-model clusters	1; 3; 22; 27; 28; 32; 34; 36; 37; 40



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2394 atoms, of which 1222 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Protein IWS1 homolog.

Mol	Chain	Residues	Atoms						Trace
1	Λ	150	Total	С	Н	N	О	S	0
	1 A	150	2394	742	1222	203	221	6	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	545	SER	-	expression tag	UNP Q96ST2
A	546	ASN	-	expression tag	UNP Q96ST2
A	547	ALA	_	expression tag	UNP Q96ST2
A	548	ALA	-	expression tag	UNP Q96ST2
A	549	SER	_	expression tag	UNP Q96ST2
A	693	GLY	_	expression tag	UNP Q96ST2
A	694	SER	-	expression tag	UNP Q96ST2




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Protein IWS1 homolog



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 9. Colouring as in section 4.1 above.

• Molecule 1: Protein IWS1 homolog





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: torsion angle dynamics, molecular dynamics.

Of the 100 calculated structures, 40 were deposited, based on the following criterion: structures with the least restraint violations.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	
YASARA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1781
Number of shifts mapped to atoms	1781
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	90%



6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	I	Bond lengths	Bond angles		
		RMSZ	#Z>5	RMSZ	#Z>5	
1	A	0.55 ± 0.01	$0\pm0/1033$ ($0.0\pm~0.0\%$)	0.71 ± 0.03	$1\pm 1/1389$ ($0.1\pm$ 0.1%)	
All	All	0.55	0/41320 (0.0%)	0.71	57/55560 (0.1%)	

There are no bond-length outliers.

5 of 7 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mal	Chain	Dag	Trino	Atoma	Z	Observed(0)	$Ideal(^{o})$	Mod	dels
Mol	Chain	Res	Type	Atoms	L	$Observed(^o)$	Ideal(*)	Worst	Total
1	A	575	ARG	NE-CZ-NH1	6.83	123.71	120.30	14	22
1	A	634	ARG	NE-CZ-NH1	6.22	123.41	120.30	34	20
1	A	626	ARG	NE-CZ-NH1	6.21	123.41	120.30	27	2
1	A	641	LEU	CB-CG-CD1	5.58	120.49	111.00	21	9
1	A	672	ARG	NE-CZ-NH1	5.09	122.85	120.30	12	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1018	1078	1078	1±1
All	All	40720	43120	43120	36

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.



5	of 10	unique	clashes	are	listed	below.	sorted	bv	their	clash	magnitude.
_	-	· · · · · · · · · · · · · · · · · · ·)		•			

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:591:LEU:HD23	1:A:633:ILE:HG23	0.55	1.76	32	1
1:A:563:MET:SD	1:A:594:VAL:HG12	0.49	2.46	36	15
1:A:616:ILE:HG21	1:A:641:LEU:HD11	0.49	1.84	27	1
1:A:616:ILE:HG23	1:A:637:LEU:HD13	0.46	1.87	27	1
1:A:638:LEU:HA	1:A:641:LEU:HD12	0.45	1.88	31	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	129/150 (86%)	125±1 (97±1%)	3±1 (2±1%)	1±0 (1±0%)	24	71
All	All	5160/6000 (86%)	5015 (97%)	103 (2%)	42 (1%)	24	71

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	645	PRO	40
1	A	600	LYS	2

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	A	115/131 (88%)	108±2 (94±2%)	7±2 (6±2%)	21	69
All	All	4600/5240 (88%)	4302 (94%)	298 (6%)	21	69

5 of 40 unique residues with a non-rotameric sidechain are listed below. They are sorted by the



frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	640	ILE	29
1	A	591	LEU	28
1	A	642	GLN	25
1	A	604	LYS	20
1	A	590	LEU	19

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 90% for the well-defined parts and 86% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: starch_output

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1781
Number of shifts mapped to atoms	1781
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	5

The following errors were found when reading this chemical shift list.

• Chemical shift has been reported more than once. All 3 occurrences are reported below.

List ID	Chain	Res	Type	Atom		Shift Dat	a
LIST ID				ites Type	Atom	Value	Uncertainty
1	A	631	LEU	HD11	1.118	0.000	•
1	A	631	LEU	HD12	1.118	0.000	
1	A	631	LEU	HD13	1.118	0.000	•

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	143	-0.88 ± 0.09	Should be checked
$^{13}C_{\beta}$	135	0.26 ± 0.14	None needed (< 0.5 ppm)
¹³ C′	136	-0.72 ± 0.09	Should be applied
^{15}N	130	0.43 ± 0.22	None needed (< 0.5 ppm)



7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 90%, i.e. 1639 atoms were assigned a chemical shift out of a possible 1821. 0 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	622/635~(98%)	251/255~(98%)	$252/258 \ (98\%)$	119/122 (98%)
Sidechain	973/1125 (86%)	671/732 (92%)	294/352 (84%)	8/41 (20%)
Aromatic	44/61 (72%)	25/31~(81%)	18/26 (69%)	1/4 (25%)
Overall	1639/1821 (90%)	947/1018 (93%)	564/636 (89%)	128/167 (77%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	587	LYS	HB3	-0.29	0.46 - 3.04	-7.9
1	A	586	LYS	Н	11.58	5.24 - 11.12	5.8
1	A	669	LYS	Н	11.26	5.24 - 11.12	5.2
1	A	671	SER	Н	11.23	5.45 - 11.10	5.2
1	A	681	LEU	HB3	-0.26	-0.26 - 3.31	-5.0

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:







8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	2916
Intra-residue ($ i-j =0$)	763
Sequential ($ i-j =1$)	689
Medium range ($ i-j >1$ and $ i-j <5$)	660
Long range (i-j ≥5)	804
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	220
Number of unmapped restraints	0
Number of restraints per residue	20.9
Number of long range restraints per residue ¹	5.4

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	10.5	0.2
0.2-0.5 (Medium)	6.8	0.5
>0.5 (Large)	7.0	1.58



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

$\mathbf{Bins}\;(^{\circ})$	Average number of violations per model	\mathbf{Max} (°)
1.0-10.0 (Small)	3.2	5.0
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Doctroints type	Count	% ¹	Vio	lated	3	Consis	tentl	${ m y~Violated^4}$
Restraints type	Count	70	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$
Intra-residue (i-j =0)	763	26.2	9	1.2	0.3	4	0.5	0.1
Backbone-Backbone	1	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	629	21.6	8	1.3	0.3	4	0.6	0.1
Sidechain-Sidechain	133	4.6	1	0.8	0.0	0	0.0	0.0
Sequential (i-j =1)	689	23.6	14	2.0	0.5	2	0.3	0.1
Backbone-Backbone	117	4.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	465	15.9	8	1.7	0.3	2	0.4	0.1
Sidechain-Sidechain	107	3.7	6	5.6	0.2	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	660	22.6	23	3.5	0.8	1	0.2	0.0
Backbone-Backbone	179	6.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	335	11.5	13	3.9	0.4	1	0.3	0.0
Sidechain-Sidechain	146	5.0	10	6.8	0.3	0	0.0	0.0
Long range ($ i-j \ge 5$)	804	27.6	46	5.7	1.6	1	0.1	0.0
Backbone-Backbone	3	0.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	284	9.7	15	5.3	0.5	0	0.0	0.0
Sidechain-Sidechain	517	17.7	31	6.0	1.1	1	0.2	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	2916	100.0	92	3.2	3.2	8	0.3	0.3
Backbone-Backbone	300	10.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	1713	58.7	44	2.6	1.5	7	0.4	0.2
Sidechain-Sidechain	903	31.0	48	5.3	1.6	1	0.1	0.0

 $^{^1}$ percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	ations	5	Mean (Å)	M (Å)	SD^6 (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
1	5	5	5	13	0	28	0.38	1.29	0.36	0.18
2	5	8	6	13	0	32	0.35	1.23	0.31	0.18
3	4	5	6	8	0	23	0.43	1.3	0.38	0.16
4	4	8	7	12	0	31	0.44	1.3	0.35	0.33
5	5	6	6	10	0	27	0.36	1.21	0.33	0.2
6	4	7	10	13	0	34	0.4	1.37	0.36	0.22
7	4	5	7	8	0	24	0.39	1.18	0.34	0.22
8	4	7	7	16	0	34	0.4	1.34	0.36	0.18
9	4	4	6	11	0	25	0.4	1.17	0.35	0.2
10	5	8	7	17	0	37	0.37	1.37	0.33	0.22
11	6	5	5	11	0	27	0.46	1.32	0.39	0.38

Continued on next page...




Continued from previous page...

Continuea ji			$\frac{pagon}{\text{ober o}}$		ations		3.5 (8)	7 (§)	GD6 (8)	7.7.1. (8)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
12	5	8	6	18	0	37	0.37	1.23	0.33	0.19
13	4	9	8	10	0	31	0.35	1.25	0.33	0.17
14	5	3	3	7	0	18	0.49	1.18	0.37	0.34
15	5	7	7	14	0	33	0.41	1.31	0.35	0.23
16	4	4	3	6	0	17	0.63	1.27	0.37	0.48
17	5	3	7	10	0	25	0.39	1.18	0.33	0.26
18	5	2	1	4	0	12	0.69	1.33	0.38	0.66
19	5	3	4	8	0	20	0.57	1.31	0.38	0.46
20	5	5	5	13	0	28	0.36	1.16	0.33	0.18
21	5	3	5	6	0	19	0.55	1.18	0.38	0.43
22	5	8	9	15	0	37	0.38	1.32	0.35	0.2
23	4	6	7	10	0	27	0.41	1.27	0.36	0.25
24	4	3	3	5	0	15	0.63	1.51	0.45	0.48
25	4	3	2	7	0	16	0.64	1.19	0.37	0.54
26	5	6	6	10	0	27	0.45	1.34	0.36	0.38
27	4	2	2	5	0	13	0.67	1.21	0.36	0.73
28	4	6	6	13	0	29	0.43	1.3	0.36	0.27
29	4	4	3	4	0	15	0.62	1.58	0.47	0.42
30	4	5	8	8	0	25	0.43	1.32	0.36	0.24
31	4	3	3	6	0	16	0.54	1.31	0.43	0.44
32	5	5	4	5	0	19	0.46	1.31	0.4	0.23
33	5	6	8	7	0	26	0.42	1.24	0.37	0.22
34	4	2	1	4	0	11	0.74	1.31	0.31	0.72
35	5	8	13	17	0	43	0.37	1.33	0.33	0.22
36	5	4	4	12	0	25	0.4	1.32	0.36	0.18
37	4	3	4	10	0	21	0.41	1.3	0.35	0.22
38	4	2	1	4	0	11	0.69	1.17	0.34	0.71
39	4	3	2	6	0	15	0.64	1.39	0.37	0.49
40	4	3	2	7	0	16	0.6	1.16	0.33	0.5

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation



9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2824(IR:754, SQ:675, MR:637, LR:758, IC:0) restraints are not violated in the ensemble.

Nu	ımber	of vio	lated	restr	aints	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Count ⁶	%		
2	2	11	16	0	31	1	2.5		
0	1	0	5	0	6	2	5.0		
0	2	0	3	0	5	3	7.5		
2	0	0	3	0	5	4	10.0		
0	1	3	2	0	6	5	12.5		
0	0	1	3	0	4	6	15.0		

Continued on next page...




Continued from previous page...

		$\frac{\mathbf{o} \mathbf{n} \mathbf{v} \mathbf{i} \mathbf{o}}{\mathbf{o} \mathbf{f} \mathbf{v} \mathbf{i} \mathbf{o}}$				Fraction	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
0	0	0	0	0	0	7	17.5
0	0	0	0	0	0	8	20.0
0	0	0	1	0	1	9	22.5
1	1	2	1	0	5	10	25.0
0	1	0	1	0	2	11	27.5
0	1	0	0	0	1	12	30.0
0	0	1	0	0	1	13	32.5
0	1	0	0	0	1	14	35.0
0	0	0	2	0	2	15	37.5
0	0	0	0	0	0	16	40.0
0	0	0	1	0	1	17	42.5
0	0	1	0	0	1	18	45.0
0	0	0	0	0	0	19	47.5
0	0	0	1	0	1	20	50.0
0	0	0	1	0	1	21	52.5
0	0	0	0	0	0	22	55.0
0	0	0	1	0	1	23	57.5
0	1	1	0	0	2	24	60.0
0	0	0	0	0	0	25	62.5
0	0	0	0	0	0	26	65.0
0	0	0	1	0	1	27	67.5
0	0	0	0	0	0	28	70.0
0	0	1	0	0	1	29	72.5
0	0	0	0	0	0	30	75.0
0	1	0	0	0	1	31	77.5
0	0	0	2	0	2	32	80.0
0	0	1	0	0	1	33	82.5
0	0	0	0	0	0	34	85.0
0	0	0	0	0	0	35	87.5
0	0	0	1	0	1	36	90.0
0	0	0	0	0	0	37	92.5
0	0	0	0	0	0	38	95.0
0	0	0	0	0	0	39	97.5
4	2	1	1	0	8	40	100.0

 $^{^1{\}rm Intra-residue}$ restraints, $^2{\rm Sequential}$ restraints, $^3{\rm Medium}$ range restraints, $^4{\rm Long}$ range restraints, $^5{\rm Inter-chain}$ restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	\mathbf{SD}^1 (Å)	Median (Å)
(1,1639)	1:A:580:GLN:H	1:A:580:GLN:HE21	40	1.23	0.08	1.25
(1,1538)	1:A:568:ASN:HD21	1:A:569:GLU:H	40	1.15	0.04	1.16
(1,1512)	1:A:567:MET:H	1:A:568:ASN:HD21	40	0.97	0.04	0.96
(1,2088)	1:A:579:ASN:H	1:A:579:ASN:HD21	40	0.95	0.18	1.0
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	40	0.91	0.4	0.97
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	40	0.91	0.4	0.97
(1,2133)	1:A:642:GLN:H	1:A:642:GLN:HE21	40	0.64	0.2	0.73
(1,2080)	1:A:576:GLN:HA	1:A:579:ASN:HD21	40	0.48	0.02	0.48
(1,1622)	1:A:578:ASN:H	1:A:578:ASN:HD21	40	0.42	0.02	0.43
(1,2122)	1:A:642:GLN:HE22	1:A:677:MET:HG3	36	0.43	0.18	0.47
(1,2135)	1:A:639:LYS:HA	1:A:642:GLN:HE21	33	0.5	0.23	0.44


¹Number of violated models, ²Standard deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	29	1.58
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	29	1.58
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	24	1.51
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	24	1.51
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	39	1.39
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	39	1.39
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	6	1.37
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	6	1.37
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	10	1.37
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	10	1.37
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	8	1.34

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	8	1.34
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	26	1.34
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	26	1.34
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD2	35	1.33
(1,2127)	1:A:642:GLN:HE22	1:A:680:LYS:HD3	35	1.33
(1,1639)	1:A:580:GLN:H	1:A:580:GLN:HE21	18	1.33
(1,1639)	1:A:580:GLN:H	1:A:580:GLN:HE21	29	1.33



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle tree	Count	$\%^{1}$	$_{oz1}$ Violated ³			Consistently Violated ⁴		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
PSI	110	50.0	10	9.1	4.5	0	0.0	0.0
PHI	110	50.0	7	6.4	3.2	0	0.0	0.0
Total	220	100.0	17	7.7	7.7	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories



10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

N. 1. 1. I.	Nun	nber o	of violations	3 (0)	3 (0)	GD (0)	3.5.11 (0)
Model ID	PSI	PHI	Total	$\mathbf{Mean} (^{\circ})$	$\mathbf{Max} \ (^{\circ})$	SD (°)	Median (°)
1	2	2	4	2.68	4.6	1.24	2.45
2	2	3	5	3.1	4.6	1.17	3.5
3	2	3	5	2.56	4.7	1.14	2.4
4	3	2	5	2.94	4.5	1.16	2.6
5	3	3	6	2.73	5.0	1.4	2.1
6	2	3	5	2.66	4.8	1.23	2.4
7	1	3	4	2.95	4.9	1.22	2.65
8	1	2	3	3.43	5.0	1.23	3.3
9	3	3	6	2.08	4.9	1.41	1.25
10	2	4	6	2.77	4.6	1.21	2.55
11	0	2	2	3.95	4.8	0.85	3.95
12	2	3	5	3.36	4.9	1.52	4.4
13	3	3	6	2.95	4.8	1.19	2.8
14	1	0	1	5.0	5.0	0.0	5.0
15	1	2	3	3.93	4.8	1.02	4.5
16	0	1	1	4.8	4.8	0.0	4.8
17	1	2	3	2.77	4.6	1.43	2.6
18	1	0	1	5.0	5.0	0.0	5.0
19	0	1	1	5.0	5.0	0.0	5.0
20	1	4	5	3.08	4.9	1.27	2.9
21	1	1	2	3.05	4.7	1.65	3.05
22	4	2	6	3.0	4.4	0.88	2.85
23	2	2	4	2.6	4.6	1.2	2.15
24	1	0	1	5.0	5.0	0.0	5.0
25	1	0	1	5.0	5.0	0.0	5.0
26	2	3	5	2.68	4.9	1.3	1.9
27	1	0	1	4.9	4.9	0.0	4.9
28	2	3	5	3.28	5.0	1.23	3.6
29	0	1	1	4.9	4.9	0.0	4.9
30	2	3	5	2.66	4.7	1.55	1.6
31	1	0	1	4.8	4.8	0.0	4.8
32	1	1	2	3.05	4.8	1.75	3.05
33	2	4	6	3.05	4.8	1.48	3.15
34	1	0	1	4.9	4.9	0.0	4.9
35	2	3	5	3.0	4.7	1.1	2.8
36	0	1	1	4.8	4.8	0.0	4.8
37	0	1	1	4.8	4.8	0.0	4.8
38	1	0	1	4.7	4.7	0.0	4.7


Continued on next page...



Continued from previous page...

Model ID	Nun	nber c	of violations	Mean (°)	Max (°)	SD (°)	Median (°)	
Wiodei 1D	PSI	PHI	Total	Mean ()	Max ()	SD ()	Median ()	
39	1	0	1	4.8	4.8	0.0	4.8	
40	1	0	1	5.0	5.0	0.0	5.0	

10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints			Fraction of the ensemble		
PSI	PHI	Total	Count ¹	%	
3	1	4	1	2.5	
2	2	4	2	5.0	
1	0	1	3	7.5	
0	1	1	4	10.0	

Continued on next page...




 $Continued\ from\ previous\ page...$

		of violated restraints	Fraction of the ensemble			
PSI	PHI	Total	$Count^1$	%		
1	0	1	5	12.5		
0	0	0	6	15.0		
0	0	0	7	17.5		
0	0	0	8	20.0		
0	0	0	9	22.5		
2	0	2	10	25.0		
0	0	0	11	27.5		
0	0	0	12	30.0		
0	1	1	13	32.5		
0	0	0	14	35.0		
0	0	0	15	37.5		
0	0	0	16	40.0		
0	0	0	17	42.5		
0	0	0	18	45.0		
0	0	0	19	47.5		
0	0	0	20	50.0		
0	0	0	21	52.5		
1	1	2	22	55.0		
0	0	0	23	57.5		
0	0	0	24	60.0		
0	0	0	25	62.5		
0	0	0	26	65.0		
0	1	1	27	67.5		
0	0	0	28	70.0		
0	0	0	29	72.5		
0	0	0	30	75.0		
0	0	0	31	77.5		
0	0	0	32	80.0		
0	0	0	33	82.5		
0	0	0	34	85.0		
0	0	0	35	87.5		
0	0	0	36	90.0		
0	0	0	37	92.5		
0	0	0	38	95.0		
0	0	0	39	97.5		
0	0	0	40	100.0		

 $^{^{1}}$ Number of models with violations



10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram: Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,77)	1:A:596:MET:C	1:A:597:HIS:N	1:A:597:HIS:CA	1:A:597:HIS:C	27	3.54	1.1	3.8
(1,121)	1:A:624:PRO:C	1:A:625:ASP:N	1:A:625:ASP:CA	1:A:625:ASP:C	22	3.84	1.28	4.55
(1,102)	1:A:612:VAL:N	1:A:612:VAL:CA	1:A:612:VAL:C	1:A:613:MET:N	22	2.02	0.67	2.0
(1,137)	1:A:635:GLU:C	1:A:636:GLU:N	1:A:636:GLU:CA	1:A:636:GLU:C	13	3.8	1.32	4.7
(1,142)	1:A:638:LEU:N	1:A:638:LEU:CA	1:A:638:LEU:C	1:A:639:LYS:N	10	4.91	0.1	4.95
(1,156)	1:A:648:SER:N	1:A:648:SER:CA	1:A:648:SER:C	1:A:649:GLN:N	10	2.11	0.96	1.6
(1,18)	1:A:563:MET:N	1:A:563:MET:CA	1:A:563:MET:C	1:A:564:ILE:N	5	1.76	0.89	1.5
(1,35)	1:A:571:ALA:C	1:A:572:GLU:N	1:A:572:GLU:CA	1:A:572:GLU:C	4	2.28	0.72	2.05
(1,84)	1:A:601:GLN:N	1:A:601:GLN:CA	1:A:601:GLN:C	1:A:602:ASP:N	3	4.03	1.23	4.9
(1,3)	1:A:555:ASP:C	1:A:556:ALA:N	1:A:556:ALA:CA	1:A:556:ALA:C	2	4.65	0.05	4.65


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram: Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,77)	1:A:596:MET:C	1:A:597:HIS:N	1:A:597:HIS:CA	1:A:597:HIS:C	5	5.0
(1,77)	1:A:596:MET:C	1:A:597:HIS:N	1:A:597:HIS:CA	1:A:597:HIS:C	19	5.0
(1,142)	1:A:638:LEU:N	1:A:638:LEU:CA	1:A:638:LEU:C	1:A:639:LYS:N	14	5.0
(1,142)	1:A:638:LEU:N	1:A:638:LEU:CA	1:A:638:LEU:C	1:A:639:LYS:N	18	5.0
(1,142)	1:A:638:LEU:N	1:A:638:LEU:CA	1:A:638:LEU:C	1:A:639:LYS:N	24	5.0
(1,142)	1:A:638:LEU:N	1:A:638:LEU:CA	1:A:638:LEU:C	1:A:639:LYS:N	25	5.0
(1,142)	1:A:638:LEU:N	1:A:638:LEU:CA	1:A:638:LEU:C	1:A:639:LYS:N	40	5.0
(1,137)	1:A:635:GLU:C	1:A:636:GLU:N	1:A:636:GLU:CA	1:A:636:GLU:C	28	5.0
(1,121)	1:A:624:PRO:C	1:A:625:ASP:N	1:A:625:ASP:CA	1:A:625:ASP:C	8	5.0
(1,84)	1:A:601:GLN:N	1:A:601:GLN:CA	1:A:601:GLN:C	1:A:602:ASP:N	9	4.9

