



# Full wwPDB X-ray Structure Validation Report i

Aug 16, 2023 – 03:07 PM EDT

PDB ID : 1ZW2  
Title : Vinculin Head (0-258) in Complex with the Talin Rod residues 2345-2369  
Authors : Gingras, A.R.; Ziegler, W.H.; Barsukov, I.L.; Roberts, G.C.; Critchley, D.R.; Emsley, J.  
Deposited on : 2005-06-03  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vinculin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	251	1957	1232	334	377	14	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	cloning artifact	UNP P12003
A	-19	GLY	-	cloning artifact	UNP P12003
A	-18	SER	-	cloning artifact	UNP P12003
A	-17	SER	-	cloning artifact	UNP P12003
A	-16	HIS	-	expression tag	UNP P12003
A	-15	HIS	-	expression tag	UNP P12003
A	-14	HIS	-	expression tag	UNP P12003
A	-13	HIS	-	expression tag	UNP P12003
A	-12	HIS	-	expression tag	UNP P12003
A	-11	HIS	-	expression tag	UNP P12003
A	-10	SER	-	cloning artifact	UNP P12003
A	-9	SER	-	cloning artifact	UNP P12003
A	-8	GLY	-	cloning artifact	UNP P12003
A	-7	LEU	-	cloning artifact	UNP P12003
A	-6	VAL	-	cloning artifact	UNP P12003
A	-5	PRO	-	cloning artifact	UNP P12003
A	-4	ARG	-	cloning artifact	UNP P12003
A	-3	GLY	-	cloning artifact	UNP P12003
A	-2	SER	-	cloning artifact	UNP P12003
A	-1	HIS	-	cloning artifact	UNP P12003
A	0	MET	-	cloning artifact	UNP P12003

- Molecule 2 is a protein called talin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	21	Total	C	N	O	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	71	Total	O	0	0
3	B	6	Total	O	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vinculin



- Molecule 2: talin



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.03Å   70.04Å   96.03Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	20.00 – 2.10 19.80 – 2.10	Depositor EDS
% Data completeness (in resolution range)	88.8 (20.00-2.10) 88.8 (19.80-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.23 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
$R$ , $R_{free}$	0.255 , 0.295 0.269 , 0.303	Depositor DCC
$R_{free}$ test set	981 reflections (5.23%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.2	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 65.3	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2170	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.63	0/1980	0.74	3/2679 (0.1%)
2	B	0.79	0/135	1.03	1/182 (0.5%)
All	All	0.64	0/2115	0.76	4/2861 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	2	2
2	B	0	2
All	All	2	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	220	GLN	N-CA-C	7.25	130.58	111.00
2	B	2364	SER	CB-CA-C	5.75	121.02	110.10
1	A	27	GLU	N-CA-C	5.16	124.92	111.00
1	A	89	ARG	CG-CD-NE	5.05	122.41	111.80

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	27	GLU	CA
1	A	220	GLN	CA

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	219	SER	Peptide
1	A	86	LYS	Mainchain
2	B	2362	ALA	Mainchain
2	B	2363	ALA	Mainchain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1957	0	2036	43	0
2	B	136	0	151	5	0
3	A	71	0	0	7	0
3	B	6	0	0	1	0
All	All	2170	0	2187	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:0:MET:N	3:A:328:HOH:O	2.11	0.84
1:A:87:LEU:HD13	1:A:110:GLY:HA3	1.70	0.72
1:A:85:THR:HB	3:A:326:HOH:O	1.89	0.71
1:A:18:GLN:HE22	2:B:2361:LYS:HD2	1.54	0.71
1:A:89:ARG:HH11	1:A:89:ARG:CG	2.05	0.70
1:A:41:ALA:HB3	1:A:42:PRO:HD3	1.76	0.66
1:A:41:ALA:HB3	1:A:42:PRO:CD	2.26	0.65
1:A:3:PHE:O	1:A:182:THR:HG21	1.96	0.64
1:A:70:LYS:NZ	3:A:284:HOH:O	2.32	0.62
1:A:18:GLN:HE22	2:B:2361:LYS:CD	2.13	0.61
1:A:153:MET:O	1:A:157:VAL:HG23	2.02	0.60
1:A:52:ASN:HD21	1:A:55:ARG:HH21	1.50	0.59
1:A:138:LYS:HE2	3:A:329:HOH:O	2.04	0.56
2:B:2346:LEU:O	2:B:2350:LYS:HG3	2.07	0.54
1:A:229:ARG:HH11	1:A:230:ASN:HD21	1.54	0.53
1:A:89:ARG:HH11	1:A:89:ARG:HG2	1.73	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:GLU:O	1:A:13:GLU:HB2	2.09	0.51
1:A:0:MET:HB2	1:A:1:PRO:HD3	1.91	0.51
1:A:25:MET:O	1:A:26:HIS:C	2.49	0.50
1:A:69:LEU:HD12	1:A:124:THR:HG22	1.93	0.49
1:A:91:ALA:HB3	3:A:259:HOH:O	2.12	0.48
1:A:31:VAL:HG12	1:A:34:LYS:HB2	1.96	0.48
1:A:156:LEU:HD11	1:A:209:LYS:HG3	1.95	0.48
1:A:246:VAL:HA	1:A:249:LEU:HD22	1.96	0.47
2:B:2364:SER:HB3	3:B:79:HOH:O	2.14	0.47
1:A:178:GLN:HG3	1:A:191:VAL:HG23	1.96	0.46
1:A:34:LYS:O	1:A:35:ALA:CB	2.63	0.46
1:A:87:LEU:CD1	1:A:110:GLY:HA3	2.42	0.46
1:A:18:GLN:NE2	2:B:2361:LYS:HD2	2.27	0.46
1:A:104:ARG:HG3	1:A:105:ASP:N	2.31	0.45
1:A:9:GLU:O	1:A:13:GLU:CB	2.65	0.45
1:A:142:GLU:O	1:A:145:THR:HB	2.16	0.45
1:A:18:GLN:HE21	1:A:18:GLN:HB3	1.57	0.44
1:A:85:THR:CG2	3:A:326:HOH:O	2.65	0.44
1:A:138:LYS:CE	3:A:329:HOH:O	2.64	0.44
1:A:101:VAL:N	1:A:102:PRO:HD2	2.33	0.44
1:A:14:PRO:O	1:A:18:GLN:HG3	2.18	0.44
1:A:152:THR:HB	1:A:155:ASP:H	1.82	0.44
1:A:121:LEU:C	1:A:121:LEU:HD23	2.37	0.43
1:A:86:LYS:O	1:A:87:LEU:C	2.53	0.42
1:A:13:GLU:HB3	1:A:14:PRO:HD3	1.99	0.42
1:A:58:LYS:O	1:A:62:GLN:HG2	2.19	0.42
1:A:3:PHE:CZ	1:A:12:LEU:HB3	2.55	0.41
1:A:211:PHE:O	1:A:215:LYS:HB2	2.20	0.41
1:A:41:ALA:CB	1:A:42:PRO:CD	2.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	249/279 (89%)	233 (94%)	10 (4%)	6 (2%)	6 2
2	B	19/25 (76%)	19 (100%)	0	0	100 100
All	All	268/304 (88%)	252 (94%)	10 (4%)	6 (2%)	6 2

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35	ALA
1	A	220	GLN
1	A	26	HIS
1	A	219	SER
1	A	25	MET
1	A	30	GLU

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	223/246 (91%)	198 (89%)	25 (11%)	6 3
2	B	12/15 (80%)	11 (92%)	1 (8%)	11 7
All	All	235/261 (90%)	209 (89%)	26 (11%)	6 3

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	HIS
1	A	12	LEU
1	A	18	GLN
1	A	27	GLU
1	A	28	GLU
1	A	32	ASP
1	A	87	LEU
1	A	89	ARG
1	A	94	LEU
1	A	101	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	104	ARG
1	A	106	TYR
1	A	109	ASP
1	A	135	ARG
1	A	141	LEU
1	A	144	LEU
1	A	152	THR
1	A	156	LEU
1	A	157	VAL
1	A	182	THR
1	A	201	LEU
1	A	204	LEU
1	A	222	ILE
1	A	235	LYS
1	A	249	LEU
2	B	2361	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	47	GLN
1	A	52	ASN
1	A	192	ASN
1	A	195	ASN
1	A	216	ASN
1	A	228	ASN
1	A	230	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	251/279 (89%)	0.89	38 (15%) <span style="border: 1px solid red; padding: 2px;">2</span> <span style="border: 1px solid red; padding: 2px;">3</span>	27, 54, 76, 89	0
2	B	21/25 (84%)	0.99	2 (9%) <span style="border: 1px solid red; padding: 2px;">8</span> <span style="border: 1px solid red; padding: 2px;">10</span>	23, 48, 62, 64	0
All	All	272/304 (89%)	0.90	40 (14%) <span style="border: 1px solid red; padding: 2px;">2</span> <span style="border: 1px solid red; padding: 2px;">3</span>	23, 54, 76, 89	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	2365	ALA	5.5
1	A	107	LEU	5.1
1	A	31	VAL	4.8
1	A	174	ILE	4.4
1	A	28	GLU	4.2
1	A	218	LYS	4.1
1	A	247	LEU	4.0
1	A	243	ILE	3.7
1	A	32	ASP	3.5
1	A	204	LEU	3.5
1	A	0	MET	3.5
1	A	29	GLY	3.4
1	A	181	LEU	3.2
1	A	210	ILE	3.1
1	A	219	SER	3.0
1	A	34	LYS	2.9
1	A	240	ILE	2.9
1	A	163	LEU	2.8
1	A	190	LEU	2.8
1	A	211	PHE	2.7
1	A	27	GLU	2.5
1	A	39	LEU	2.5
1	A	90	ALA	2.5
1	A	250	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	223	GLU	2.4
1	A	114	ILE	2.4
1	A	33	GLY	2.4
1	A	208	MET	2.4
1	A	1	PRO	2.4
1	A	217	THR	2.3
2	B	2364	SER	2.3
1	A	170	MET	2.3
1	A	135	ARG	2.3
1	A	99	TYR	2.2
1	A	220	GLN	2.1
1	A	108	ILE	2.1
1	A	249	LEU	2.1
1	A	80	VAL	2.1
1	A	71	ARG	2.0
1	A	26	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.