

Full wwPDB X-ray Structure Validation Report (i)

May 13, 2020 – 07:08 pm BST

PDB ID : 5ZXD

Title : Crystal structure of ATP-bound human ABCF1

Authors : Qu, L.; Jiang, Y.; Liu, Z.J.

Deposited on : 2018-05-18

Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Mol Probity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	546	75%	10% •	15%
1	В	546	79%	11%	10%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP-binding cassette sub-family F member 1.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
1	A	464	Total 3713	C					0	5	0
			3713	2301	650	082	1	13			
1	R	491	Total	$^{\mathrm{C}}$	N	Ο	\mathbf{S}	Se	0	9	0
1	ם	491	3917	2479	693	727	7	11		2	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	544	HIS	_	expression tag	UNP Q8NE71
A	545	HIS	-	expression tag	UNP Q8NE71
A	546	HIS	-	expression tag	UNP Q8NE71
A	547	HIS	-	expression tag	UNP Q8NE71
В	544	HIS	-	expression tag	UNP Q8NE71
В	545	HIS	-	expression tag	UNP Q8NE71
В	546	HIS	-	expression tag	UNP Q8NE71
В	547	HIS	-	expression tag	UNP Q8NE71

• Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	Р	0	0	
2	Α	1	31	10	5	13	3	U	0	
2	Λ	1	Total C N O F		Р	0	0			
2	Α	1	31	10	5	13	3	U		
9	D	1	Total	С	N	О	Р	0	0	
2	2 B	1	31	10	5	13	3	U	U	
9	0 D	1	Total	С	N	О	Р	0	0	
$\begin{vmatrix} 2 & B \end{vmatrix}$		31	10	5	13	3	U	0		

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	$\mathbf{AltConf}$
3	A	186	Total O 186 186	0	0
3	В	123	Total O 123 123	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-binding cassette sub-family F member 1





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	66.07Å 132.81Å 148.33Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.42 - 2.29	Depositor
Resolution (A)	54.95 - 2.29	EDS
% Data completeness	98.3 (42.42-2.29)	Depositor
(in resolution range)	95.9 (54.95-2.29)	EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.15 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.13-2998	Depositor
D D.	0.194 , 0.224	Depositor
R, R_{free}	0.194 , 0.224	DCC
R_{free} test set	2003 reflections (3.41%)	wwPDB-VP
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.308	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 51.4	EDS
L-test for twinning ²	$ < L > = 0.48, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8063	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.27	0/3777	0.45	0/5082	
1	В	0.26	0/3983	0.45	0/5366	
All	All	0.27	0/7760	0.45	0/10448	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3713	0	3745	42	1
1	В	3917	0	3919	51	1
2	A	62	0	24	2	0
2	В	62	0	24	4	0
3	A	186	0	0	2	0
3	В	123	0	0	3	0
All	All	8063	0	7712	93	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

		Interatomic	Clash
Atom-1	Atom-2	${ m distance} \; ({ m \AA})$	overlap (Å)
1:B:94:GLU:CG	1:B:98:ARG:HH12	1.34	1.39
1:B:94:GLU:HG3	1:B:98:ARG:NH1	1.42	1.32
1:A:71:VAL:HG23	1:A:152:GLN:HA	1.31	1.07
1:A:66:LEU:HD11	1:A:164:LEU:HA	1.42	1.02
1:A:308:LYS:NZ	3:A:701:HOH:O	2.09	0.86
1:A:71:VAL:CG2	1:A:152:GLN:HA	2.08	0.83
1:B:94:GLU:HG3	1:B:98:ARG:HH12	0.63	0.78
1:B:94:GLU:OE2	1:B:98:ARG:NH1	2.18	0.76
1:B:94:GLU:CD	1:B:98:ARG:HH12	1.89	0.75
1:A:80:GLN:HE22	1:A:166:ARG:HB2	1.50	0.74
1:A:227:ARG:NH1	3:A:702:HOH:O	2.19	0.73
1:B:94:GLU:HG3	1:B:98:ARG:CZ	2.17	0.72
1:B:407:GLU:OE1	1:B:415:ARG:NH1	2.22	0.72
1:B:94:GLU:CG	1:B:98:ARG:NH1	2.19	0.72
1:B:94:GLU:CD	1:B:98:ARG:NH1	2.44	0.71
2:B:601:ATP:O2G	3:B:701:HOH:O	2.08	0.70
1:B:322:LEU:HD23	1:B:352:MSE:HE3	1.75	0.69
1:B:8:LEU:HD11	1:B:56:LEU:HB3	1.75	0.68
1:A:149:ARG:HD3	1:A:153:LYS:HB2	1.77	0.67
1:A:66:LEU:HD12	1:A:167:ALA:HB3	1.78	0.65
1:A:367:SER:OG	1:A:469:ASP:OD2	2.12	0.64
1:B:98:ARG:NH2	3:B:704:HOH:O	2.28	0.64
1:A:75:GLU:OE2	1:A:75:GLU:N	2.16	0.64
1:A:80:GLN:NE2	1:A:80:GLN:HA	2.13	0.63
1:A:244:GLN:HA	1:A:247:LYS:HE2	1.80	0.62
1:A:77:PRO:HD3	1:A:147:GLN:O	1.99	0.62
1:A:72:VAL:H	1:A:151:THR:HG23	1.65	0.61
1:B:358:ILE:HB	1:B:497:VAL:HG22	1.82	0.60
1:B:383:GLU:OE1	1:B:385:ARG:NE	2.33	0.59
1:A:62:ILE:HG23	1:A:173:THR:H	1.67	0.59
1:B:32:ARG:NE	1:B:232:ARG:HH21	2.00	0.59
1:B:245:LYS:HA	1:B:245:LYS:HE3	1.86	0.58
1:A:65:LEU:HB2	1:A:176:MSE:HE3	1.84	0.57
1:B:333:THR:HB	1:B:381:HIS:HB2	1.86	0.57
1:B:481:ASP:O	1:B:485:GLU:HG3	2.05	0.57
1:A:498:SER:HB3	1:A:504:ILE:HD11	1.86	0.56
1:A:71:VAL:HG22	1:A:152:GLN:HG2	1.85	0.56
1:A:72:VAL:H	1:A:151:THR:CG2	2.20	0.55
1:A:71:VAL:HA	1:A:151:THR:HG23	1.88	0.54
1:B:45:THR:OG1	2:B:601:ATP:O2G	2.26	0.54
1:A:80:GLN:NE2	1:A:166:ARG:HB2	2.21	0.53



Continued from previous page...

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ ({\rm \AA})$	$overlap(\AA)$
1:B:350:ILE:HD13	1:B:495:ILE:HD11	1.90	0.53
1:A:72:VAL:O	1:A:152:GLN:HG3	2.08	0.53
1:A:367:SER:OG	2:A:602:ATP:O1G	2.27	0.52
1:B:367:SER:OG	1:B:469:ASP:OD2	2.20	0.52
1:B:71:VAL:CG1	1:B:159:ARG:HG2	2.39	0.52
1:A:350:ILE:HG21	1:A:495:ILE:HD11	1.91	0.51
1:B:350:ILE:HG21	1:B:495:ILE:HD11	1.94	0.49
1:B:6:ILE:HB	1:B:28:ILE:HB	1.94	0.49
1:B:176:MSE:SE	1:B:204:LEU:HD23	2.62	0.49
1:A:214:ASP:HA	1:A:234:ASN:HB2	1.93	0.49
1:A:66:LEU:HD12	1:A:167:ALA:CB	2.42	0.49
1:B:390:LEU:HD11	1:B:465:VAL:HG23	1.94	0.49
1:B:75:GLU:H	1:B:75:GLU:CD	2.15	0.49
1:A:244:GLN:O	1:A:247:LYS:HG2	2.13	0.48
1:B:175:LEU:HG	1:B:177:LEU:HD13	1.96	0.48
1:B:367:SER:OG	2:B:602:ATP:O1G	2.31	0.48
1:A:359:VAL:HG11	1:A:511:LEU:HD23	1.96	0.47
1:B:468:LEU:HB2	1:B:496:VAL:HG23	1.96	0.47
1:B:19:GLU:HG2	1:B:226:GLN:OE1	2.15	0.46
1:A:45:THR:OG1	1:A:178:ASP:OD2	2.22	0.46
1:A:80:GLN:OE1	1:A:162:VAL:HG13	2.16	0.46
1:B:231:TYR:CD2	1:B:238:PHE:HB2	2.51	0.46
1:B:45:THR:OG1	1:B:178:ASP:OD2	2.24	0.45
1:A:186:LEU:HD21	1:A:526:PHE:HE1	1.81	0.45
1:B:113:ARG:NH1	3:B:717:HOH:O	2.49	0.45
1:A:176:MSE:HG2	1:A:204:LEU:HB3	1.99	0.45
1:A:510:GLN:NE2	1:A:521:GLN:OE1	2.50	0.44
1:B:71:VAL:HG12	1:B:159:ARG:HG2	1.99	0.44
1:B:175:LEU:HG	1:B:177:LEU:CD1	2.47	0.43
1:B:19:GLU:O	1:B:19:GLU:HG2	2.19	0.43
1:A:358:ILE:HB	1:A:497:VAL:HG22	2.01	0.42
1:B:421:TYR:O	1:B:425:ARG:HG3	2.19	0.42
1:A:244:GLN:OE1	1:B:415:ARG:NH1	2.52	0.42
1:B:94:GLU:HG3	1:B:98:ARG:NH2	2.34	0.42
1:A:258:LYS:HE2	1:A:262:GLU:OE2	2.18	0.42
1:B:351:ASP:O	1:B:493:ALA:HB2	2.19	0.42
1:B:16:HIS:ND1	2:B:601:ATP:C4	2.88	0.42
1:B:71:VAL:HG12	1:B:159:ARG:HB3	2.01	0.42
1:A:448:GLY:HA2	1:A:474:ASN:HB3	2.01	0.42
1:A:200:ARG:HA	1:A:200:ARG:HD3	1.83	0.41
1:B:528:ASP:OD1	1:B:531:ARG:NH2	2.52	0.41



$\alpha \cdots \tau$	e	•	
Continued	trom	meaningile	maaa
-	110116	piculuas	puyc

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} & (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:B:71:VAL:HG12	1:B:159:ARG:CB	2.50	0.41
1:A:405:MSE:HB3	1:A:443:CYS:HB3	2.02	0.41
1:A:45:THR:OG1	2:A:601:ATP:O2G	2.38	0.41
1:B:224:ASP:OD2	1:B:225:ALA:N	2.54	0.41
1:B:251:LYS:HE2	1:B:251:LYS:HB3	1.59	0.41
1:A:69:GLN:HE21	1:A:69:GLN:HB3	1.72	0.41
1:B:65:LEU:HB3	1:B:172:PRO:HG3	2.03	0.41
1:A:256:GLN:OE1	1:A:304:LEU:HD12	2.21	0.41
1:B:29:VAL:HB	1:B:32:ARG:HD2	2.01	0.41
1:B:245:LYS:HA	1:B:245:LYS:CE	2.45	0.40
1:B:235:TYR:OH	1:B:477:ILE:HG21	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$	
1:A:159:ARG:NH2	1:B:9:GLU:O[3_455]	2.15	0.05	

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	463/546 (85%)	454 (98%)	8 (2%)	1 (0%)	47	57
1	В	489/546~(90%)	479 (98%)	10 (2%)	0	100	100
All	All	952/1092 (87%)	933 (98%)	18 (2%)	1 (0%)	51	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	VAL



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	400/459 (87%)	393 (98%)	7 (2%)	59 72
1	В	420/459 (92%)	418 (100%)	2 (0%)	88 94
All	All	820/918 (89%)	811 (99%)	9 (1%)	76 84

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	SER
1	A	69	GLN
1	A	160[A]	MSE
1	A	160[B]	MSE
1	A	251	LYS
1	A	308	LYS
1	A	466	LEU
1	В	183	HIS
1	В	232	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	80	GLN
1	A	152	GLN
1	В	345	ASN
1	В	397	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	$\mid \# Z > 2$	Counts	RMSZ	# Z > 2
2	ATP	A	601	-	26,33,33	0.92	1 (3%)	31,52,52	1.32	4 (12%)
2	ATP	В	602	-	26,33,33	0.91	1 (3%)	31,52,52	1.40	4 (12%)
2	ATP	A	602	-	26,33,33	0.91	1 (3%)	31,52,52	1.44	4 (12%)
2	ATP	В	601	-	26,33,33	0.93	1 (3%)	31,52,52	1.38	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	601	-	-	3/18/38/38	0/3/3/3
2	ATP	В	602	-	-	6/18/38/38	0/3/3/3
2	ATP	A	602	-	-	6/18/38/38	0/3/3/3
2	ATP	В	601	-	-	6/18/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	${ m Observed}({ m \AA})$	$\mathbf{Ideal}(\mathbf{\AA})$
2	В	601	ATP	C5-C4	2.33	1.47	1.40



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	Z	${ m Observed}(m \AA)$	$Ideal(\AA)$
2	A	601	ATP	C5-C4	2.33	1.47	1.40
2	В	602	ATP	C5-C4	2.32	1.47	1.40
2	A	602	ATP	C5-C4	2.32	1.47	1.40

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	В	602	ATP	N3-C2-N1	-3.55	123.12	128.68
2	A	602	ATP	N3-C2-N1	-3.55	123.12	128.68
2	В	601	ATP	N3-C2-N1	-3.51	123.19	128.68
2	A	601	ATP	N3-C2-N1	-3.51	123.19	128.68
2	A	602	ATP	PB-O3B-PG	-3.25	121.69	132.83
2	В	602	ATP	PB-O3B-PG	-2.88	122.93	132.83
2	A	602	ATP	C4-C5-N7	-2.65	106.64	109.40
2	В	602	ATP	C4-C5-N7	-2.62	106.67	109.40
2	A	601	ATP	C4-C5-N7	-2.59	106.70	109.40
2	В	601	ATP	PB-O3B-PG	-2.56	124.06	132.83
2	В	601	ATP	C4-C5-N7	-2.52	106.77	109.40
2	A	602	ATP	PA-O3A-PB	-2.37	124.71	132.83
2	В	602	ATP	PA-O3A-PB	-2.36	124.72	132.83
2	A	601	ATP	PB-O3B-PG	-2.30	124.94	132.83
2	A	601	ATP	PA-O3A-PB	-2.24	125.13	132.83
2	В	601	ATP	C2-N1-C6	2.03	122.22	118.75

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	602	ATP	C5'-O5'-PA-O1A
2	В	602	ATP	C5'-O5'-PA-O3A
2	A	602	ATP	C5'-O5'-PA-O1A
2	A	602	ATP	C5'-O5'-PA-O3A
2	В	601	ATP	C5'-O5'-PA-O2A
2	В	602	ATP	PB-O3B-PG-O1G
2	A	601	ATP	PA-O3A-PB-O3B
2	В	601	ATP	PA-O3A-PB-O3B
2	A	602	ATP	PB-O3B-PG-O1G
2	В	602	ATP	PB-O3B-PG-O2G
2	В	602	ATP	PG-O3B-PB-O2B
2	A	602	ATP	PA-O3A-PB-O1B
2	В	601	ATP	PA-O3A-PB-O1B
2	В	601	ATP	C5'-O5'-PA-O3A



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	601	ATP	PA-O3A-PB-O1B
2	A	601	ATP	PA-O3A-PB-O2B
2	A	602	ATP	PG-O3B-PB-O2B
2	В	601	ATP	PA-O3A-PB-O2B
2	В	602	ATP	C5'-O5'-PA-O2A
2	A	602	ATP	C5'-O5'-PA-O2A
2	В	601	ATP	C5'-O5'-PA-O1A


There are no ring outliers.

4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	ATP	1	0
2	В	602	ATP	1	0
2	A	602	ATP	1	0
2	В	601	ATP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	454/546 (83%)	-0.05	17 (3%) 41 47	20, 39, 76, 105	0
1	В	481/546 (88%)	0.03	13 (2%) 54 60	29, 48, 79, 115	0
All	All	935/1092~(85%)	-0.01	30 (3%) 47 53	20, 45, 78, 115	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	67	CYS	5.1
1	В	250	LEU	4.6
1	A	82	VAL	4.2
1	В	249	LEU	4.0
1	A	70	GLU	3.6
1	A	403	LEU	3.6
1	В	71	VAL	3.5
1	A	400	ALA	3.4
1	A	80	GLN	3.4
1	В	246	GLN	3.3
1	В	251	LYS	3.0
1	В	252	GLN	2.9
1	A	2	ASN	2.8
1	В	183	HIS	2.8
1	A	309	GLU	2.7
1	A	69	GLN	2.7
1	A	402	GLN	2.6
1	В	3	ALA	2.6
1	В	248	GLU	2.6
1	A	302	GLU	2.5
1	A	127	ALA	2.4
1	A	76	THR	2.4
1	A	310	TYR	2.3
1	A	523	ASP	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ	
1	A	125	GLY	2.2	
1	В	72	VAL	2.2	
1	В	337	GLN	2.2	
1	В	151	THR	2.2	
1	В	75	GLU	2.1	
1	A	308	LYS	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	ATP	В	601	31/31	0.94	0.14	38,74,84,85	0
2	ATP	A	601	31/31	0.95	0.15	39,61,71,73	0
2	ATP	A	602	31/31	0.95	0.12	30,42,56,67	0
2	ATP	В	602	31/31	0.95	0.12	39,51,65,72	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ATP B 601:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ATP A 601:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around ATP A 602:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ATP B 602:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

