



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2024 – 12:25 PM JST

PDB ID : 5ZXH  
Title : The structure of MT189-tubulin complex  
Authors : Li, Z.P.; Wang, Y.X.; Meng, T.; Yang, J.L.; Chen, Q.  
Deposited on : 2018-05-21  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	93% . .
1	C	450	91% 7% .
2	B	445	87% 7% . 5%
2	D	445	5% 89% 6% 5%
3	E	143	83% . 16%
4	F	384	5% 76% . 20%

## 2 Entry composition i

There are 12 unique types of molecules in this entry. The entry contains 17367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	437	Total	C	N	O	S	0	4	0
			3441	2179	586	652	24			
1	C	440	Total	C	N	O	S	0	7	0
			3469	2193	588	663	25			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	424	Total	C	N	O	S	0	2	0
			3351	2106	574	645	26			
2	D	421	Total	C	N	O	S	0	1	0
			3304	2078	562	638	26			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	170	VAL	MET	engineered mutation	UNP Q6B856
B	316	VAL	ILE	engineered mutation	UNP Q6B856
D	170	VAL	MET	engineered mutation	UNP Q6B856
D	316	VAL	ILE	engineered mutation	UNP Q6B856

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	120	Total	C	N	O	S	0	0	0
			994	614	180	195	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	-	expression tag	UNP P63043
E	4	ALA	-	expression tag	UNP P63043


- Molecule 4 is a protein called Tubulin tyrosine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	F	309	2553	1645	438	455	15	0	3	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	32	10	5	14	3	0	0
5	C	1	32	10	5	14	3	0	0
5	D	1	32	10	5	14	3	0	0


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Mg 1 1	0	0
6	B	1	Total Mg 1 1	0	0
6	C	1	Total Mg 1 1	0	0
6	D	1	Total Mg 1 1	0	0

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Ca 1 1	0	0
7	B	1	Total Ca 1 1	0	0
7	C	1	Total Ca 1 1	0	0


- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C N O P 28 10 5 11 2	0	0


- Molecule 9 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES)

(formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	S			
9	B	1	Total	12	6	1	4	1	0	0
9	B	1	Total	12	6	1	4	1	0	0

- Molecule 10 is 2-(6-fluoro-3-[[[(4-methoxyphenyl)methyl]amino]imidazo[1,2-a]pyridin-2-yl]phenol (three-letter code: 9LX) (formula: C<sub>21</sub>H<sub>18</sub>FN<sub>3</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
10	B	1	27	21	1	3	2	0	0
10	D	1	27	21	1	3	2	0	0

- Molecule 11 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula:  $C_{11}H_{18}N_5O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
11	F	1	31	11	5	12	3	0	0

- Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	7	Total 7 7	0	0
12	B	4	Total 4 4	0	0
12	C	4	Total 4 4	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain

Chain A:  93%





- Molecule 1: Tubulin alpha-1B chain

Chain C:  91% 7%





- Molecule 2: Tubulin beta-2B chain

Chain B:  87% 7% 5%



- Molecule 2: Tubulin beta-2B chain

Chain D:  89% 6% 5%







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.91Å 157.06Å 182.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 46.84 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-2.80) 99.5 (46.84-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.40 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.208 , 0.259 0.210 , 0.254	Depositor DCC
$R_{free}$ test set	3753 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.8	Xtrriage
Anisotropy	0.062	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 29.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	17367	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MES, 9LX, CA, GDP, GTP, ACP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/3525	0.73	0/4784
1	C	0.55	0/3556	0.74	4/4828 (0.1%)
2	B	0.53	0/3425	0.72	0/4639
2	D	0.49	0/3377	0.68	0/4576
3	E	0.52	0/1002	0.68	0/1329
4	F	0.49	0/2618	0.65	0/3537
All	All	0.52	0/17503	0.71	4/23693 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	39	ASP	CB-CG-OD1	5.69	123.42	118.30
1	C	243	ARG	NE-CZ-NH1	5.33	122.96	120.30
1	C	79	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	C	422	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3441	0	3362	15	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3469	0	3377	13	0
2	B	3351	0	3226	17	0
2	D	3304	0	3179	10	0
3	E	994	0	1013	1	0
4	F	2553	0	2519	6	0
5	A	32	0	12	0	0
5	C	32	0	12	0	0
5	D	32	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
8	B	28	0	12	0	0
9	B	24	0	26	1	0
10	B	27	0	0	0	0
10	D	27	0	0	0	0
11	F	31	0	14	1	0
12	A	7	0	0	0	0
12	B	4	0	0	0	0
12	C	4	0	0	0	0
All	All	17367	0	16764	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229[A]:ARG:CG	1:A:229[A]:ARG:HH11	1.83	0.90
1:A:229[A]:ARG:HH11	1:A:229[A]:ARG:HG2	1.52	0.74
1:A:174:ALA:O	1:A:178:SER:HB2	1.96	0.66
1:A:229[A]:ARG:HH11	1:A:229[A]:ARG:HG3	1.60	0.66
2:B:251:ARG:NH1	9:B:503:MES:O2S	2.27	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/450 (98%)	429 (98%)	10 (2%)	0	100	100
1	C	444/450 (99%)	434 (98%)	9 (2%)	1 (0%)	47	78
2	B	422/445 (95%)	404 (96%)	15 (4%)	3 (1%)	22	53
2	D	417/445 (94%)	393 (94%)	20 (5%)	4 (1%)	15	44
3	E	116/143 (81%)	109 (94%)	7 (6%)	0	100	100
4	F	300/384 (78%)	289 (96%)	10 (3%)	1 (0%)	41	72
All	All	2138/2317 (92%)	2058 (96%)	71 (3%)	9 (0%)	34	66

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	243	PRO
2	D	39	ASP
2	D	248	ALA
2	D	392	LYS
2	B	245	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/378 (98%)	362 (97%)	10 (3%)	44	78
1	C	377/378 (100%)	368 (98%)	9 (2%)	49	81
2	B	367/383 (96%)	358 (98%)	9 (2%)	47	80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	362/383 (94%)	356 (98%)	6 (2%)	60	87
3	E	108/127 (85%)	107 (99%)	1 (1%)	78	94
4	F	281/342 (82%)	275 (98%)	6 (2%)	53	84
All	All	1867/1991 (94%)	1826 (98%)	41 (2%)	53	83

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	2	ARG
4	F	1	MET
2	D	137	HIS
2	D	253	LEU
4	F	87	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	133	GLN
1	C	249	ASN
4	F	252	ASN
1	C	372	GLN
2	B	83	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 7 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
9	MES	B	505	-	12,12,12	2.10	1 (8%)	14,16,16	6.37	9 (64%)
10	9LX	D	503	-	26,30,30	2.07	2 (7%)	27,42,42	1.26	1 (3%)
5	GTP	D	501	6	26,34,34	0.88	1 (3%)	32,54,54	1.47	9 (28%)
10	9LX	B	506	-	26,30,30	2.53	4 (15%)	27,42,42	1.15	2 (7%)
11	ACP	F	401	-	27,33,33	1.78	6 (22%)	32,52,52	1.34	3 (9%)
8	GDP	B	501	6	24,30,30	1.00	1 (4%)	30,47,47	1.04	3 (10%)
5	GTP	A	501	6	26,34,34	0.97	1 (3%)	32,54,54	1.45	4 (12%)
9	MES	B	503	-	12,12,12	2.21	1 (8%)	14,16,16	6.68	9 (64%)
5	GTP	C	501	6	26,34,34	0.94	1 (3%)	32,54,54	1.26	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	MES	B	505	-	-	2/6/14/14	0/1/1/1
10	9LX	D	503	-	-	4/8/11/11	0/4/4/4
5	GTP	D	501	6	-	5/18/38/38	0/3/3/3
10	9LX	B	506	-	-	5/8/11/11	0/4/4/4
11	ACP	F	401	-	-	6/15/38/38	0/3/3/3
8	GDP	B	501	6	-	3/12/32/32	0/3/3/3
5	GTP	A	501	6	-	5/18/38/38	0/3/3/3
9	MES	B	503	-	-	3/6/14/14	0/1/1/1
5	GTP	C	501	6	-	6/18/38/38	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	506	9LX	C11-C10	-11.01	1.38	1.49
10	D	503	9LX	C11-C10	-8.92	1.40	1.49
9	B	503	MES	C8-S	-7.07	1.67	1.77
9	B	505	MES	C8-S	-6.76	1.67	1.77
11	F	401	ACP	PG-O1G	5.53	1.61	1.50

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	B	505	MES	O3S-S-O1S	-12.92	79.72	111.27
9	B	503	MES	O3S-S-C8	-12.09	86.22	105.77
9	B	503	MES	O3S-S-O2S	-11.40	83.42	111.27
9	B	503	MES	O3S-S-O1S	-11.13	84.07	111.27
9	B	505	MES	O3S-S-O2S	-11.12	84.11	111.27

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O2G
5	A	501	GTP	C5'-O5'-PA-O1A
5	C	501	GTP	PB-O3B-PG-O3G
5	C	501	GTP	C5'-O5'-PA-O1A
5	C	501	GTP	C5'-O5'-PA-O2A

There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	F	401	ACP	1	0
9	B	503	MES	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient





equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	437/450 (97%)	-0.21	2 (0%) 91 88	19, 39, 66, 81	0
1	C	440/450 (97%)	-0.35	0 100 100	13, 29, 55, 73	1 (0%)
2	B	424/445 (95%)	-0.24	5 (1%) 79 73	17, 36, 67, 89	1 (0%)
2	D	421/445 (94%)	0.24	23 (5%) 25 16	22, 55, 89, 113	4 (0%)
3	E	120/143 (83%)	0.26	2 (1%) 70 63	26, 54, 91, 120	0
4	F	309/384 (80%)	0.24	21 (6%) 17 10	27, 59, 100, 118	0
All	All	2151/2317 (92%)	-0.07	53 (2%) 57 47	13, 42, 81, 120	6 (0%)

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	233	PHE	5.4
2	B	55	THR	4.0
4	F	161	LEU	3.9
4	F	182	ILE	3.9
4	F	232	ASN	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	MG	B	502	1/1	0.82	0.35	37,37,37,37	0
11	ACP	F	401	31/31	0.84	0.29	76,86,136,142	0
7	CA	B	504	1/1	0.89	0.15	82,82,82,82	0
9	MES	B	505	12/12	0.92	0.25	66,72,81,81	0
9	MES	B	503	12/12	0.93	0.18	66,71,73,80	0
6	MG	D	502	1/1	0.93	0.11	42,42,42,42	0
7	CA	C	503	1/1	0.93	0.09	56,56,56,56	0
10	9LX	D	503	27/27	0.94	0.18	40,52,56,58	0
7	CA	A	503	1/1	0.94	0.05	62,62,62,62	0
5	GTP	D	501	32/32	0.95	0.13	39,45,50,58	0
10	9LX	B	506	27/27	0.96	0.17	28,29,33,34	0
8	GDP	B	501	28/28	0.97	0.15	17,22,25,28	0
6	MG	A	502	1/1	0.97	0.14	25,25,25,25	0
5	GTP	A	501	32/32	0.98	0.17	19,22,24,24	0
6	MG	C	502	1/1	0.98	0.12	17,17,17,17	0
5	GTP	C	501	32/32	0.98	0.14	18,20,23,24	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ACP F 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 9LX D 503:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









**Electron density around GTP D 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)


**Electron density around 9LX B 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









## 6.5 Other polymers [i](#)

There are no such residues in this entry.