



# wwPDB X-ray Structure Validation Summary Report

Jun 22, 2022 – 01:25 pm BST

PDB ID : 7ZXG  
Title : Pfs48/45 bound to Fab fragment of monoclonal antibody 10D8  
Authors : Ko, K.T.; Lennartz, F.L.; Higgins, M.K.  
Deposited on : 2022-05-20  
Resolution : 4.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1005 (4.62-3.78)
Clashscore	141614	1044 (4.60-3.80)
Ramachandran outliers	138981	1000 (4.60-3.80)
Sidechain outliers	138945	1007 (4.62-3.78)
RSRZ outliers	127900	1063 (4.70-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	428	
1	D	428	
2	B	466	
2	E	466	
3	C	240	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	240	 % 68% 18% 13%
4	G	4	 50% 50%
5	H	2	 100%
5	I	2	 100%
5	K	2	 50% 50%
6	J	3	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	K	2	-	-	-	X
7	NAG	A	502	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 12599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gametocyte surface protein P45/48.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	363	Total 2942	C 1888	N 469	O 568	S 17	0	0	0
1	D	352	Total 2861	C 1842	N 453	O 549	S 17	0	0	0

- Molecule 2 is a protein called 10D8 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	218	Total 1652	C 1039	N 279	O 324	S 10	0	0	0
2	E	215	Total 1626	C 1021	N 274	O 321	S 10	0	0	0

- Molecule 3 is a protein called 10D8 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	217	Total 1690	C 1061	N 278	O 344	S 7	0	0	0
3	F	209	Total 1627	C 1026	N 265	O 329	S 7	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	4	Total 50	C 28	N 2	O 20	0	0	0

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	H	2	28	16	2	10	0	0	0
5	I	2	28	16	2	10	0	0	0
5	K	2	28	16	2	10	0	0	0

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	J	3	39	22	2	15	0	0	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>				<b>ZeroOcc</b>	<b>AltConf</b>
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		









- Molecule 4: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  



- Molecule 6: beta-D-mannopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.77Å 125.80Å 216.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.36 – 4.20 65.36 – 4.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (65.36-4.20) 99.8 (65.36-4.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.14 (at 4.14Å)	Xtrriage
Refinement program	PHENIX 1.19_4092, PHENIX 1.19_4092	Depositor
R, $R_{free}$	0.259 , 0.276 0.257 , 0.274	Depositor DCC
$R_{free}$ test set	1691 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	171.1	Xtrriage
Anisotropy	0.151	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12599	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	212.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/3007	0.47	0/4064
1	D	0.25	0/2924	0.47	0/3950
2	B	0.25	0/1694	0.52	0/2312
2	E	0.25	0/1667	0.51	0/2275
3	C	0.25	0/1731	0.49	0/2350
3	F	0.25	0/1667	0.49	0/2263
All	All	0.25	0/12690	0.49	0/17214

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2942	0	2866	38	0
1	D	2861	0	2789	45	0
2	B	1652	0	1612	22	0
2	E	1626	0	1581	22	0
3	C	1690	0	1623	22	0
3	F	1627	0	1566	28	0
4	G	50	0	43	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	28	0	25	2	0
5	I	28	0	25	3	0
5	K	28	0	25	2	0
6	J	39	0	34	1	0
7	A	28	0	26	0	0
All	All	12599	0	12215	169	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 169 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:196:PHE:HB2	5:I:1:NAG:H82	1.54	0.89
2:B:11:LEU:HD23	2:B:157:PRO:HG3	1.60	0.84
1:D:234:CYS:SG	1:D:235:PHE:N	2.50	0.84
3:C:156:ILE:HD12	3:C:161:ARG:HD3	1.63	0.80
2:E:11:LEU:HD21	2:E:126:THR:HG22	1.64	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/428 (83%)	318 (90%)	37 (10%)	0	100	100
1	D	342/428 (80%)	318 (93%)	24 (7%)	0	100	100
2	B	214/466 (46%)	198 (92%)	16 (8%)	0	100	100
2	E	211/466 (45%)	193 (92%)	18 (8%)	0	100	100
3	C	215/240 (90%)	198 (92%)	17 (8%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	F	205/240 (85%)	191 (93%)	14 (7%)	0	100	100
All	All	1542/2268 (68%)	1416 (92%)	126 (8%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/402 (85%)	334 (98%)	8 (2%)	50	70
1	D	332/402 (83%)	324 (98%)	8 (2%)	49	69
2	B	189/417 (45%)	187 (99%)	2 (1%)	73	84
2	E	186/417 (45%)	184 (99%)	2 (1%)	73	84
3	C	194/214 (91%)	189 (97%)	5 (3%)	46	67
3	F	187/214 (87%)	183 (98%)	4 (2%)	53	71
All	All	1430/2066 (69%)	1401 (98%)	29 (2%)	55	73

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	190	ASP
3	F	143	ASN
1	D	197	THR
2	E	205	CYS
1	D	156	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	E	103	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	G	1	4,1	14,14,15	0.44	0	17,19,21	0.42	0
4	NAG	G	2	4	14,14,15	0.21	0	17,19,21	0.66	0
4	BMA	G	3	4	11,11,12	0.60	0	15,15,17	0.86	1 (6%)
4	MAN	G	4	4	11,11,12	0.72	0	15,15,17	1.09	2 (13%)
5	NAG	H	1	1,5	14,14,15	0.90	1 (7%)	17,19,21	1.09	1 (5%)
5	NAG	H	2	5	14,14,15	0.43	0	17,19,21	0.68	1 (5%)
5	NAG	I	1	1,5	14,14,15	0.32	0	17,19,21	1.13	2 (11%)
5	NAG	I	2	5	14,14,15	0.38	0	17,19,21	0.99	1 (5%)
6	NAG	J	1	1,6	14,14,15	0.23	0	17,19,21	0.78	0
6	NAG	J	2	6	14,14,15	0.61	1 (7%)	17,19,21	0.72	0
6	BMA	J	3	6	11,11,12	0.58	0	15,15,17	1.17	1 (6%)
5	NAG	K	1	1,5	14,14,15	0.43	0	17,19,21	1.32	1 (5%)
5	NAG	K	2	5	14,14,15	0.32	0	17,19,21	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
4	BMA	G	3	4	-	1/2/19/22	0/1/1/1
4	MAN	G	4	4	-	1/2/19/22	1/1/1/1
5	NAG	H	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	H	2	5	-	1/6/23/26	0/1/1/1
5	NAG	I	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	I	2	5	-	2/6/23/26	0/1/1/1
6	NAG	J	1	1,6	-	3/6/23/26	0/1/1/1
6	NAG	J	2	6	-	0/6/23/26	0/1/1/1
6	BMA	J	3	6	-	2/2/19/22	0/1/1/1
5	NAG	K	1	1,5	-	4/6/23/26	0/1/1/1
5	NAG	K	2	5	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	1	NAG	O5-C1	-2.33	1.40	1.43
6	J	2	NAG	O5-C1	2.07	1.47	1.43

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	K	1	NAG	C1-O5-C5	4.79	118.69	112.19
6	J	3	BMA	C1-O5-C5	3.27	116.63	112.19
4	G	4	MAN	C1-O5-C5	3.17	116.49	112.19
5	H	1	NAG	C4-C3-C2	2.82	115.15	111.02
5	I	1	NAG	O4-C4-C3	2.49	116.10	110.35

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	J	1	NAG	O5-C5-C6-O6
4	G	1	NAG	O5-C5-C6-O6
5	K	1	NAG	C8-C7-N2-C2
5	K	1	NAG	O7-C7-N2-C2
6	J	1	NAG	C4-C5-C6-O6


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	4	MAN	C1-C2-C3-C4-C5-O5


6 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	2	NAG	1	0
5	I	1	NAG	3	0
5	K	1	NAG	2	0
6	J	1	NAG	1	0
5	H	1	NAG	2	0
5	I	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	NAG	A	501	1	14,14,15	0.39	0	17,19,21	0.34	0
7	NAG	A	502	1	14,14,15	0.29	0	17,19,21	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	501	1	-	1/6/23/26	0/1/1/1
7	NAG	A	502	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	502	NAG	O5-C5-C6-O6
7	A	501	NAG	O5-C5-C6-O6
7	A	502	NAG	C4-C5-C6-O6
7	A	502	NAG	C1-C2-N2-C7
7	A	502	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	363/428 (84%)	-0.27	4 (1%) 80 72	157, 224, 296, 333	0
1	D	352/428 (82%)	-0.29	0 100 100	151, 204, 306, 345	0
2	B	218/466 (46%)	-0.43	0 100 100	138, 169, 235, 248	0
2	E	215/466 (46%)	-0.18	4 (1%) 66 58	126, 182, 330, 354	0
3	C	217/240 (90%)	-0.21	1 (0%) 91 86	159, 192, 227, 242	0
3	F	209/240 (87%)	-0.23	3 (1%) 75 65	134, 229, 317, 339	0
All	All	1574/2268 (69%)	-0.27	12 (0%) 86 79	126, 203, 304, 354	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	151	LEU	5.3
2	E	150	CYS	4.4
1	A	153	GLU	3.0
1	A	154	PHE	3.0
1	A	178	ARG	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	K	2	14/15	0.54	0.45	254,300,311,314	0
5	NAG	I	2	14/15	0.62	0.38	279,291,296,298	0
4	BMA	G	3	11/12	0.78	0.26	259,262,269,271	0
5	NAG	I	1	14/15	0.80	0.22	227,279,291,292	0
5	NAG	K	1	14/15	0.81	0.21	249,267,306,314	0
5	NAG	H	2	14/15	0.83	0.27	271,278,287,289	0
6	BMA	J	3	11/12	0.83	0.67	261,262,263,267	0
4	MAN	G	4	11/12	0.84	0.40	274,278,285,290	0
6	NAG	J	2	14/15	0.87	0.32	262,265,272,274	0
6	NAG	J	1	14/15	0.88	0.27	198,218,266,268	0
4	NAG	G	1	14/15	0.91	0.20	246,255,267,268	0
5	NAG	H	1	14/15	0.92	0.15	242,257,271,272	0
4	NAG	G	2	14/15	0.95	0.17	253,264,273,275	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)













**Electron density around Chain K:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain J:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	NAG	A	501	14/15	0.52	0.26	223,256,278,279	0
7	NAG	A	502	14/15	0.66	0.43	282,286,287,287	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.