

wwPDB X-ray Structure Validation Summary Report (i)

Oct 23, 2023 – 07:52 PM EDT

PDB ID	:	2ZY4
Title	:	dodecameric L-aspartate beta-decarboxylase
Authors	:	Chen, HJ.; Ko, TP.; Lee, CY.; Wang, NC.; Wang, A.HJ.
Deposited on	:	2009-01-13
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	Λ	546	2%	150/	
	Π	040	<u>2%</u>	15%	• •
1	В	546	76%	16%	• 6%
1	C	546	3%	100/	60/
1	C	040	3%	18%	• 6%
1	D	546	79%	15%	• 6%
1	Б	F 4 C	5%		
	E	540	73%	19%	• 7%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
			4%		
1	F	546	78%	18%	••



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 27823 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	594	Total	С	Ν	0	\mathbf{S}	0	0	0
1	Л	524	4131	2630	701	782	18	0	0	0
1	В	519	Total	С	Ν	0	S	0	0	0
1	D	512	4032	2567	687	761	17	0	0	0
1	С	511	Total	С	Ν	0	S	0	0	0
		511	4019	2557	685	760	17	0	0	0
1	Л	515	Total	С	Ν	0	S	0	0	0
1	D	515	4046	2573	692	764	17	0	0	U
1	F	500	Total	С	Ν	0	S	0	0	0
1		509	4007	2551	683	756	17	0	0	0
1	Б	525	Total	С	Ν	0	S	0	0	0
I F	535	4202	2675	716	794	17	0	0		

• Molecule 1 is a protein called L-aspartate beta-decarboxylase.

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	534	LYS	-	expression tag	UNP Q93QX0
А	535	LEU	-	expression tag	UNP Q93QX0
А	536	ALA	-	expression tag	UNP Q93QX0
А	537	ALA	-	expression tag	UNP Q93QX0
А	538	ALA	-	expression tag	UNP Q93QX0
А	539	LEU	-	expression tag	UNP Q93QX0
А	540	GLU	-	expression tag	UNP Q93QX0
А	541	HIS	-	expression tag	UNP Q93QX0
А	542	HIS	-	expression tag	UNP Q93QX0
А	543	HIS	-	expression tag	UNP Q93QX0
А	544	HIS	-	expression tag	UNP Q93QX0
А	545	HIS	-	expression tag	UNP Q93QX0
А	546	HIS	-	expression tag	UNP Q93QX0
В	534	LYS	-	expression tag	UNP Q93QX0
B	535	LEU	-	expression tag	UNP Q93QX0
В	536	ALA	-	expression tag	UNP Q93QX0
В	537	ALA	-	expression tag	UNP Q93QX0



Continu	ieu jioni pre	vious puye			
Chain	Residue	Modelled	Actual	Comment	Reference
В	538	ALA	-	expression tag	UNP Q93QX0
В	539	LEU	-	expression tag	UNP Q93QX0
В	540	GLU	-	expression tag	UNP Q93QX0
В	541	HIS	-	expression tag	UNP Q93QX0
В	542	HIS	-	expression tag	UNP Q93QX0
В	543	HIS	-	expression tag	UNP Q93QX0
В	544	HIS	-	expression tag	UNP Q93QX0
В	545	HIS	-	expression tag	UNP Q93QX0
В	546	HIS	-	expression tag	UNP Q93QX0
С	534	LYS	-	expression tag	UNP Q93QX0
С	535	LEU	-	expression tag	UNP Q93QX0
С	536	ALA	-	expression tag	UNP Q93QX0
С	537	ALA	-	expression tag	UNP Q93QX0
С	538	ALA	-	expression tag	UNP Q93QX0
С	539	LEU	-	expression tag	UNP Q93QX0
С	540	GLU	-	expression tag	UNP Q93QX0
С	541	HIS	-	expression tag	UNP Q93QX0
С	542	HIS	-	expression tag	UNP Q93QX0
С	543	HIS	-	expression tag	UNP Q93QX0
С	544	HIS	-	expression tag	UNP Q93QX0
С	545	HIS	-	expression tag	UNP Q93QX0
С	546	HIS	-	expression tag	UNP Q93QX0
D	534	LYS	-	expression tag	UNP Q93QX0
D	535	LEU	-	expression tag	UNP Q93QX0
D	536	ALA	-	expression tag	UNP Q93QX0
D	537	ALA	-	expression tag	UNP Q93QX0
D	538	ALA	-	expression tag	UNP Q93QX0
D	539	LEU	-	expression tag	UNP Q93QX0
D	540	GLU	-	expression tag	UNP Q93QX0
D	541	HIS	-	expression tag	UNP Q93QX0
D	542	HIS	-	expression tag	UNP Q93QX0
D	543	HIS	-	expression tag	UNP Q93QX0
D	544	HIS	-	expression tag	UNP Q93QX0
D	545	HIS	-	expression tag	UNP Q93QX0
D	546	HIS	-	expression tag	UNP Q93QX0
Е	534	LYS	-	expression tag	UNP Q93QX0
Е	535	LEU	-	expression tag	UNP Q93QX0
Е	536	ALA	-	expression tag	UNP Q93QX0
Е	537	ALA	-	expression tag	UNP Q93QX0
Е	538	ALA	-	expression tag	UNP Q93QX0
Е	539	LEU	-	expression tag	UNP Q93QX0
Е	540	GLU	-	expression tag	UNP Q93QX0

Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
Е	541	HIS	-	expression tag	UNP Q93QX0
Е	542	HIS	-	expression tag	UNP Q93QX0
Е	543	HIS	-	expression tag	UNP Q93QX0
Е	544	HIS	-	expression tag	UNP Q93QX0
Е	545	HIS	-	expression tag	UNP Q93QX0
Е	546	HIS	-	expression tag	UNP Q93QX0
F	534	LYS	-	expression tag	UNP Q93QX0
F	535	LEU	-	expression tag	UNP Q93QX0
F	536	ALA	-	expression tag	UNP Q93QX0
F	537	ALA	-	expression tag	UNP Q93QX0
F	538	ALA	-	expression tag	UNP Q93QX0
F	539	LEU	-	expression tag	UNP Q93QX0
F	540	GLU	-	expression tag	UNP Q93QX0
F	541	HIS	-	expression tag	UNP Q93QX0
F	542	HIS	-	expression tag	UNP Q93QX0
F	543	HIS	-	expression tag	UNP Q93QX0
F	544	HIS	-	expression tag	UNP Q93QX0
F	545	HIS	-	expression tag	UNP Q93QX0
F	546	HIS	-	expression tag	UNP Q93QX0

Continued from previous page...

• Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
9	Λ	1	Total	С	Ν	0	Р	0	0	
	Z A	1	15	8	1	5	1	0	0	
0	Р	1	Total	С	Ν	0	Р	0	0	
	D	1	15	8	1	5	1	0	0	



Continued from previous page...

Mol	Chain	Residues		Ate	\mathbf{oms}			ZeroOcc	AltConf
0	С	1	Total	С	Ν	0	Р	0	0
	U	L	15	8	1	5	1	0	0
0	Л	1	Total	С	Ν	0	Р	0	0
	D	L	15	8	1	5	1	0	0
0	F	1	Total	С	Ν	0	Р	0	0
	Ľ	I	15	8	1	5	1	0	0
0	Б	1	Total	С	Ν	0	Р	0	0
	Г		15	8	1	5	1	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Cl 1 1	0	0
3	В	1	Total Cl 1 1	0	0
3	С	1	Total Cl 1 1	0	0
3	D	1	Total Cl 1 1	0	0
3	Ε	1	Total Cl 1 1	0	0
3	F	1	Total Cl 1 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	565	Total O 565 565	0	0
4	В	611	Total O 611 611	0	0
4	С	593	Total O 593 593	0	0
4	D	573	Total O 573 573	0	0
4	Е	462	Total O 462 462	0	0
4	F	486	Total O 486 486	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: L-aspartate beta-decarboxylase







S490 1191 MET M493 V348 1191 MET M493 H352 V348 1195 NR M493 H352 V348 1195 NR M493 H352 T206 S7 VR L503 H352 F204 S6 YR V508 P359 F204 S7 VR K517 Q414 W231 F14 S7 K534 H35 H35 K1 P249 F14 K53 H43 W230 F14 P236 F14 K53 H43 W230 F34 F14 F14 K53 H43 W230 F34 F14 F14 H1S H44 W233 F43 F14 F14 H1S H44 W233 F43 F14 F14 H1S H44 W233 F34 F34 F14 H1S H44 M235



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	149.95Å 217.15Å 207.15Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution(A)	30.00 - 2.00	Depositor
Resolution (A)	29.71 - 2.00	EDS
% Data completeness	94.7 (30.00-2.00)	Depositor
(in resolution range)	94.8 (29.71-2.00)	EDS
R_{merge}	0.07	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.58 (at 2.00 \text{\AA})$	Xtriage
Refinement program	CNS	Depositor
P. P.	0.157 , 0.204	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.155 , 0.203	DCC
R_{free} test set	10656 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.35 , 65.8	EDS
L-test for $twinning^2$	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	27823	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.42% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Chain		Bond lengths		Bond angles	
MIOI	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.71	0/4215	0.80	2/5705~(0.0%)
1	В	0.72	0/4114	0.80	3/5571~(0.1%)
1	С	0.70	0/4101	0.79	2/5554~(0.0%)
1	D	0.70	0/4128	0.76	1/5591~(0.0%)
1	Е	0.68	0/4089	0.78	2/5538~(0.0%)
1	F	0.70	0/4286	0.79	3/5803~(0.1%)
All	All	0.70	0/24933	0.79	13/33762~(0.0%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	С	257	PRO	N-CA-C	7.55	131.73	112.10
1	В	257	PRO	N-CA-C	7.18	130.77	112.10
1	D	257	PRO	N-CA-C	7.16	130.71	112.10
1	F	257	PRO	N-CA-C	6.68	129.48	112.10
1	Е	257	PRO	N-CA-C	6.67	129.45	112.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4131	0	4097	77	1
1	В	4032	0	4004	83	0
1	С	4019	0	3983	82	0
1	D	4046	0	4011	71	0
1	Е	4007	0	3975	91	0
1	F	4202	0	4181	90	0
2	А	15	0	6	0	0
2	В	15	0	6	0	0
2	С	15	0	7	0	0
2	D	15	0	7	0	0
2	Е	15	0	6	0	0
2	F	15	0	7	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
3	Е	1	0	0	0	0
3	F	1	0	0	0	0
4	А	565	0	0	17	0
4	В	611	0	0	20	0
4	С	593	0	0	17	1
4	D	573	0	0	19	0
4	Е	462	0	0	16	0
4	F	486	0	0	15	1
All	All	27823	0	24290	481	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 481 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:429:MET:HG3	4:A:2840:HOH:O	1.35	1.26
1:F:332:GLN:HG2	4:F:3116:HOH:O	1.54	1.08
1:E:26:SER:HB3	1:E:29:ASN:HB2	1.41	1.02
1:F:17:LYS:HE3	1:F:487:ARG:HH22	1.27	0.97
1:D:448:ASP:HB2	4:D:1323:HOH:O	1.64	0.94

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:1730:HOH:O	4:F:1730:HOH:O[4_566]	2.09	0.11
1:A:425:ARG:NH2	4:C:1033:HOH:O[4_566]	2.15	0.05

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	520/546~(95%)	509~(98%)	11 (2%)	0	100	100
1	В	510/546~(93%)	500~(98%)	9~(2%)	1 (0%)	47	44
1	С	509/546~(93%)	493 (97%)	15 (3%)	1 (0%)	47	44
1	D	513/546~(94%)	504 (98%)	8 (2%)	1 (0%)	47	44
1	Е	507/546~(93%)	481 (95%)	24~(5%)	2(0%)	34	30
1	F	533/546~(98%)	522 (98%)	10 (2%)	1 (0%)	47	44
All	All	3092/3276~(94%)	3009~(97%)	77 (2%)	6~(0%)	47	44

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
1	В	492	ASN
1	D	492	ASN
1	Е	455	GLY
1	С	492	ASN
1	F	492	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	439/455~(96%)	426~(97%)	13 (3%)	41 41
1	В	427/455~(94%)	415 (97%)	12 (3%)	43 44
1	\mathbf{C}	425/455~(93%)	415 (98%)	10 (2%)	49 51
1	D	426/455~(94%)	415 (97%)	11 (3%)	46 48
1	Ε	424/455~(93%)	411 (97%)	13 (3%)	40 40
1	F	444/455~(98%)	431 (97%)	13 (3%)	42 43
All	All	2585/2730~(95%)	2513 (97%)	72 (3%)	43 44

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	Е	333	ASN
1	F	492	ASN
1	F	41	ASN
1	F	259	SER
1	В	456	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 69 such sidechains are listed below:

Mol	Chain	\mathbf{Res}	Type
1	F	34	ASN
1	F	230	ASN
1	F	333	ASN
1	В	447	GLN
1	В	380	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Tuno Chain		Dec	Tink	Bond lengths			Bond angles		
1VIOI	Moi Type	Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	PLP	А	900	1	$15,\!15,\!16$	2.58	6 (40%)	$20,\!22,\!23$	1.87	4 (20%)
2	PLP	В	900	1	$15,\!15,\!16$	2.21	7 (46%)	$20,\!22,\!23$	1.69	5 (25%)
2	PLP	Е	900	1	15,15,16	2.83	7 (46%)	20,22,23	1.41	2 (10%)
2	PLP	С	900	1	15,15,16	2.18	7 (46%)	20,22,23	1.63	3 (15%)
2	PLP	D	900	1	15,15,16	2.26	5 (33%)	20,22,23	2.06	4 (20%)
2	PLP	F	900	1	15,15,16	2.71	6 (40%)	20,22,23	2.02	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	А	900	1	-	1/6/6/8	0/1/1/1
2	PLP	В	900	1	-	0/6/6/8	0/1/1/1
2	PLP	Е	900	1	-	0/6/6/8	0/1/1/1
2	PLP	С	900	1	-	0/6/6/8	0/1/1/1
2	PLP	D	900	1	-	0/6/6/8	0/1/1/1
2	PLP	F	900	1	-	0/6/6/8	0/1/1/1

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Е	900	PLP	C2A-C2	6.85	1.62	1.50
2	А	900	PLP	C3-C2	-6.01	1.34	1.40
2	F	900	PLP	C5-C4	5.39	1.46	1.40
2	F	900	PLP	C4A-C4	5.36	1.62	1.51
2	Е	900	PLP	C5-C4	4.59	1.45	1.40

The worst 5 of 22 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	F	900	PLP	C4A-C4-C5	6.92	128.07	120.94
2	D	900	PLP	C4A-C4-C5	5.41	126.51	120.94
2	А	900	PLP	C4A-C4-C5	5.25	126.34	120.94
2	В	900	PLP	C4A-C4-C5	5.01	126.09	120.94
2	С	900	PLP	C4A-C4-C5	4.87	125.95	120.94

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	900	PLP	C6-C5-C5A-O4P

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.





6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$OWAB(Å^2)$	Q<0.9
1	А	524/546~(95%)	-0.32	11 (2%) 63	62	10, 20, 42, 65	0
1	В	512/546~(93%)	-0.38	11 (2%) 63	62	9, 19, 43, 81	0
1	С	511/546~(93%)	-0.33	19 (3%) 41	41	9, 20, 46, 76	0
1	D	515/546~(94%)	-0.32	19 (3%) 41	41	9, 21, 43, 76	0
1	Ε	509/546~(93%)	-0.01	29 (5%) 23	23	11, 25, 57, 88	0
1	F	535/546~(97%)	-0.22	23 (4%) 35	34	9, 22, 52, 78	0
All	All	3106/3276~(94%)	-0.26	112 (3%) 42	42	9, 21, 48, 88	0

The worst 5 of 112 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	490	GLY	11.0
1	Е	490	GLY	9.3
1	С	490	GLY	8.0
1	F	491	SER	7.2
1	D	490	GLY	7.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
2	PLP	Е	900	15/16	0.98	0.09	17,21,23,24	0
3	CL	D	4004	1/1	0.98	0.06	21,21,21,21	0
3	CL	F	4006	1/1	0.98	0.05	24,24,24,24	0
2	PLP	D	900	15/16	0.99	0.11	14,16,17,18	0
2	PLP	А	900	15/16	0.99	0.08	14,15,18,18	0
2	PLP	F	900	15/16	0.99	0.08	13,15,18,18	0
3	CL	А	4001	1/1	0.99	0.05	18,18,18,18	0
3	CL	В	4002	1/1	0.99	0.04	20,20,20,20	0
3	CL	С	4003	1/1	0.99	0.04	19,19,19,19	0
2	PLP	В	900	15/16	0.99	0.08	12,15,17,18	0
3	CL	Е	4005	1/1	0.99	0.05	20,20,20,20	0
2	PLP	С	900	15/16	0.99	0.09	$13,\!15,\!17,\!17$	0

6.5 Other polymers (i)

There are no such residues in this entry.

