



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 08:44 pm GMT

PDB ID : 1A00
Title : HEMOGLOBIN (VAL BETA1 MET, TRP BETA37 TYR) MUTANT
Authors : Kavanaugh, J.S.; Arnone, A.
Deposited on : 1997-12-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	141	82% 16% .
1	C	141	86% 9% 5% .
2	B	146	82% 15% ..
2	D	146	78% 19% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4770 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMOGLOBIN (ALPHA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1069	685	187	194	3	0	0	0
1	C	141	1069	685	187	194	3	0	0	0

- Molecule 2 is a protein called HEMOGLOBIN (BETA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	146	1122	722	194	202	4	0	0	0
2	D	146	1122	722	194	202	4	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	37	TYR	TRP	ENGINEERED	UNP P68871
D	37	TYR	TRP	ENGINEERED	UNP P68871

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	52	Total	O	0	0
			52	52		
4	C	66	Total	O	0	0
			66	66		
4	D	46	Total	O	0	0
			46	46		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HEMOGLOBIN (ALPHA CHAIN)




- Molecule 1: HEMOGLOBIN (ALPHA CHAIN)



- Molecule 2: HEMOGLOBIN (BETA CHAIN)



- Molecule 2: HEMOGLOBIN (BETA CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.10Å 112.00Å 63.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00 46.61 – 1.73	Depositor EDS
% Data completeness (in resolution range)	97.1 (8.00-2.00) 67.5 (46.61-1.73)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.73Å)	Xtrriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.169 , 0.223 0.531 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	14.8	Xtrriage
Anisotropy	0.955	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 999.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.40	EDS
Total number of atoms	4770	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	0/1097	1.33	7/1491 (0.5%)
1	C	0.83	0/1097	1.50	12/1491 (0.8%)
2	B	0.84	0/1151	1.33	3/1561 (0.2%)
2	D	0.82	0/1151	1.39	8/1561 (0.5%)
All	All	0.82	0/4496	1.39	30/6104 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	92	ARG	NE-CZ-NH2	-20.75	109.92	120.30
2	D	40	ARG	NE-CZ-NH1	12.78	126.69	120.30
1	C	85	ASP	CB-CG-OD2	-8.22	110.90	118.30
1	C	85	ASP	CB-CG-OD1	7.70	125.23	118.30
2	D	66	LYS	CA-CB-CG	7.68	130.29	113.40
1	C	99	LYS	CB-CA-C	7.13	124.67	110.40
1	A	75	ASP	CB-CG-OD2	-7.09	111.92	118.30
2	D	40	ARG	CD-NE-CZ	7.08	133.51	123.60
1	C	1	VAL	CG1-CB-CG2	7.00	122.10	110.90
1	C	105	LEU	CA-CB-CG	6.76	130.86	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	92	ARG	NE-CZ-NH1	6.73	123.67	120.30
1	A	92	ARG	NE-CZ-NH2	-6.53	117.03	120.30
2	D	52	ASP	CB-CG-OD2	-6.44	112.50	118.30
1	C	92	ARG	NH1-CZ-NH2	6.37	126.41	119.40
1	A	92	ARG	NE-CZ-NH1	6.29	123.44	120.30
1	C	24	TYR	CB-CG-CD2	-6.17	117.30	121.00
2	D	2	HIS	CA-CB-CG	-6.08	103.25	113.60
1	C	99	LYS	CA-CB-CG	6.02	126.64	113.40
2	B	6	GLU	CA-CB-CG	6.00	126.59	113.40
2	B	40	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	C	106	LEU	CA-CB-CG	5.81	128.66	115.30
1	A	30	GLU	OE1-CD-OE2	5.58	130.00	123.30
1	A	23	GLU	CA-CB-CG	5.51	125.52	113.40
1	A	64	ASP	CB-CG-OD1	5.25	123.03	118.30
1	A	74	ASP	CB-CG-OD2	-5.21	113.61	118.30
2	D	52	ASP	CB-CG-OD1	5.17	122.95	118.30
2	D	2	HIS	N-CA-CB	5.16	119.88	110.60
1	C	47	ASP	O-C-N	5.15	130.94	122.70
2	D	141	LEU	CA-CB-CG	5.09	127.00	115.30
2	B	143	HIS	CA-CB-CG	-5.02	105.07	113.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	104	ARG	Sidechain
2	D	40	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	14	0
1	C	1069	0	1073	10	0
2	B	1122	0	1117	11	0
2	D	1122	0	1117	12	0
3	A	43	0	30	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	43	0	30	4	0
3	C	43	0	30	0	0
3	D	43	0	30	1	0
4	A	52	0	0	1	0
4	B	52	0	0	0	0
4	C	66	0	0	0	0
4	D	46	0	0	1	0
All	All	4770	0	4500	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:147:HEM:HBC2	3:B:147:HEM:HMC1	1.71	0.72
3:B:147:HEM:HBC2	3:B:147:HEM:CMC	2.20	0.71
1:A:96:VAL:O	1:A:99:LYS:HG2	1.96	0.66
2:D:124:PRO:HB2	2:D:125:PRO:HD3	1.80	0.63
1:C:49:SER:O	1:C:52:SER:HB3	2.03	0.58
1:A:96:VAL:HB	1:A:99:LYS:NZ	2.18	0.57
2:B:51:PRO:O	2:B:55:MET:HG2	2.07	0.55
2:B:1:MET:HG3	2:B:136:GLY:HA3	1.88	0.55
1:A:75:ASP:OD2	1:A:78:ASN:HB2	2.07	0.54
2:D:80:ASN:ND2	2:D:83:GLY:HA3	2.22	0.54
1:C:35:SER:HB3	2:D:131:GLN:HG3	1.89	0.54
2:D:143:HIS:HB3	4:D:414:HOH:O	2.07	0.53
2:B:124:PRO:HB2	2:B:125:PRO:HD3	1.89	0.53
1:A:13:ALA:O	1:A:17:VAL:HG23	2.10	0.51
1:C:90:LYS:HB3	1:C:90:LYS:NZ	2.26	0.50
2:B:6:GLU:CD	2:B:6:GLU:H	2.15	0.50
1:A:76:MET:N	1:A:77:PRO:CD	2.75	0.50
2:D:91:LEU:HD12	2:D:95:LYS:HB2	1.94	0.49
1:C:3:SER:HB2	1:C:4:PRO:HD2	1.94	0.49
1:C:90:LYS:O	1:C:92:ARG:HD3	2.13	0.49
1:A:3:SER:O	1:A:7:LYS:HG3	2.13	0.48
1:C:3:SER:HB2	1:C:4:PRO:CD	2.44	0.47
2:B:86:ALA:O	2:B:90:GLU:HG3	2.14	0.47
1:A:51:GLY:O	1:A:52:SER:C	2.51	0.47
2:B:82:LYS:HE3	2:B:143:HIS:CD2	2.50	0.46
2:D:50:THR:HB	2:D:51:PRO:HD2	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:LYS:HG3	1:A:116:GLU:HG2	1.97	0.45
3:B:147:HEM:CBC	3:B:147:HEM:HMC1	2.44	0.45
1:A:113:LEU:HB3	1:A:116:GLU:HB2	1.99	0.44
2:D:51:PRO:O	2:D:55:MET:HG2	2.18	0.44
2:B:3:LEU:HA	2:B:7:GLU:OE1	2.18	0.44
2:D:88:LEU:HD23	2:D:91:LEU:HD23	2.00	0.43
2:D:3:LEU:HD23	2:D:3:LEU:N	2.33	0.43
2:B:18:VAL:HG13	2:B:23:VAL:HG21	2.00	0.43
2:B:95:LYS:HA	2:B:95:LYS:HD3	1.72	0.43
1:A:21:ALA:HB1	1:A:63:ALA:HB1	2.01	0.43
1:C:31:ARG:HD3	2:D:127:GLN:OE1	2.19	0.43
3:D:147:HEM:HBC2	3:D:147:HEM:CMC	2.49	0.42
2:D:21:ASP:HA	2:D:65:LYS:HG3	2.00	0.42
1:C:20:HIS:HB3	1:C:24:TYR:CE1	2.55	0.42
1:C:17:VAL:HG13	1:C:24:TYR:CD2	2.55	0.42
2:D:4:THR:HB	2:D:5:PRO:HD2	2.02	0.41
1:C:43:PHE:N	1:C:44:PRO:CD	2.84	0.41
1:A:49:SER:O	1:A:52:SER:HB3	2.21	0.41
1:A:76:MET:HB2	1:A:77:PRO:HD3	2.03	0.41
1:A:96:VAL:HB	1:A:99:LYS:HZ3	1.85	0.41
1:A:4:PRO:HD2	4:A:308:HOH:O	2.21	0.40
2:B:96:LEU:HD13	3:B:147:HEM:C3D	2.57	0.40
2:B:57:ASN:HA	2:B:58:PRO:HD3	1.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	135 (97%)	4 (3%)	0	100	100
1	C	139/141 (99%)	138 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	144/146 (99%)	141 (98%)	3 (2%)	0	100	100
2	D	144/146 (99%)	140 (97%)	4 (3%)	0	100	100
All	All	566/574 (99%)	554 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	108 (96%)	5 (4%)	31	27
1	C	113/113 (100%)	105 (93%)	8 (7%)	16	11
2	B	118/118 (100%)	110 (93%)	8 (7%)	17	12
2	D	118/118 (100%)	106 (90%)	12 (10%)	8	4
All	All	462/462 (100%)	429 (93%)	33 (7%)	16	11

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LYS
1	A	16	LYS
1	A	52	SER
1	A	90	LYS
1	A	109	LEU
2	B	1	MET
2	B	6	GLU
2	B	12	THR
2	B	22	GLU
2	B	26	GLU
2	B	32	LEU
2	B	66	LYS
2	B	104	ARG
1	C	1	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	16	LYS
1	C	52	SER
1	C	85	ASP
1	C	90	LYS
1	C	92	ARG
1	C	99	LYS
1	C	105	LEU
2	D	1	MET
2	D	3	LEU
2	D	6	GLU
2	D	8	LYS
2	D	37	TYR
2	D	43	GLU
2	D	59	LYS
2	D	61	LYS
2	D	65	LYS
2	D	66	LYS
2	D	75	LEU
2	D	104	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	HIS
2	B	63	HIS
2	B	139	ASN
2	D	80	ASN
2	D	117	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	142	1	27,50,50	1.69	4 (14%)	17,82,82	1.98	6 (35%)
3	HEM	B	147	2	27,50,50	2.01	7 (25%)	17,82,82	1.88	6 (35%)
3	HEM	C	142	1	27,50,50	1.85	4 (14%)	17,82,82	1.91	7 (41%)
3	HEM	D	147	2	27,50,50	2.03	7 (25%)	17,82,82	1.56	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	142	1	-	0/6/54/54	0/0/8/8
3	HEM	B	147	2	-	0/6/54/54	0/0/8/8
3	HEM	C	142	1	-	0/6/54/54	0/0/8/8
3	HEM	D	147	2	-	0/6/54/54	0/0/8/8

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	147	HEM	C3C-C2C	-5.15	1.33	1.40
3	D	147	HEM	C3C-C2C	-5.15	1.33	1.40
3	C	142	HEM	C3C-C2C	-4.90	1.33	1.40
3	A	142	HEM	C3C-C2C	-4.39	1.34	1.40
3	C	142	HEM	C3B-C2B	-4.08	1.34	1.40
3	D	147	HEM	C3B-C2B	-4.01	1.34	1.40
3	B	147	HEM	C3B-C2B	-3.83	1.35	1.40
3	A	142	HEM	C3B-C2B	-3.53	1.35	1.40
3	D	147	HEM	CMB-C2B	2.04	1.56	1.51
3	B	147	HEM	CMC-C2C	2.07	1.56	1.51
3	B	147	HEM	CMA-C3A	2.09	1.55	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	147	HEM	CAA-C2A	2.16	1.55	1.52
3	D	147	HEM	CMA-C3A	2.46	1.56	1.51
3	B	147	HEM	C3B-CAB	2.73	1.53	1.47
3	B	147	HEM	CAA-C2A	2.90	1.56	1.52
3	A	142	HEM	C3C-CAC	3.06	1.53	1.47
3	A	142	HEM	C3B-CAB	3.07	1.54	1.47
3	C	142	HEM	C3B-CAB	3.31	1.54	1.47
3	D	147	HEM	C3C-CAC	3.55	1.54	1.47
3	C	142	HEM	C3C-CAC	3.86	1.55	1.47
3	D	147	HEM	C3B-CAB	3.91	1.55	1.47
3	B	147	HEM	C3C-CAC	4.00	1.55	1.47

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	142	HEM	CMA-C3A-C4A	-4.08	122.20	128.46
3	B	147	HEM	CMD-C2D-C1D	-3.79	122.64	128.46
3	A	142	HEM	CMD-C2D-C1D	-3.25	123.48	128.46
3	A	142	HEM	CMA-C3A-C4A	-3.23	123.50	128.46
3	D	147	HEM	CMD-C2D-C1D	-2.90	124.00	128.46
3	B	147	HEM	CMA-C3A-C4A	-2.88	124.03	128.46
3	C	142	HEM	CMD-C2D-C1D	-2.57	124.51	128.46
3	C	142	HEM	CBD-CAD-C3D	-2.29	108.11	112.47
3	D	147	HEM	CAD-CBD-CGD	-2.21	108.88	112.66
3	D	147	HEM	CMA-C3A-C4A	-2.07	125.28	128.46
3	D	147	HEM	CMD-C2D-C3D	2.02	128.75	124.94
3	B	147	HEM	CMB-C2B-C3B	2.02	128.56	124.88
3	C	142	HEM	CMC-C2C-C3C	2.15	128.79	124.88
3	D	147	HEM	CMB-C2B-C3B	2.15	128.80	124.88
3	C	142	HEM	CMD-C2D-C3D	2.20	129.10	124.94
3	A	142	HEM	CMA-C3A-C2A	2.24	129.17	124.94
3	D	147	HEM	CAA-CBA-CGA	2.30	116.59	112.66
3	C	142	HEM	CBA-CAA-C2A	2.31	116.91	112.48
3	B	147	HEM	CMA-C3A-C2A	2.39	129.44	124.94
3	B	147	HEM	C4C-C3C-C2C	2.56	108.68	106.90
3	A	142	HEM	CMB-C2B-C3B	2.57	129.56	124.88
3	A	142	HEM	CMD-C2D-C3D	2.59	129.82	124.94
3	B	147	HEM	CMD-C2D-C3D	2.85	130.31	124.94
3	C	142	HEM	CMA-C3A-C2A	3.10	130.78	124.94
3	A	142	HEM	CBA-CAA-C2A	4.22	120.56	112.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	147	HEM	4	0
3	D	147	HEM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.