



Full wwPDB X-ray Structure Validation Report i

Apr 19, 2023 – 10:47 pm BST

PDB ID : 8A4J
Title : Human GDAP1, A247V mutant
Authors : Sutinen, A.; Kursula, P.
Deposited on : 2022-06-12
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.32.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 8010 atoms, of which 4006 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ganglioside-induced differentiation-associated protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	249	Total	C 4140	H 1319	N 2073	O 360	S 379	9	0	0
1	B	232	Total	C 3859	H 1232	N 1933	O 336	S 349	9	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	247	VAL	ALA	variant	UNP Q8TB36
B	247	VAL	ALA	variant	UNP Q8TB36


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	10	Total O 10 10	0	0
2	B	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ganglioside-induced differentiation-associated protein 1



- Molecule 1: Ganglioside-induced differentiation-associated protein 1



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.29 Å 115.69 Å 115.94 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.28 – 2.68 42.28 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.6 (42.28-2.68) 99.6 (42.28-2.68)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.67 (at 2.69 Å)	Xtriage
Refinement program	PHENIX 1.20_4459	Depositor
R , R_{free}	0.252 , 0.284 0.244 , 0.277	Depositor DCC
R_{free} test set	1997 reflections (7.08%)	wwPDB-VP
Wilson B-factor (Å ²)	85.0	Xtriage
Anisotropy	0.528	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 79.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8010	wwPDB-VP
Average B, all atoms (Å ²)	127.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	3/2112 (0.1%)	0.93	0/2857
1	B	0.59	1/1968 (0.1%)	0.75	1/2663 (0.0%)
All	All	0.74	4/4080 (0.1%)	0.85	1/5520 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	76	GLU	CD-OE2	7.92	1.34	1.25
1	A	247	VAL	CB-CG1	-6.95	1.38	1.52
1	A	31	TRP	CB-CG	-5.89	1.39	1.50
1	B	31	TRP	CB-CG	-5.17	1.41	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	226	ARG	NE-CZ-NH2	-5.35	117.62	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2067	2073	2072	36	0
1	B	1926	1933	1931	36	0
2	A	10	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
All	All	4004	4006	4003	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:ASP:OD2	2:A:401:HOH:O	1.93	0.87
1:B:226:ARG:HH21	1:B:227:ASN:HB2	1.48	0.77
1:A:26:LEU:CD2	1:A:80:LEU:HD11	2.16	0.75
1:A:103:ASP:OD1	1:A:104:GLU:N	2.22	0.72
1:A:222:GLU:OE2	1:A:226:ARG:NH2	2.27	0.67
1:B:245:THR:OG1	1:B:247:VAL:HG12	1.96	0.66
1:A:26:LEU:HD22	1:A:80:LEU:HD11	1.79	0.64
1:B:251:LEU:HD23	1:B:279:TYR:CE1	2.32	0.63
1:A:66:PRO:HD2	1:A:69:MET:HG3	1.81	0.62
1:B:251:LEU:HD23	1:B:279:TYR:CZ	2.35	0.62
1:B:36:SER:HB2	1:B:90:ALA:HB2	1.83	0.60
1:B:77:VAL:HG13	1:B:77:VAL:O	2.01	0.60
1:A:34:SER:OG	1:A:37:SER:HB3	2.03	0.59
1:B:141:ILE:HD11	1:B:152:ILE:HD13	1.85	0.58
1:A:30:HIS:HB2	1:A:37:SER:HB2	1.86	0.58
1:B:219:VAL:HG12	1:B:223:LEU:HD12	1.86	0.58
1:B:226:ARG:NH1	1:B:236:GLN:NE2	2.52	0.57
1:A:66:PRO:HD2	1:A:69:MET:HE3	1.88	0.56
1:B:226:ARG:NH1	1:B:236:GLN:HE22	2.03	0.55
1:A:69:MET:HE1	1:A:155:TYR:HE1	1.71	0.54
1:B:226:ARG:NH2	1:B:227:ASN:HB2	2.21	0.54
1:A:36:SER:HB2	1:A:90:ALA:HB2	1.90	0.54
1:B:66:PRO:HD2	1:B:69:MET:HG3	1.89	0.53
1:A:66:PRO:HD2	1:A:69:MET:CG	2.39	0.53
1:A:96:TYR:O	1:A:100:THR:HG23	2.09	0.52
1:A:292:VAL:HG12	1:A:293:LEU:HD23	1.91	0.52
1:B:31:TRP:CH2	1:B:77:VAL:N	2.78	0.52
1:A:69:MET:CE	1:A:155:TYR:HE1	2.23	0.52
1:A:293:LEU:O	1:A:296:VAL:HG22	2.10	0.50
1:B:59:PRO:HB3	1:B:302:SER:HB3	1.93	0.50
1:A:133:MET:O	1:A:137:THR:HG23	2.12	0.50
1:A:26:LEU:HD23	1:A:80:LEU:HD11	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:240:CYS:SG	1:B:247:VAL:HG11	2.54	0.48
1:A:103:ASP:OD1	1:A:105:ARG:N	2.39	0.48
1:A:84:GLU:OE2	1:B:58:LEU:N	2.45	0.48
1:B:219:VAL:HG12	1:B:223:LEU:CD1	2.44	0.48
1:B:272:LYS:HG2	1:B:273:ARG:HG3	1.96	0.47
1:B:124:TYR:CE2	1:B:222:GLU:HG2	2.50	0.47
1:A:69:MET:HE1	1:A:155:TYR:CE1	2.50	0.46
1:B:220:GLU:OE1	1:B:275:ASN:HB2	2.15	0.46
1:B:255:LEU:HD11	1:B:276:LEU:HD11	1.97	0.46
1:B:135:ALA:HB1	1:B:208:ILE:CD1	2.46	0.46
1:A:96:TYR:CE1	1:A:100:THR:HG21	2.51	0.45
1:B:103:ASP:OD1	1:B:104:GLU:N	2.50	0.45
1:B:137:THR:HG22	1:B:261:LEU:HD11	1.98	0.45
1:B:38:GLN:HG2	1:B:293:LEU:CD2	2.47	0.45
1:B:82:HIS:ND1	1:B:96:TYR:OH	2.36	0.44
1:B:289:PHE:C	1:B:289:PHE:CD1	2.90	0.44
1:A:264:ALA:O	1:A:265:ARG:C	2.55	0.44
1:A:292:VAL:HG12	1:A:293:LEU:CD2	2.48	0.44
1:A:246:LEU:HA	1:A:249:VAL:HG13	2.00	0.44
1:A:118:TYR:HB3	1:A:119:PRO:HD3	2.00	0.43
1:B:292:VAL:HG12	1:B:293:LEU:HG	2.00	0.43
1:B:116:MET:O	1:B:119:PRO:HD2	2.19	0.43
1:B:38:GLN:HG2	1:B:293:LEU:HD22	1.99	0.43
1:A:235:GLN:O	1:A:235:GLN:HG3	2.18	0.43
1:B:220:GLU:OE1	1:B:220:GLU:HA	2.19	0.43
1:A:226:ARG:HB3	1:A:237:PRO:HG2	2.00	0.42
1:B:69:MET:CE	1:B:155:TYR:CE2	3.02	0.42
1:A:289:PHE:CE2	1:A:293:LEU:HD12	2.55	0.42
1:B:31:TRP:HH2	1:B:77:VAL:N	2.16	0.42
1:B:88:CYS:O	1:B:89:GLU:HB2	2.20	0.42
1:A:84:GLU:HG2	1:B:56:VAL:O	2.20	0.41
1:A:226:ARG:NH1	1:A:229:GLU:OE1	2.54	0.41
1:A:137:THR:HG22	1:A:261:LEU:HD11	2.02	0.41
1:A:103:ASP:OD1	1:A:103:ASP:C	2.59	0.41
1:A:155:TYR:CD1	1:A:155:TYR:C	2.95	0.41
1:A:213:GLU:HG3	1:A:266:ARG:HH12	1.87	0.40
1:A:47:LYS:NZ	1:A:98:GLU:OE1	2.42	0.40
1:B:27:ILE:O	1:B:80:LEU:HD12	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	243/280 (87%)	235 (97%)	8 (3%)	0	100 100
1	B	224/280 (80%)	215 (96%)	8 (4%)	1 (0%)	34 58
All	All	467/560 (83%)	450 (96%)	16 (3%)	1 (0%)	47 71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	225	ARG

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	233/259 (90%)	227 (97%)	6 (3%)	46 73
1	B	217/259 (84%)	213 (98%)	4 (2%)	59 81
All	All	450/518 (87%)	440 (98%)	10 (2%)	52 77

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	SER
1	A	76	GLU
1	A	77	VAL
1	A	137	THR
1	A	193	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	227	ASN
1	B	37	SER
1	B	200	ASP
1	B	210	ASP
1	B	226	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	236	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/280 (88%)	0.46	10 (4%) 38 36	59, 81, 165, 209	0
1	B	232/280 (82%)	0.92	41 (17%) 1 1	71, 123, 240, 335	0
All	All	481/560 (85%)	0.68	51 (10%) 6 4	59, 94, 221, 335	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	72	ASN	6.3
1	B	228	GLU	5.7
1	B	263	PHE	5.3
1	A	192	LEU	5.0
1	B	205	LEU	4.8
1	B	117	TYR	4.4
1	B	136	TYR	3.9
1	B	138	HIS	3.9
1	B	212	LEU	3.9
1	A	191	ARG	3.7
1	B	135	ALA	3.6
1	A	60	LEU	3.5
1	B	68	PHE	3.4
1	B	60	LEU	3.4
1	B	113	LYS	3.4
1	B	146	LEU	3.4
1	B	141	ILE	3.4
1	A	190	LYS	3.3
1	B	118	TYR	3.3
1	B	139	GLY	3.2
1	B	223	LEU	3.1
1	A	23	GLU	3.0
1	B	239	LEU	2.9
1	B	240	CYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	225	ARG	2.8
1	B	216	LEU	2.8
1	B	23	GLU	2.7
1	A	24	VAL	2.6
1	B	209	LEU	2.6
1	B	202	VAL	2.5
1	B	279	TYR	2.5
1	B	142	LEU	2.5
1	B	71	LEU	2.4
1	B	104	GLU	2.4
1	B	44	ILE	2.4
1	B	203	LYS	2.3
1	B	26	LEU	2.3
1	B	267	ASN	2.2
1	B	58	LEU	2.2
1	B	50	LYS	2.2
1	B	302	SER	2.2
1	B	265	ARG	2.2
1	A	189	GLN	2.1
1	B	266	ARG	2.1
1	B	49	LEU	2.1
1	A	69	MET	2.1
1	B	215	VAL	2.0
1	A	195	LYS	2.0
1	B	247	VAL	2.0
1	B	206	LYS	2.0
1	B	116	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.