



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2023 – 10:35 AM EDT

PDB ID : 2A5V  
Title : Crystal structure of *M. tuberculosis* beta carbonic anhydrase, Rv3588c, tetrameric form  
Authors : Covarrubias, A.S.; Bergfors, T.; Jones, T.A.; Hogbom, M.  
Deposited on : 2005-07-01  
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	213	88% 10% .
1	B	213	86% 11% ..
1	C	213	84% 9% . 6%
1	D	213	83% 15% .

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CARBONIC ANHYDRASE (CARBONATE DEHYDRATASE) (CARBONIC DEHYDRATASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	210	1562	972	297	289	4	0	0	0
1	B	210	1561	970	294	293	4	0	0	0
1	C	201	1484	925	276	279	4	0	0	0
1	D	209	1553	966	295	288	4	0	0	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	cloning artifact	UNP O53573
A	-4	HIS	-	cloning artifact	UNP O53573
A	-3	HIS	-	cloning artifact	UNP O53573
A	-2	HIS	-	cloning artifact	UNP O53573
A	-1	HIS	-	cloning artifact	UNP O53573
A	0	HIS	-	cloning artifact	UNP O53573
A	1	GLY	-	cloning artifact	UNP O53573
B	-5	HIS	-	cloning artifact	UNP O53573
B	-4	HIS	-	cloning artifact	UNP O53573
B	-3	HIS	-	cloning artifact	UNP O53573
B	-2	HIS	-	cloning artifact	UNP O53573
B	-1	HIS	-	cloning artifact	UNP O53573
B	0	HIS	-	cloning artifact	UNP O53573
B	1	GLY	-	cloning artifact	UNP O53573
C	-5	HIS	-	cloning artifact	UNP O53573
C	-4	HIS	-	cloning artifact	UNP O53573
C	-3	HIS	-	cloning artifact	UNP O53573
C	-2	HIS	-	cloning artifact	UNP O53573
C	-1	HIS	-	cloning artifact	UNP O53573
C	0	HIS	-	cloning artifact	UNP O53573

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	GLY	-	cloning artifact	UNP O53573
D	-5	HIS	-	cloning artifact	UNP O53573
D	-4	HIS	-	cloning artifact	UNP O53573
D	-3	HIS	-	cloning artifact	UNP O53573
D	-2	HIS	-	cloning artifact	UNP O53573
D	-1	HIS	-	cloning artifact	UNP O53573
D	0	HIS	-	cloning artifact	UNP O53573
D	1	GLY	-	cloning artifact	UNP O53573

- Molecule 2 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	N	S	0	0
			3	1	1	1		
2	B	1	Total	C	N	S	0	0
			3	1	1	1		
2	C	1	Total	C	N	S	0	0
			3	1	1	1		
2	C	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Zn	0	0
			4	4		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total 2	Zn 2	0	0
3	C	1	Total 1	Zn 1	0	0
3	D	4	Total 4	Zn 4	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	99	Total 99	O 99	0	0
4	B	110	Total 110	O 110	0	0
4	C	49	Total 49	O 49	0	0
4	D	53	Total 53	O 53	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CARBONIC ANHYDRASE (CARBONATE DEHYDRATASE) (CARBONIC DEHYDRATASE)

Chain A:  88% 10%





- Molecule 1: CARBONIC ANHYDRASE (CARBONATE DEHYDRATASE) (CARBONIC DEHYDRATASE)

Chain B:  86% 11%





- Molecule 1: CARBONIC ANHYDRASE (CARBONATE DEHYDRATASE) (CARBONIC DEHYDRATASE)

Chain C:  84% 9% 6%



- Molecule 1: CARBONIC ANHYDRASE (CARBONATE DEHYDRATASE) (CARBONIC DEHYDRATASE)

Chain D:  83% 15%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.88Å 70.30Å 84.36Å 90.00° 93.41° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.20 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-2.20) 99.5 (29.20-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.48 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.162 , 0.214 0.163 , 0.214	Depositor DCC
$R_{free}$ test set	2016 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.9	Xtrriage
Anisotropy	0.479	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6494	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SCN, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/1592	0.67	0/2164
1	B	0.67	0/1589	0.70	0/2160
1	C	0.57	0/1507	0.68	1/2048 (0.0%)
1	D	0.54	0/1582	0.65	1/2151 (0.0%)
All	All	0.60	0/6270	0.68	2/8523 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	167	ARG	CG-CD-NE	-5.88	99.45	111.80
1	C	80	ASP	CB-CG-OD1	5.62	123.36	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	79	ILE	Peptide



## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1562	0	1544	17	0
1	B	1561	0	1546	20	0
1	C	1484	0	1487	19	0
1	D	1553	0	1541	25	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
2	C	6	0	0	0	0
3	A	4	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
3	D	4	0	0	0	0
4	A	99	0	0	0	0
4	B	110	0	0	3	0
4	C	49	0	0	1	0
4	D	53	0	0	0	0
All	All	6494	0	6118	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:80:ASP:HB3	1:D:80:ASP:OD2	1.60	1.00
1:B:111:ASN:HB2	4:B:594:HOH:O	1.62	0.97
1:A:126:TYR:OH	1:C:126:TYR:OH	1.97	0.81
1:B:80:ASP:HB2	4:B:555:HOH:O	1.89	0.73
1:C:80:ASP:HB3	1:D:80:ASP:CG	2.11	0.71
1:C:4:THR:HG23	1:C:5:ASN:H	1.57	0.70
1:B:0:HIS:O	4:B:599:HOH:O	2.15	0.65
1:C:42:GLN:HE22	1:D:54:SER:H	1.46	0.62
1:A:42:GLN:HE22	1:B:54:SER:H	1.47	0.62
1:D:61:ILE:HD11	1:D:102:LEU:HD21	1.84	0.59
1:C:64:GLN:HG3	1:C:69:MET:HE3	1.83	0.59
1:D:183:VAL:HG23	1:D:196:LEU:CD1	2.34	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:ASP:CG	1:B:80:ASP:OD1	2.43	0.57
1:A:42:GLN:NE2	1:B:54:SER:H	2.04	0.55
1:B:4:THR:HG21	1:B:45:THR:HG22	1.89	0.55
1:B:2:PRO:O	1:B:4:THR:HG22	2.08	0.54
1:D:183:VAL:CG2	1:D:196:LEU:HD13	2.38	0.54
1:A:183:VAL:HG13	1:A:185:VAL:HG23	1.91	0.53
1:B:129:ASP:OD2	1:C:167:ARG:NH2	2.37	0.53
1:A:61:ILE:HD11	1:A:102:LEU:HD21	1.90	0.53
1:A:148:ARG:NH1	1:A:150:ASP:OD2	2.42	0.52
1:B:47:VAL:HG12	1:B:96:VAL:HG21	1.92	0.52
1:D:48:ILE:HG13	1:D:100:VAL:HB	1.91	0.51
1:C:54:SER:H	1:D:42:GLN:HE22	1.58	0.51
1:A:146:LEU:HD13	1:A:151:GLU:HB3	1.92	0.51
1:D:33:ASP:N	1:D:33:ASP:OD1	2.44	0.51
1:B:187:TYR:CE1	1:B:192:GLY:HA2	2.46	0.51
1:B:4:THR:HG21	1:B:45:THR:CG2	2.42	0.49
1:C:14:LYS:NZ	4:C:551:HOH:O	2.45	0.49
1:D:-3:HIS:O	1:D:40:ALA:HA	2.13	0.48
1:C:4:THR:HG23	1:C:5:ASN:N	2.28	0.48
1:D:28:PRO:O	1:D:65:GLY:HA3	2.14	0.48
1:D:185:VAL:HG11	1:D:194:ALA:HB1	1.96	0.47
1:A:45:THR:O	1:A:97:PRO:HD2	2.14	0.47
1:A:80:ASP:OD2	1:B:80:ASP:OD1	2.32	0.47
1:A:54:SER:H	1:B:42:GLN:HE22	1.62	0.47
1:B:70:PHE:CD2	1:B:90:ALA:HB2	2.50	0.47
1:C:42:GLN:NE2	1:D:54:SER:H	2.10	0.46
1:D:199:HIS:HD2	1:D:201:GLY:O	1.99	0.46
1:C:188:GLN:NE2	1:C:195:VAL:HG21	2.31	0.46
1:C:165:MET:HG3	1:C:200:ILE:CD1	2.46	0.46
1:D:183:VAL:CG2	1:D:196:LEU:CD1	2.95	0.45
1:A:122:LEU:HD12	1:D:140:LEU:HG	1.97	0.45
1:A:28:PRO:O	1:A:30:GLN:HG3	2.15	0.45
1:D:70:PHE:CD2	1:D:90:ALA:HB2	2.52	0.44
1:D:47:VAL:HG12	1:D:96:VAL:HG21	1.98	0.44
1:C:54:SER:H	1:D:42:GLN:NE2	2.16	0.44
1:D:183:VAL:HG23	1:D:196:LEU:HD13	1.98	0.44
1:D:100:VAL:HG22	1:D:183:VAL:CG1	2.47	0.44
1:D:113:ALA:O	1:D:117:ILE:HD12	2.19	0.43
1:C:64:GLN:HG3	1:C:69:MET:CE	2.48	0.43
1:A:73:ARG:CZ	1:B:71:VAL:HB	2.49	0.43
1:C:138:VAL:O	1:C:142:ARG:HB2	2.19	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:4:THR:HG23	1:D:97:PRO:HG3	2.01	0.42
1:D:43:LYS:HG2	1:D:43:LYS:O	2.19	0.42
1:A:78:VAL:HG23	1:B:82:ALA:HB1	2.02	0.42
1:C:165:MET:HG3	1:C:200:ILE:HD13	2.01	0.41
1:C:70:PHE:CD2	1:C:90:ALA:HB2	2.56	0.41
1:B:61:ILE:HD11	1:B:102:LEU:HD21	2.03	0.41
1:B:148:ARG:HB2	1:B:151:GLU:HG3	2.03	0.41
1:A:140:LEU:HG	1:D:122:LEU:HD12	2.03	0.40
1:B:4:THR:CG2	1:B:45:THR:HG22	2.50	0.40
1:C:148:ARG:HB3	1:C:150:ASP:OD1	2.20	0.40
1:A:188:GLN:NE2	1:A:195:VAL:HG21	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/213 (98%)	204 (98%)	4 (2%)	0	100	100
1	B	208/213 (98%)	202 (97%)	6 (3%)	0	100	100
1	C	199/213 (93%)	196 (98%)	3 (2%)	0	100	100
1	D	207/213 (97%)	199 (96%)	8 (4%)	0	100	100
All	All	822/852 (96%)	801 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/164 (98%)	158 (99%)	2 (1%)	69	81
1	B	161/164 (98%)	154 (96%)	7 (4%)	29	36
1	C	153/164 (93%)	150 (98%)	3 (2%)	55	69
1	D	160/164 (98%)	156 (98%)	4 (2%)	47	60
All	All	634/656 (97%)	618 (98%)	16 (2%)	47	60

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	129	ASP
1	A	183	VAL
1	B	3	ASN
1	B	4	THR
1	B	47	VAL
1	B	48	ILE
1	B	61	ILE
1	B	142	ARG
1	B	167	ARG
1	C	69	MET
1	C	84	LEU
1	C	129	ASP
1	D	24	ARG
1	D	33	ASP
1	D	84	LEU
1	D	129	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	188	GLN
1	B	-3	HIS
1	B	42	GLN
1	B	95	ASN
1	B	188	GLN
1	C	42	GLN
1	C	154	GLN
1	C	188	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	-2	HIS
1	D	42	GLN
1	D	199	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 11 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SCN	C	503	3	1,2,2	1.21	0	0,1,1	-	-
2	SCN	C	504	-	1,2,2	0.82	0	0,1,1	-	-
2	SCN	A	501	3	1,2,2	1.48	0	0,1,1	-	-
2	SCN	B	502	3	1,2,2	1.45	0	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.