



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 02:24 PM EDT

PDB ID : 1A7T
Title : METALLO-BETA-LACTAMASE WITH MES
Authors : Fitzgerald, P.M.D.; Wu, J.K.; Toney, J.H.
Deposited on : 1998-03-17
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	 83% 13% ..
1	B	232	 81% 16% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METALLO-BETA-LACTAMASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	227	1744	1105	289	341	9	0	0	0
1	B	227	1744	1105	289	341	9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	171	THR	ALA	engineered mutation	UNP P25910
A	208	ASN	ASP	engineered mutation	UNP P25910
B	171	THR	ALA	engineered mutation	UNP P25910
B	208	ASN	ASP	engineered mutation	UNP P25910

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Zn 2	0	0
2	B	2	Total 2	Zn 2	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Na 1	0	0
3	B	1	Total 1	Na 1	0	0

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	12	6	1	4	1	0	0
4	B	1	12	6	1	4	1	0	0


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	167	167	167	0	0
5	B	170	170	170	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: METALLO-BETA-LACTAMASE

Chain A:  83% 13% ..



- Molecule 1: METALLO-BETA-LACTAMASE

Chain B:  81% 16% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.27Å 66.96Å 69.48Å 90.00° 103.77° 90.00°	Depositor
Resolution (Å)	10.00 – 1.85 31.87 – 1.80	Depositor EDS
% Data completeness (in resolution range)	91.3 (10.00-1.85) 85.9 (31.87-1.80)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 1.79Å)	Xtrriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.151 , (Not available) 0.146 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	12.6	Xtrriage
Anisotropy	0.124	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 85.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3855	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MES, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	0/1784	1.54	10/2433 (0.4%)
1	B	0.91	0/1784	1.49	13/2433 (0.5%)
All	All	0.89	0/3568	1.51	23/4866 (0.5%)

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	179	ASP	CB-CG-OD2	14.98	131.78	118.30
1	A	179	ASP	CB-CG-OD1	-10.84	108.54	118.30
1	B	200	ARG	NE-CZ-NH2	-9.39	115.61	120.30
1	B	9	ASP	CB-CG-OD1	9.30	126.67	118.30
1	B	57	ASP	CB-CG-OD1	7.47	125.02	118.30
1	B	18	ASP	CB-CG-OD1	7.40	124.96	118.30
1	A	109	ASP	CB-CG-OD1	6.80	124.42	118.30
1	A	148	ASP	CB-CG-OD1	6.79	124.41	118.30
1	B	148	ASP	CB-CG-OD1	6.77	124.40	118.30
1	A	120	GLU	CA-CB-CG	6.75	128.25	113.40
1	B	12	SER	N-CA-CB	6.09	119.64	110.50
1	B	120	GLU	CB-CA-C	-6.05	98.31	110.40
1	B	46	HIS	CA-CB-CG	-5.79	103.75	113.60
1	B	52	ASP	N-CA-CB	5.68	120.82	110.60
1	A	173	SER	N-CA-CB	-5.67	102.00	110.50
1	A	201	TYR	CB-CG-CD1	-5.60	117.64	121.00
1	B	125	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	113	GLU	OE1-CD-OE2	-5.44	116.77	123.30
1	A	10	ASP	CA-CB-CG	5.26	124.96	113.40
1	A	10	ASP	CB-CA-C	-5.24	99.92	110.40
1	B	120	GLU	CA-CB-CG	5.20	124.84	113.40
1	B	30	GLU	OE1-CD-OE2	-5.02	117.27	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	58	ALA	N-CA-CB	-5.00	103.10	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1744	0	1703	17	0
1	B	1744	0	1703	27	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	12	0	12	0	0
4	B	12	0	13	0	0
5	A	167	0	0	7	0
5	B	170	0	0	6	0
All	All	3855	0	3431	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ILE:CD1	1:B:71:LEU:HD13	1.78	1.13
1:A:106:MET:HA	5:A:394:HOH:O	1.69	0.93
1:B:43:ILE:CD1	1:B:71:LEU:CD1	2.57	0.80
1:B:43:ILE:HD11	1:B:71:LEU:CD1	2.13	0.79
1:A:183:THR:HG21	5:A:406:HOH:O	1.90	0.71
1:B:124:THR:O	5:B:321:HOH:O	2.09	0.69
1:A:223:ASN:O	1:A:227:GLU:HG3	1.94	0.68
1:B:43:ILE:HD12	1:B:71:LEU:HD13	1.72	0.66
1:B:193:LYS:HE2	1:B:216:GLU:OE2	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ILE:HD13	1:B:71:LEU:HD13	1.75	0.60
1:A:182:VAL:HG23	5:A:379:HOH:O	2.02	0.59
1:B:43:ILE:HD11	1:B:71:LEU:HD12	1.83	0.59
1:A:43:ILE:CD1	1:A:71:LEU:HD13	2.33	0.59
1:A:83:TRP:CH2	1:A:114:LYS:HE3	2.39	0.58
1:A:74:LYS:HE3	5:A:350:HOH:O	2.05	0.56
1:A:39:GLY:C	1:A:40:MET:HG2	2.26	0.55
1:A:169:ASN:ND2	5:A:396:HOH:O	2.38	0.55
1:A:170:GLN:HG3	5:A:413:HOH:O	2.09	0.52
1:B:23:TYR:CE2	1:B:39:GLY:HA3	2.47	0.49
1:B:183:THR:CG2	5:B:377:HOH:O	2.59	0.49
1:A:83:TRP:CZ2	1:A:114:LYS:HE3	2.48	0.49
1:B:200:ARG:HG3	5:B:331:HOH:O	2.12	0.49
1:B:45:ASN:ND2	1:B:132:ASP:OD2	2.34	0.47
1:A:41:ILE:HD11	1:A:63:LEU:HD11	1.97	0.47
1:B:7:ILE:HD13	1:B:13:ILE:CG1	2.45	0.47
1:B:41:ILE:HD11	1:B:63:LEU:HD11	1.96	0.46
1:A:43:ILE:HD11	1:A:71:LEU:HD13	1.98	0.45
1:B:139:TYR:HE1	5:B:424:HOH:O	2.00	0.44
1:B:183:THR:HG21	5:B:377:HOH:O	2.16	0.44
1:B:29:ILE:HD12	1:B:35:VAL:HG21	1.99	0.44
1:B:7:ILE:HD13	1:B:13:ILE:HG13	1.99	0.44
1:A:135:PRO:HB3	5:A:292:HOH:O	2.18	0.44
1:B:23:TYR:CZ	1:B:39:GLY:HA3	2.53	0.44
1:A:43:ILE:HD11	1:A:71:LEU:CD1	2.47	0.43
1:A:55:ILE:HD12	1:A:55:ILE:HG21	1.68	0.43
1:B:195:LYS:HE2	5:B:397:HOH:O	2.18	0.43
1:A:187:LYS:HD3	1:A:187:LYS:HA	1.40	0.43
1:B:88:ILE:HD13	1:B:88:ILE:HG21	1.69	0.43
1:B:29:ILE:HD12	1:B:35:VAL:CG2	2.49	0.42
1:B:55:ILE:HG23	1:B:55:ILE:HD13	1.84	0.42
1:B:29:ILE:HD12	1:B:35:VAL:HB	2.03	0.41
1:B:226:ILE:O	1:B:230:SER:OG	2.39	0.41
1:B:103:ALA:O	1:B:122:GLY:HA2	2.21	0.41
1:B:92:GLY:O	1:B:96:ARG:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/232 (97%)	218 (97%)	7 (3%)	0	100	100
1	B	225/232 (97%)	219 (97%)	6 (3%)	0	100	100
All	All	450/464 (97%)	437 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/198 (98%)	184 (95%)	10 (5%)	23	8
1	B	194/198 (98%)	190 (98%)	4 (2%)	53	38
All	All	388/396 (98%)	374 (96%)	14 (4%)	35	18

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LYS
1	A	9	ASP
1	A	10	ASP
1	A	17	SER
1	A	55	ILE
1	A	62	MET
1	A	114	LYS
1	A	125	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	149	ASN
1	A	195	LYS
1	B	9	ASP
1	B	55	ILE
1	B	149	ASN
1	B	173	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	220	GLN
1	B	158	ASN
1	B	220	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MES	A	250	-	12,12,12	0.93	0	14,16,16	2.39	3 (21%)
4	MES	B	250	-	12,12,12	0.90	0	14,16,16	2.19	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	A	250	-	-	1/6/14/14	0/1/1/1
4	MES	B	250	-	-	1/6/14/14	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	250	MES	O2S-S-C8	6.99	115.33	106.92
4	B	250	MES	O2S-S-C8	4.63	112.50	106.92
4	B	250	MES	C2-C3-N4	-3.80	104.35	110.10
4	A	250	MES	C5-N4-C3	3.33	116.33	108.83
4	B	250	MES	O1S-S-C8	2.58	110.02	106.92
4	A	250	MES	C7-N4-C3	2.45	117.51	111.23
4	B	250	MES	O1-C2-C3	-2.43	106.45	111.80
4	B	250	MES	C5-N4-C3	2.12	113.60	108.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	250	MES	N4-C7-C8-S
4	B	250	MES	N4-C7-C8-S

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/232 (97%)	-0.64	1 (0%) 92 92	5, 13, 25, 41	0
1	B	227/232 (97%)	-0.69	0 100 100	5, 12, 25, 33	0
All	All	454/464 (97%)	-0.66	1 (0%) 95 94	5, 12, 25, 41	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	230	SER	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MES	A	250	12/12	0.97	0.09	27,28,29,29	0
4	MES	B	250	12/12	0.97	0.11	24,25,29,29	0
3	NA	A	255	1/1	0.99	0.49	37,37,37,37	0
3	NA	B	255	1/1	0.99	0.49	35,35,35,35	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	A	251	1/1	1.00	0.02	10,10,10,10	0
2	ZN	A	252	1/1	1.00	0.02	13,13,13,13	0
2	ZN	B	251	1/1	1.00	0.01	9,9,9,9	0
2	ZN	B	252	1/1	1.00	0.02	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.