



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 07:41 PM EDT

PDB ID : 7A76
Title : Bacillithiol Disulfide Reductase Bdr (YpdA) from Bacillus cereus
Authors : Hammerstad, M.; Gudim, I.; Hersleth, H.-P.
Deposited on : 2020-08-27
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	
1	B	326	
1	C	326	
1	D	326	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 11737 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THIOREDOXIN REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	326	Total 2658	C 1687	N 460	O 503	S 8	0	9	0
1	B	326	Total 2695	C 1717	N 465	O 506	S 7	0	13	0
1	C	325	Total 2657	C 1688	N 456	O 505	S 8	0	11	0
1	D	324	Total 2654	C 1688	N 455	O 504	S 7	0	11	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	B	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	D	1	Total	Na	0	0
			1	1		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	283	Total	O	0	7
			283	283		
4	B	251	Total	O	0	7
			251	251		
4	C	182	Total	O	0	4
			182	182		
4	D	141	Total	O	0	2
			141	141		



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: THIOREDOXIN REDUCTASE





- Molecule 1: THIOREDOXIN REDUCTASE



- Molecule 1: THIOREDOXIN REDUCTASE



- Molecule 1: THIOREDOXIN REDUCTASE



LYS

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.73Å 114.90Å 146.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.23 – 1.65 73.37 – 1.65	Depositor EDS
% Data completeness (in resolution range)	97.5 (45.23-1.65) 97.5 (73.37-1.65)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.65Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660, REFMAC 5.8.0253	Depositor
R, R_{free}	0.190 , 0.218 0.192 , 0.220	Depositor DCC
R_{free} test set	8788 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtrriage
Anisotropy	0.401	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11737	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/2713	0.57	0/3666
1	B	0.38	0/2753	0.57	0/3725
1	C	0.35	0/2713	0.53	0/3671
1	D	0.33	0/2709	0.53	0/3667
All	All	0.36	0/10888	0.55	0/14729

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2658	0	2618	15	0
1	B	2695	0	2652	29	0
1	C	2657	0	2607	25	0
1	D	2654	0	2604	37	0
2	A	53	0	31	0	0
2	B	53	0	31	4	0
2	C	53	0	31	2	0
2	D	53	0	31	4	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	283	0	0	1	1
4	B	251	0	0	6	1
4	C	182	0	0	6	1
4	D	141	0	0	2	1
All	All	11737	0	10605	101	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:285:THR:HG23	1:D:287:ALA:H	1.53	0.74
1:A:273:THR:HG23	1:A:275:ARG:H	1.54	0.73
1:A:47:HIS:CE1	1:B:74[B]:ASN:HD21	2.08	0.72
1:A:99:LYS:NZ	1:A:101:GLN:OE1	2.25	0.69
1:D:48[A]:GLN:HG2	2:D:401:FAD:HM72	1.76	0.67
1:A:124:LYS:HE3	1:A:326:LYS:HG2	1.78	0.66
1:D:282:ARG:NH1	1:D:284:GLU:OE1	2.30	0.64
1:D:51:PHE:N	2:D:401:FAD:O4	2.31	0.63
1:B:48[A]:GLN:HG2	2:B:401:FAD:HM72	1.81	0.63
1:D:47:HIS:HD2	4:D:539:HOH:O	1.83	0.62
1:B:47:HIS:HD2	4:B:547:HOH:O	1.83	0.60
1:D:100:VAL:HG22	1:D:109:VAL:HG12	1.84	0.59
1:B:133[A]:TYR:OH	2:B:401:FAD:H9	2.03	0.59
1:B:273:THR:O	1:B:275:ARG:N	2.33	0.59
1:B:112:THR:OG1	1:B:118:LYS:NZ	2.30	0.58
1:D:313:HIS:O	1:D:317:ILE:HG12	2.04	0.58
1:B:24:LYS:NZ	4:B:502:HOH:O	2.33	0.58
1:B:226:LYS:NZ	4:B:504:HOH:O	2.36	0.58
1:D:283:MET:HB2	1:D:317:ILE:HD11	1.85	0.58
1:B:47:HIS:HE1	1:B:175:ASP:OD1	1.87	0.57
1:B:51:PHE:N	2:B:401:FAD:O4	2.35	0.56
1:C:133[A]:TYR:OH	2:C:401:FAD:H9	2.07	0.55
1:D:115[A]:ASP:OD1	1:D:115[A]:ASP:N	2.31	0.55
1:D:285:THR:HG22	1:D:290:ILE:O	2.07	0.54
1:B:68:ASN:HB2	4:B:597:HOH:O	2.08	0.54
1:C:274:GLY:HA3	1:C:300:ASN:ND2	2.23	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:51:PHE:N	2:C:401:FAD:O4	2.40	0.53
1:D:47:HIS:HE1	1:D:175[B]:ASP:OD1	1.92	0.53
1:C:115:ASP:OD2	1:D:231:HIS:HE1	1.90	0.53
1:D:234:THR:HG23	4:D:540:HOH:O	2.07	0.53
1:D:267:VAL:HG22	1:D:285:THR:HG21	1.92	0.51
1:D:283:MET:HB2	1:D:317:ILE:CD1	2.40	0.51
1:C:47:HIS:ND1	4:C:505:HOH:O	2.34	0.51
1:D:234:THR:HG22	1:D:243:THR:OG1	2.10	0.51
1:D:259:HIS:HB3	1:D:269:ILE:HD13	1.91	0.51
1:C:304:GLU:HG3	1:C:305:ILE:HG13	1.93	0.50
1:B:100:VAL:HB	1:B:265:MET:HG3	1.93	0.50
1:C:298:ALA:HB1	1:C:302:ALA:HA	1.92	0.50
1:D:4:GLU:HG3	1:D:7[B]:ILE:HD11	1.92	0.50
1:A:47:HIS:CE1	1:B:74[B]:ASN:ND2	2.78	0.50
1:C:137:ASN:OD1	1:C:256:HIS:HD2	1.95	0.50
1:C:259:HIS:CE1	1:C:300:ASN:HD21	2.30	0.50
1:D:4:GLU:O	1:D:5:THR:OG1	2.20	0.49
1:D:285:THR:HG23	1:D:287:ALA:N	2.26	0.49
1:A:231:HIS:HE1	1:B:115:ASP:OD2	1.96	0.49
1:C:5:THR:HG22	1:C:6:VAL:HG23	1.95	0.49
1:A:69[B]:ARG:NE	1:C:207:GLU:OE2	2.46	0.49
1:D:47:HIS:HE1	1:D:175[A]:ASP:OD2	1.95	0.49
1:D:194:ILE:HD13	1:D:194:ILE:H	1.77	0.48
1:D:27:ILE:O	1:D:29:PRO:HD3	2.14	0.48
1:D:48[B]:GLN:HG3	2:D:401:FAD:HM72	1.96	0.48
1:D:139[B]:MET:HE2	1:D:251:ALA:HB1	1.96	0.48
1:C:114:ARG:NH2	4:C:504:HOH:O	2.26	0.48
1:D:278:TYR:CD1	1:D:292:ILE:HD11	2.48	0.47
1:C:278:TYR:CD2	1:C:292:ILE:HD11	2.49	0.47
1:D:262:LEU:O	1:D:265:MET:HG2	2.14	0.47
1:C:68:ASN:HB2	4:C:554:HOH:O	2.14	0.47
1:A:256:HIS:HE1	4:A:742:HOH:O	1.99	0.46
1:B:132:TYR:HE1	1:B:133[A]:TYR:CE1	2.33	0.46
1:B:56:LYS:HB3	1:B:308:GLU:HG2	1.98	0.45
1:A:1:MET:HG3	1:A:122:ILE:HG12	1.99	0.45
1:B:268:GLN:HG3	1:B:286:ASN:ND2	2.32	0.45
1:A:219[B]:MET:HG2	1:A:221:PHE:CE2	2.52	0.45
1:A:271:GLU:O	1:A:272:GLU:HB3	2.16	0.45
1:C:69:ARG:O	4:C:502:HOH:O	2.21	0.45
1:A:270:ASP:HB3	1:A:273:THR:HG22	1.99	0.45
1:B:226:LYS:NZ	4:B:507:HOH:O	2.45	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:291:PHE:HB3	1:D:317:ILE:HD12	1.98	0.44
1:C:51:PHE:HB3	1:C:201:LYS:HE2	2.00	0.44
1:C:282:ARG:NH1	1:C:284:GLU:OE1	2.50	0.44
1:B:4:GLU:O	1:B:123:ALA:HA	2.18	0.43
1:D:63:ALA:HB1	1:D:65:ILE:HD13	2.00	0.43
1:D:275:ARG:O	1:D:275:ARG:HG2	2.18	0.43
1:C:300:ASN:N	1:C:300:ASN:HD22	2.16	0.43
1:C:262:LEU:HD11	1:C:296:ILE:HD11	2.01	0.43
1:C:135:ASN:HB2	1:C:256:HIS:O	2.19	0.43
1:D:139[B]:MET:HE1	1:D:167:ILE:HG21	2.01	0.43
1:B:213:ARG:HE	1:B:213:ARG:HB3	1.59	0.43
1:D:291:PHE:CE2	1:D:321:ILE:HD11	2.54	0.43
1:C:46:THR:O	4:C:503:HOH:O	2.22	0.42
1:B:65[B]:ILE:HG23	4:C:570:HOH:O	2.17	0.42
1:D:273:THR:HG23	1:D:275:ARG:H	1.83	0.42
1:B:133[A]:TYR:OH	2:B:401:FAD:H1'1	2.20	0.42
1:B:278:TYR:CD1	1:B:292:ILE:HD11	2.53	0.42
1:A:108:GLN:HG2	1:A:122:ILE:HD13	2.02	0.42
1:B:162[A]:ARG:NH2	4:B:511:HOH:O	2.51	0.41
1:B:265:MET:HG2	1:B:290:ILE:HD13	2.02	0.41
1:C:102:LYS:HD2	1:C:104:GLY:O	2.21	0.41
1:C:262:LEU:O	1:C:265[B]:MET:HG2	2.20	0.41
1:B:63:ALA:HB1	1:B:65[A]:ILE:HD13	2.03	0.41
1:A:137:ASN:OD1	1:A:256:HIS:HD2	2.04	0.41
1:B:205:LEU:HG	1:B:209:GLU:HG3	2.03	0.41
1:C:4:GLU:O	1:C:123:ALA:HA	2.20	0.41
1:D:270:ASP:OD2	1:D:275:ARG:NH2	2.54	0.41
1:C:298:ALA:CB	1:C:302:ALA:HA	2.49	0.41
1:D:133[A]:TYR:OH	2:D:401:FAD:H9	2.20	0.41
1:B:65[B]:ILE:O	1:B:65[B]:ILE:HG13	2.22	0.40
1:A:194:ILE:H	1:A:194:ILE:HG13	1.77	0.40
1:B:110:GLU:HG2	1:B:120:ILE:HG12	2.03	0.40
1:D:67[A]:GLU:HG3	1:D:68:ASN:N	2.36	0.40
1:D:273:THR:HG21	1:D:275:ARG:CZ	2.51	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:654:HOH:O	4:D:629:HOH:O[4_754]	2.15	0.05
4:A:737:HOH:O	4:B:524:HOH:O[2_555]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/326 (102%)	324 (97%)	8 (2%)	1 (0%)	41	22
1	B	337/326 (103%)	325 (96%)	10 (3%)	2 (1%)	25	8
1	C	334/326 (102%)	327 (98%)	7 (2%)	0	100	100
1	D	333/326 (102%)	320 (96%)	11 (3%)	2 (1%)	25	8
All	All	1337/1304 (102%)	1296 (97%)	36 (3%)	5 (0%)	34	16

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	300	ASN
1	D	5	THR
1	B	274	GLY
1	D	271	GLU
1	A	325	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/273 (103%)	275 (98%)	7 (2%)	47	22
1	B	286/273 (105%)	278 (97%)	8 (3%)	43	18
1	C	283/273 (104%)	278 (98%)	5 (2%)	59	36
1	D	282/273 (103%)	266 (94%)	16 (6%)	20	4
All	All	1133/1092 (104%)	1097 (97%)	36 (3%)	43	13

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	PHE
1	A	132	TYR
1	A	219[A]	MET
1	A	219[B]	MET
1	A	260	SER
1	A	281	ASP
1	A	325	GLU
1	B	64	PHE
1	B	65[A]	ILE
1	B	65[B]	ILE
1	B	102	LYS
1	B	103	ASP
1	B	132	TYR
1	B	213	ARG
1	B	306	PHE
1	C	64	PHE
1	C	268	GLN
1	C	271	GLU
1	C	303	ASN
1	C	306	PHE
1	D	64	PHE
1	D	67[A]	GLU
1	D	67[B]	GLU
1	D	68	ASN
1	D	108	GLN
1	D	115[A]	ASP
1	D	115[B]	ASP
1	D	132	TYR
1	D	173	SER
1	D	175[A]	ASP
1	D	175[B]	ASP
1	D	194	ILE
1	D	260	SER
1	D	275	ARG
1	D	279	THR
1	D	306	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	231	HIS
1	A	256	HIS
1	B	47	HIS
1	C	39	ASN
1	C	47	HIS
1	C	101	GLN
1	C	256	HIS
1	C	300	ASN
1	C	319	GLN
1	D	2	GLN
1	D	39	ASN
1	D	47	HIS
1	D	231	HIS
1	D	256	HIS
1	D	300	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	C	401	-	54,58,58	0.74	1 (1%)	71,89,89	0.57	1 (1%)
2	FAD	A	401	-	54,58,58	0.55	0	71,89,89	0.67	2 (2%)
2	FAD	D	401	-	54,58,58	0.60	0	71,89,89	0.62	1 (1%)
2	FAD	B	401	-	54,58,58	0.74	1 (1%)	71,89,89	0.64	2 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	C	401	-	-	0/30/50/50	0/6/6/6
2	FAD	A	401	-	-	1/30/50/50	0/6/6/6
2	FAD	D	401	-	-	1/30/50/50	0/6/6/6
2	FAD	B	401	-	-	0/30/50/50	0/6/6/6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	FAD	P-O3P	-4.00	1.55	1.59
2	B	401	FAD	P-O3P	3.27	1.63	1.59

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	FAD	C4B-O4B-C1B	-2.41	107.72	109.92
2	C	401	FAD	C5A-C6A-N6A	2.35	123.88	120.31
2	D	401	FAD	C5A-C6A-N6A	2.28	123.79	120.31
2	A	401	FAD	C5A-C6A-N6A	2.24	123.72	120.31
2	A	401	FAD	O2A-PA-O1A	2.18	122.59	112.44
2	B	401	FAD	C5A-C6A-N6A	2.10	123.51	120.31

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	FAD	PA-O3P-P-O5'
2	D	401	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.


3 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	401	FAD	2	0
2	D	401	FAD	4	0
2	B	401	FAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/326 (100%)	0.06	8 (2%) 57 58	17, 33, 56, 82	0
1	B	326/326 (100%)	0.15	13 (3%) 38 38	17, 32, 62, 92	0
1	C	325/326 (99%)	0.41	23 (7%) 16 15	18, 41, 77, 94	0
1	D	324/326 (99%)	0.68	42 (12%) 3 2	18, 49, 92, 132	0
All	All	1301/1304 (99%)	0.32	86 (6%) 18 17	17, 38, 78, 132	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	271	GLU	9.9
1	B	301	ASN	9.4
1	D	300	ASN	9.3
1	C	302	ALA	8.9
1	C	277	PHE	8.8
1	B	273	THR	8.6
1	D	273	THR	8.5
1	D	299	GLY	8.3
1	C	269	ILE	7.9
1	D	304	GLU	7.8
1	D	274	GLY	7.7
1	D	269	ILE	7.5
1	C	301	ASN	7.4
1	C	303	ASN	6.2
1	C	273	THR	5.9
1	D	301	ASN	5.8
1	D	277	PHE	5.7
1	D	305	ILE	5.7
1	C	275	ARG	5.3
1	A	326	LYS	5.2
1	D	303	ASN	5.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	270	ASP	5.1
1	D	298	ALA	5.1
1	C	272	GLU	5.0
1	C	270	ASP	5.0
1	C	268	GLN	4.9
1	B	300	ASN	4.9
1	B	303	ASN	4.8
1	A	272	GLU	4.7
1	D	268	GLN	4.2
1	D	272	GLU	4.1
1	C	263	THR	4.0
1	A	271	GLU	3.8
1	D	239	GLY	3.7
1	D	275	ARG	3.6
1	B	277[A]	PHE	3.6
1	D	306	PHE	3.6
1	C	299	GLY	3.5
1	B	326	LYS	3.5
1	B	271	GLU	3.5
1	C	271	GLU	3.4
1	D	259	HIS	3.4
1	D	120	ILE	3.4
1	D	302	ALA	3.3
1	B	302	ALA	3.2
1	D	276	PRO	3.1
1	D	295	VAL	3.0
1	D	238	ASP	2.9
1	C	283	MET	2.9
1	A	300	ASN	2.8
1	D	267	VAL	2.7
1	D	242	LEU	2.7
1	B	299	GLY	2.6
1	D	237	VAL	2.6
1	D	263	THR	2.6
1	D	285	THR	2.6
1	C	276	PRO	2.6
1	D	194	ILE	2.6
1	C	261	PHE	2.6
1	D	283	MET	2.5
1	B	306	PHE	2.5
1	D	260	SER	2.4
1	C	51	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	298	ALA	2.4
1	B	274	GLY	2.4
1	C	288	GLU	2.4
1	D	125	TYR	2.3
1	D	123	ALA	2.3
1	A	301	ASN	2.3
1	C	323	SER	2.3
1	C	306	PHE	2.3
1	D	262	LEU	2.2
1	B	304	GLU	2.2
1	B	305	ILE	2.2
1	D	323	SER	2.2
1	D	257	PRO	2.2
1	A	69[A]	ARG	2.2
1	D	51	PHE	2.1
1	D	291	PHE	2.1
1	A	273	THR	2.1
1	D	133[A]	TYR	2.1
1	C	242[A]	LEU	2.1
1	D	290	ILE	2.1
1	C	274	GLY	2.0
1	A	299	GLY	2.0
1	D	288	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FAD	D	401	53/53	0.93	0.14	26,44,74,79	0
2	FAD	C	401	53/53	0.95	0.12	22,30,39,44	0
2	FAD	A	401	53/53	0.96	0.10	21,27,33,39	0
3	NA	D	402	1/1	0.96	0.17	29,29,29,29	0
2	FAD	B	401	53/53	0.97	0.10	17,26,48,50	0
3	NA	C	402	1/1	0.98	0.14	26,26,26,26	0
3	NA	A	402	1/1	0.98	0.23	25,25,25,25	0
3	NA	B	402	1/1	0.99	0.19	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around FAD C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.