



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2023 – 01:32 pm GMT

PDB ID : 8A8Z
Title : Crystal structure of Danio rerio HDAC6 CD2 in complex with in situ enzymatically hydrolyzed DFMO-based ITF5924
Authors : Zrubek, K.; Sandrone, G.; Cukier, C.D.; Stevenazzi, A.
Deposited on : 2022-06-27
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.3
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	
1	B	365	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6377 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histone deacetylase 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2824	1775	504	527	18	0	8	0
1	B	357	2805	1763	502	522	18	0	4	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	434	GLY	-	cloning artifact	UNP F8W4B7
A	435	SER	-	cloning artifact	UNP F8W4B7
A	436	ASN	-	cloning artifact	UNP F8W4B7
A	437	ALA	-	cloning artifact	UNP F8W4B7
A	438	GLY	-	cloning artifact	UNP F8W4B7
A	439	GLY	-	cloning artifact	UNP F8W4B7
B	434	GLY	-	cloning artifact	UNP F8W4B7
B	435	SER	-	cloning artifact	UNP F8W4B7
B	436	ASN	-	cloning artifact	UNP F8W4B7
B	437	ALA	-	cloning artifact	UNP F8W4B7
B	438	GLY	-	cloning artifact	UNP F8W4B7
B	439	GLY	-	cloning artifact	UNP F8W4B7

- Molecule 2 is 4-[[4-[4-(imidazolidin-2-ylideneamino)phenyl]-1,2,3-triazol-1-yl]methyl]benzo hydrazide (three-letter code: LDL) (formula: C₁₉H₂₀N₈O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	19	8	1		
2	B	1	Total	C	N	O	0	0
			28	19	8	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	K	0	0
			2	2		
4	B	2	Total	K	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	362	Total	O	0	0
			362	362		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	324	Total 324	O 324	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.37Å 118.92Å 60.26Å 90.00° 93.87° 90.00°	Depositor
Resolution (Å)	47.88 – 1.60 47.84 – 1.60	Depositor EDS
% Data completeness (in resolution range)	95.4 (47.88-1.60) 95.4 (47.84-1.60)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.155 , 0.190 0.170 , 0.199	Depositor DCC
R_{free} test set	4695 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtrriage
Anisotropy	0.290	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6377	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, K, LDL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/2910	0.81	0/3946
1	B	0.72	0/2885	0.83	1/3913 (0.0%)
All	All	0.73	0/5795	0.82	1/7859 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	788	ARG	CG-CD-NE	-6.34	98.48	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2824	0	2749	10	0
1	B	2805	0	2727	6	0
2	A	28	0	0	0	0
2	B	28	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	362	0	0	5	2
5	B	324	0	0	3	2
All	All	6377	0	5476	16	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:MET:HE3	5:B:912:HOH:O	1.59	1.00
1:A:442:PRO:N	5:A:901:HOH:O	2.05	0.87
1:B:727:HIS:HD2	5:B:1166:HOH:O	1.75	0.68
1:A:565[B]:VAL:HG23	1:A:761:LEU:HD12	1.81	0.63
1:B:452:MET:CE	5:B:912:HOH:O	2.32	0.56
1:A:530:ASN:HB2	5:A:1176:HOH:O	2.06	0.55
1:B:720:GLU:HA	1:B:756:MET:HE3	1.95	0.48
1:A:641:ALA:O	1:A:645:ASN:ND2	2.47	0.48
1:A:510:ILE:HG13	5:A:902:HOH:O	2.15	0.47
1:B:704:PHE:HB2	1:B:750:ILE:HG22	1.97	0.46
1:A:772:LEU:HB2	5:A:942:HOH:O	2.15	0.46
1:A:782:SER:O	1:A:786:VAL:HG23	2.16	0.45
1:A:459:TRP:HZ2	5:A:1071:HOH:O	1.99	0.45
1:A:791:ALA:N	1:A:792:PRO:CD	2.81	0.44
1:A:454:LEU:HB3	1:A:537:GLU:HG2	1.99	0.44
1:B:459:TRP:CZ2	1:B:523:ASN:HB2	2.54	0.42

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1228:HOH:O	5:B:1130:HOH:O[2_554]	2.12	0.08
5:A:934:HOH:O	5:B:1152:HOH:O[1_656]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/365 (100%)	352 (97%)	11 (3%)	0	100	100
1	B	359/365 (98%)	346 (96%)	12 (3%)	1 (0%)	41	21
All	All	722/730 (99%)	698 (97%)	23 (3%)	1 (0%)	51	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	443	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/305 (101%)	304 (98%)	5 (2%)	62	41
1	B	305/305 (100%)	300 (98%)	5 (2%)	62	41
All	All	614/610 (101%)	604 (98%)	10 (2%)	60	41

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	465	GLU
1	A	573	HIS
1	A	611	TRP
1	A	658	LYS
1	A	765	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	465	GLU
1	B	573	HIS
1	B	611	TRP
1	B	771	HIS
1	B	777	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	536	ASN
1	A	645	ASN
1	A	784	ASN
1	B	536	ASN
1	B	716	GLN
1	B	727	HIS
1	B	771	HIS
1	B	784	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LDL	B	801	3	30,31,31	0.85	2 (6%)	38,42,42	1.26	5 (13%)
2	LDL	A	801	3	30,31,31	0.77	2 (6%)	38,42,42	1.17	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LDL	B	801	3	-	1/18/25/25	0/4/4/4
2	LDL	A	801	3	-	1/18/25/25	0/4/4/4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	LDL	C15-N5	-2.58	1.31	1.35
2	B	801	LDL	C15-N6	-2.43	1.31	1.35
2	A	801	LDL	C15-N4	2.29	1.37	1.31
2	A	801	LDL	C15-N6	-2.27	1.32	1.35

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	LDL	C14-N4-C15	4.24	132.21	121.25
2	A	801	LDL	C14-N4-C15	3.56	130.44	121.25
2	B	801	LDL	N6-C15-N5	3.43	111.25	108.94
2	A	801	LDL	O1-C19-N7	-2.94	118.83	122.50
2	A	801	LDL	C7-N1-C18	2.85	132.82	129.19
2	A	801	LDL	N6-C15-N5	2.67	110.74	108.94
2	B	801	LDL	O1-C19-N7	-2.35	119.57	122.50
2	B	801	LDL	C13-C14-N4	2.34	126.70	120.44
2	B	801	LDL	C11-C14-N4	-2.25	114.43	120.44

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	LDL	C13-C14-N4-C15

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	801	LDL	C11-C14-N4-C15

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/365 (97%)	-0.51	5 (1%) 75 75	15, 21, 39, 92	0
1	B	357/365 (97%)	-0.51	6 (1%) 70 69	16, 22, 42, 101	0
All	All	714/730 (97%)	-0.51	11 (1%) 73 73	15, 21, 42, 101	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	771	HIS	6.9
1	B	773	THR	5.0
1	A	771	HIS	4.8
1	B	770	ASP	4.3
1	A	770	ASP	4.0
1	A	459	TRP	3.0
1	B	442	PRO	2.9
1	B	772	LEU	2.5
1	B	777	THR	2.4
1	A	772	LEU	2.2
1	A	798	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	LDL	B	801	28/28	0.89	0.14	26,39,51,52	0
2	LDL	A	801	28/28	0.93	0.12	26,32,50,51	0
3	ZN	A	802	1/1	1.00	0.09	16,16,16,16	0
3	ZN	B	802	1/1	1.00	0.10	17,17,17,17	0
4	K	A	803	1/1	1.00	0.06	19,19,19,19	0
4	K	A	804	1/1	1.00	0.06	15,15,15,15	0
4	K	B	803	1/1	1.00	0.06	17,17,17,17	0
4	K	B	804	1/1	1.00	0.08	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.