



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 08:46 pm BST

PDB ID : 6AAG
Title : Crystal structure of budding yeast Atg8 complexed with the helical AIM of Hfl1.
Authors : Yamasaki, A.; Noda, N.N.
Deposited on : 2018-07-18
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	
1	B	140	
1	C	140	
1	D	140	
1	E	140	
1	F	140	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	140	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into two segments: a green segment on the left labeled '83%' and a yellow segment on the right labeled '14%'. A small red square is at the beginning of the bar, and two small black dots are at the end. A '%' symbol is positioned above the bar.</p>

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 7859 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	137	Total	C	N	O	S	0	0	0
			1087	698	175	211	3			
1	B	138	Total	C	N	O	S	0	0	0
			1119	722	179	214	4			
1	C	139	Total	C	N	O	S	0	0	0
			1103	710	178	212	3			
1	D	137	Total	C	N	O	S	0	0	0
			1092	703	177	209	3			
1	E	136	Total	C	N	O	S	0	0	0
			1074	694	173	204	3			
1	F	138	Total	C	N	O	S	0	0	0
			1104	713	180	207	4			
1	G	138	Total	C	N	O	S	0	0	0
			1085	698	176	208	3			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	GLY	-	expression tag	UNP P36142
A	-22	PRO	-	expression tag	UNP P36142
A	26	PRO	LYS	engineered mutation	UNP A6ZKM4
B	-23	GLY	-	expression tag	UNP P36142
B	-22	PRO	-	expression tag	UNP P36142
B	26	PRO	LYS	engineered mutation	UNP A6ZKM4
C	-23	GLY	-	expression tag	UNP P36142
C	-22	PRO	-	expression tag	UNP P36142
C	26	PRO	LYS	engineered mutation	UNP A6ZKM4
D	-23	GLY	-	expression tag	UNP P36142
D	-22	PRO	-	expression tag	UNP P36142
D	26	PRO	LYS	engineered mutation	UNP A6ZKM4
E	-23	GLY	-	expression tag	UNP P36142
E	-22	PRO	-	expression tag	UNP P36142

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	26	PRO	LYS	engineered mutation	UNP A6ZKM4
F	-23	GLY	-	expression tag	UNP P36142
F	-22	PRO	-	expression tag	UNP P36142
F	26	PRO	LYS	engineered mutation	UNP A6ZKM4
G	-23	GLY	-	expression tag	UNP P36142
G	-22	PRO	-	expression tag	UNP P36142
G	26	PRO	LYS	engineered mutation	UNP A6ZKM4


- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	33	Total O 33 33	0	0
2	B	33	Total O 33 33	0	0
2	C	28	Total O 28 28	0	0
2	D	22	Total O 22 22	0	0
2	E	24	Total O 24 24	0	0
2	F	27	Total O 27 27	0	0
2	G	28	Total O 28 28	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8

Chain A: 




- Molecule 1: Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8

Chain B: 




- Molecule 1: Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8

Chain C: 




- Molecule 1: Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8

Chain D: 




- Molecule 1: Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8


Chain E: 



- Molecule 1: Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8



- Molecule 1: Transmembrane protein 184 homolog YKR051W, Autophagy-related protein 8



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.47Å 61.65Å 100.11Å 90.00° 96.37° 90.00°	Depositor
Resolution (Å)	45.73 – 2.44 46.47 – 2.44	Depositor EDS
% Data completeness (in resolution range)	99.0 (45.73-2.44) 99.1 (46.47-2.44)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.45Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.193 , 0.244 0.193 , 0.244	Depositor DCC
R_{free} test set	2011 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	47.7	Xtrriage
Anisotropy	0.416	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7859	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/1116	0.55	1/1521 (0.1%)
1	B	0.31	0/1150	0.52	0/1561
1	C	0.33	0/1133	0.55	1/1541 (0.1%)
1	D	0.31	0/1121	0.56	1/1525 (0.1%)
1	E	0.28	0/1103	0.49	0/1501
1	F	0.30	0/1134	0.53	0/1540
1	G	0.31	0/1115	0.54	0/1521
All	All	0.31	0/7872	0.53	3/10710 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	-22	PRO	N-CA-CB	5.99	110.48	103.30
1	C	-22	PRO	N-CA-CB	5.90	110.38	103.30
1	D	-22	PRO	N-CA-CB	5.85	110.32	103.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	-10	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1087	0	1024	23	0
1	B	1119	0	1085	25	0
1	C	1103	0	1043	24	0
1	D	1092	0	1044	16	0
1	E	1074	0	1027	16	0
1	F	1104	0	1077	22	0
1	G	1085	0	1027	14	0
2	A	33	0	0	4	0
2	B	33	0	0	3	0
2	C	28	0	0	3	0
2	D	22	0	0	0	0
2	E	24	0	0	1	0
2	F	27	0	0	1	0
2	G	28	0	0	0	0
All	All	7859	0	7327	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:4:THR:HG22	1:G:7:SER:H	1.46	0.81
1:G:31:VAL:HG22	1:G:49:TYR:HB2	1.63	0.80
1:C:45:ASP:N	1:C:45:ASP:OD1	2.23	0.71
1:G:44:ILE:HD13	1:G:67:ARG:HG3	1.72	0.70
1:A:-6:GLU:OE1	2:A:201:HOH:O	2.11	0.69
1:C:-17:GLU:O	2:C:201:HOH:O	2.10	0.69
1:A:-11:GLN:O	2:A:202:HOH:O	2.11	0.67
1:E:-7:PRO:O	2:E:201:HOH:O	2.12	0.67
1:A:0:VAL:HG12	1:B:84:LEU:HD13	1.76	0.67
1:F:113:ASN:O	1:F:113:ASN:ND2	2.25	0.66
1:F:52:PRO:HB2	1:F:55:LEU:HD13	1.78	0.64
1:C:8:GLU:CD	1:C:47:ARG:HH22	2.04	0.60
1:C:81:ASN:ND2	2:C:203:HOH:O	2.35	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:ASP:OD1	1:B:45:ASP:N	2.35	0.57
1:A:-7:PRO:HG2	1:B:62:TYR:CE2	2.40	0.57
1:E:25:PHE:CG	1:E:50:LEU:HD21	2.41	0.56
1:B:82:ASP:OD2	2:B:201:HOH:O	2.18	0.55
1:C:8:GLU:OE2	1:C:47:ARG:NH2	2.37	0.55
1:A:97:GLU:OE2	1:F:67:ARG:HD2	2.07	0.55
1:A:97:GLU:HG2	1:F:67:ARG:HH11	1.72	0.55
1:A:-6:GLU:OE2	1:B:87:THR:OG1	2.25	0.54
1:G:8:GLU:CD	1:G:47:ARG:HH22	2.10	0.54
1:A:52:PRO:HB2	1:A:55:LEU:HD13	1.89	0.54
1:C:1:MET:CB	1:E:20:ARG:HG2	2.38	0.54
1:A:114:THR:OG1	2:A:203:HOH:O	2.19	0.53
1:F:15:LYS:HD3	1:F:102:ASP:HA	1.90	0.53
1:B:-7:PRO:HG2	1:C:62:TYR:CD2	2.45	0.52
1:C:29:ILE:HD12	1:C:53:ALA:HA	1.92	0.51
1:D:52:PRO:HB2	1:D:55:LEU:HD13	1.92	0.51
1:D:77:PHE:CE1	1:E:-1:PRO:HG2	2.45	0.51
1:C:93:ALA:O	1:C:97:GLU:HG2	2.12	0.50
1:B:6:LYS:NZ	1:B:100:ASP:OD2	2.44	0.49
1:D:62:TYR:CE2	1:E:-7:PRO:HG2	2.47	0.49
1:G:45:ASP:N	1:G:45:ASP:OD1	2.41	0.49
1:D:95:TYR:CE2	1:D:99:LYS:HD3	2.48	0.48
1:A:62:TYR:CE2	1:F:-7:PRO:HG2	2.49	0.48
1:B:65:ARG:HH21	1:B:76:ILE:HG22	1.79	0.48
1:E:45:ASP:OD1	1:E:45:ASP:N	2.39	0.48
1:A:-7:PRO:HG2	1:B:62:TYR:CD2	2.49	0.47
1:A:14:ARG:NH2	2:A:208:HOH:O	2.46	0.47
1:G:78:ILE:O	1:G:84:LEU:HD12	2.15	0.47
1:A:45:ASP:OD1	1:A:45:ASP:N	2.30	0.47
1:G:78:ILE:HA	1:G:108:THR:O	2.14	0.47
1:B:53:ALA:HB1	1:B:92:SER:HB3	1.97	0.46
1:A:84:LEU:HD13	1:F:0:VAL:HG12	1.97	0.46
1:C:62:TYR:CE2	1:C:66:LYS:HE2	2.51	0.46
1:E:29:ILE:CD1	1:E:53:ALA:HA	2.46	0.46
1:A:25:PHE:CD2	1:A:50:LEU:HD21	2.50	0.46
1:B:28:ARG:HB3	1:B:50:LEU:HD22	1.97	0.46
1:B:-18:TRP:CE2	1:C:48:LYS:HD3	2.51	0.46
1:E:-14:ILE:O	1:E:-8:PHE:HE2	1.99	0.46
1:B:6:LYS:HG3	1:B:14:ARG:CZ	2.46	0.45
1:E:25:PHE:CD2	1:E:50:LEU:HD21	2.51	0.45
1:C:10:PRO:HG3	1:G:-17:GLU:OE2	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:33:CYS:HA	1:F:107:VAL:O	2.15	0.45
1:B:-7:PRO:HG2	1:C:62:TYR:CE2	2.50	0.45
1:F:45:ASP:N	1:F:45:ASP:OD1	2.42	0.45
1:G:95:TYR:HD1	1:G:105:LEU:HB2	1.82	0.45
1:C:31:VAL:HG13	1:C:107:VAL:HG13	1.99	0.45
1:D:97:GLU:OE2	1:E:67:ARG:HD2	2.17	0.45
1:G:43:GLU:HA	1:G:109:TYR:OH	2.16	0.45
1:C:14:ARG:NH2	2:C:207:HOH:O	2.50	0.45
1:E:78:ILE:HA	1:E:108:THR:O	2.17	0.44
1:A:97:GLU:HG2	1:F:67:ARG:NH1	2.33	0.44
1:F:73:GLU:CD	1:F:73:GLU:H	2.21	0.44
1:B:-1:PRO:HG2	1:C:77:PHE:CE1	2.53	0.44
1:D:73:GLU:H	1:D:73:GLU:CD	2.21	0.43
1:A:62:TYR:CD2	1:F:-7:PRO:HG2	2.53	0.43
1:B:-4:PRO:HD3	1:C:62:TYR:CD1	2.54	0.43
1:B:50:LEU:HA	1:B:50:LEU:HD23	1.88	0.43
1:F:-14:ILE:O	1:F:-11:GLN:HG2	2.19	0.43
1:C:29:ILE:CD1	1:C:53:ALA:HA	2.49	0.43
1:D:78:ILE:HA	1:D:108:THR:O	2.18	0.43
1:F:65:ARG:NH2	1:F:76:ILE:HG22	2.34	0.43
1:G:93:ALA:O	1:G:97:GLU:HG2	2.19	0.43
1:B:60:PHE:CZ	1:B:64:ILE:HD11	2.53	0.43
1:C:10:PRO:HG3	1:G:-17:GLU:CD	2.39	0.43
1:B:20:ARG:NH2	2:B:202:HOH:O	2.27	0.42
1:C:73:GLU:H	1:C:73:GLU:CD	2.23	0.42
1:A:71:PRO:HB2	1:A:73:GLU:OE1	2.19	0.42
1:B:44:ILE:HG12	1:B:67:ARG:HG3	2.00	0.42
1:C:35:LYS:HD3	1:C:43:GLU:HG3	2.01	0.42
1:G:-11:GLN:HB2	1:G:-8:PHE:CZ	2.54	0.42
1:F:41:ILE:HD11	1:F:70:LEU:HD22	2.01	0.42
1:A:35:LYS:HG2	1:A:36:ALA:O	2.20	0.42
1:B:73:GLU:CD	1:B:73:GLU:H	2.23	0.42
1:F:95:TYR:CE2	1:F:99:LYS:HD3	2.55	0.42
1:A:67:ARG:HD2	1:B:97:GLU:OE1	2.20	0.42
1:D:46:LYS:HD3	1:D:49:TYR:CZ	2.55	0.42
1:F:25:PHE:CD2	1:F:50:LEU:HD21	2.55	0.42
1:F:28:ARG:NH1	2:F:203:HOH:O	2.47	0.42
1:C:97:GLU:HG3	1:C:98:HIS:CD2	2.55	0.41
1:B:-2:TYR:HA	1:B:-1:PRO:HD3	1.91	0.41
1:D:29:ILE:CD1	1:D:53:ALA:HA	2.50	0.41
1:D:17:GLU:HG3	1:D:20:ARG:NH2	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:34:GLU:O	1:F:108:THR:HA	2.20	0.41
1:B:15:LYS:HE3	1:B:19:GLU:OE2	2.20	0.41
1:C:41:ILE:HD13	1:C:109:TYR:HD2	1.85	0.41
1:D:50:LEU:HD12	1:D:50:LEU:HA	1.84	0.41
1:A:5:PHE:CE2	1:A:14:ARG:HG2	2.56	0.41
1:A:29:ILE:CD1	1:A:53:ALA:HA	2.51	0.41
1:B:14:ARG:NH2	2:B:206:HOH:O	2.53	0.41
1:D:51:VAL:HG11	1:D:91:MET:HG3	2.02	0.41
1:A:50:LEU:HD12	1:A:50:LEU:HA	1.89	0.41
1:D:66:LYS:HE2	1:D:66:LYS:HB3	1.82	0.41
1:D:65:ARG:NH2	1:D:76:ILE:HG22	2.36	0.41
1:E:8:GLU:CD	1:E:47:ARG:HH22	2.24	0.41
1:E:46:LYS:HG3	1:G:-20:GLU:OE2	2.21	0.41
1:E:73:GLU:CD	1:E:73:GLU:H	2.23	0.40
1:D:86:PRO:HB2	1:E:43:GLU:OE2	2.21	0.40
1:C:41:ILE:HD13	1:C:109:TYR:CD2	2.56	0.40
1:D:48:LYS:HD3	1:E:-18:TRP:CE2	2.57	0.40
1:F:100:ASP:HB3	1:F:102:ASP:OD1	2.21	0.40
1:F:65:ARG:HH21	1:F:76:ILE:HG22	1.86	0.40
1:F:79:PHE:HZ	1:F:113:ASN:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/140 (96%)	128 (95%)	7 (5%)	0	100	100
1	B	136/140 (97%)	132 (97%)	4 (3%)	0	100	100
1	C	137/140 (98%)	132 (96%)	5 (4%)	0	100	100
1	D	135/140 (96%)	129 (96%)	6 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	132/140 (94%)	128 (97%)	4 (3%)	0	100	100
1	F	136/140 (97%)	133 (98%)	3 (2%)	0	100	100
1	G	136/140 (97%)	133 (98%)	3 (2%)	0	100	100
All	All	947/980 (97%)	915 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/125 (92%)	114 (99%)	1 (1%)	78	87
1	B	122/125 (98%)	120 (98%)	2 (2%)	62	74
1	C	116/125 (93%)	116 (100%)	0	100	100
1	D	116/125 (93%)	114 (98%)	2 (2%)	60	73
1	E	113/125 (90%)	112 (99%)	1 (1%)	78	87
1	F	118/125 (94%)	115 (98%)	3 (2%)	47	60
1	G	114/125 (91%)	112 (98%)	2 (2%)	59	71
All	All	814/875 (93%)	803 (99%)	11 (1%)	67	78

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	THR
1	B	82	ASP
1	B	87	THR
1	D	46	LYS
1	D	69	MET
1	E	0	VAL
1	F	69	MET
1	F	113	ASN
1	F	114	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	4	THR
1	G	31	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	137/140 (97%)	-0.15	0 100 100	36, 51, 73, 86	0
1	B	138/140 (98%)	-0.10	1 (0%) 87 87	34, 49, 75, 95	0
1	C	139/140 (99%)	-0.06	1 (0%) 87 87	31, 50, 75, 90	0
1	D	137/140 (97%)	0.01	0 100 100	32, 52, 77, 91	0
1	E	136/140 (97%)	-0.01	0 100 100	38, 56, 87, 104	0
1	F	138/140 (98%)	-0.05	1 (0%) 87 87	34, 52, 77, 109	0
1	G	138/140 (98%)	-0.05	2 (1%) 75 73	38, 55, 76, 102	0
All	All	963/980 (98%)	-0.06	5 (0%) 91 91	31, 52, 79, 109	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	-23	GLY	4.6
1	G	113	ASN	2.8
1	G	76	ILE	2.7
1	C	1	MET	2.2
1	B	76	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.