



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 07:10 am BST

PDB ID : 1AD9
Title : IGG-FAB FRAGMENT OF ENGINEERED HUMAN MONOCLONAL AN-TIBODY CTM01
Authors : Banfield, M.J.; Brady, R.L.
Deposited on : 1997-02-24
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	
1	L	219	
2	B	219	
2	H	219	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6760 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG CTM01 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	219	1696	1063	284	340	9	0	0	0
1	A	219	1696	1063	284	340	9	0	0	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	25	SER	ALA	CONFLICT	GB 468243
L	27	LYS	GLN	CONFLICT	GB 468243
L	27B	LEU	-	INSERTION	GB 468243
L	27C	LEU	-	INSERTION	GB 468243
L	27D	HIS	ILE	CONFLICT	GB 468243
L	28	ASN	-	INSERTION	GB 468243
L	29	GLY	-	INSERTION	GB 468243
L	30	ASP	-	INSERTION	GB 468243
L	31	THR	PRO	CONFLICT	GB 468243
L	32	PHE	TRP	CONFLICT	GB 468243
L	34	TYR	PRO	CONFLICT	GB 468243
L	36	PHE	TYR	CONFLICT	GB 468243
L	48	MET	ILE	CONFLICT	GB 468243
L	50	ARG	LYS	CONFLICT	GB 468243
L	51	MET	ALA	CONFLICT	GB 468243
L	53	ASN	SER	CONFLICT	GB 468243
L	55	ALA	GLU	CONFLICT	GB 468243
L	76	SER	THR	CONFLICT	GB 468243
L	87	TYR	PHE	CONFLICT	GB 468243
L	89	MET	-	INSERTION	GB 468243
L	92	LEU	TYR	CONFLICT	GB 468243
L	93	GLU	ASN	CONFLICT	GB 468243
L	94	TYR	ARG	CONFLICT	GB 468243
L	96	PHE	TRP	CONFLICT	GB 468243
L	106	VAL	ILE	CONFLICT	GB 468243

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	SER	ALA	CONFLICT	GB 468243
A	27	LYS	GLN	CONFLICT	GB 468243
A	27B	LEU	-	INSERTION	GB 468243
A	27C	LEU	-	INSERTION	GB 468243
A	27D	HIS	ILE	CONFLICT	GB 468243
A	28	ASN	-	INSERTION	GB 468243
A	29	GLY	-	INSERTION	GB 468243
A	30	ASP	-	INSERTION	GB 468243
A	31	THR	PRO	CONFLICT	GB 468243
A	32	PHE	TRP	CONFLICT	GB 468243
A	34	TYR	PRO	CONFLICT	GB 468243
A	36	PHE	TYR	CONFLICT	GB 468243
A	48	MET	ILE	CONFLICT	GB 468243
A	50	ARG	LYS	CONFLICT	GB 468243
A	51	MET	ALA	CONFLICT	GB 468243
A	53	ASN	SER	CONFLICT	GB 468243
A	55	ALA	GLU	CONFLICT	GB 468243
A	76	SER	THR	CONFLICT	GB 468243
A	87	TYR	PHE	CONFLICT	GB 468243
A	89	MET	-	INSERTION	GB 468243
A	92	LEU	TYR	CONFLICT	GB 468243
A	93	GLU	ASN	CONFLICT	GB 468243
A	94	TYR	ARG	CONFLICT	GB 468243
A	96	PHE	TRP	CONFLICT	GB 468243
A	106	VAL	ILE	CONFLICT	GB 468243

- Molecule 2 is a protein called IGG CTM01 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	219	Total	C	N	O	S	0	0	0
			1674	1057	273	336	8			
2	B	219	Total	C	N	O	S	0	0	0
			1674	1057	273	336	8			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	3	Total	O	0	0
			3	3		
4	H	6	Total	O	0	0
			6	6		
4	A	2	Total	O	0	0
			2	2		
4	B	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: IGG CTM01 FAB (LIGHT CHAIN)




- Molecule 1: IGG CTM01 FAB (LIGHT CHAIN)



- Molecule 2: IGG CTM01 FAB (HEAVY CHAIN)



- Molecule 2: IGG CTM01 FAB (HEAVY CHAIN)





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.39Å 104.63Å 119.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80 19.90 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (15.00-2.80) 99.8 (19.90-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.45 (at 2.79Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.233 , 0.284 0.228 , 0.270	Depositor DCC
R_{free} test set	1306 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	46.0	Xtrriage
Anisotropy	0.214	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6760	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1734	0.63	0/2348
1	L	0.42	0/1734	0.65	1/2348 (0.0%)
2	B	0.43	0/1716	0.64	0/2336
2	H	0.43	0/1716	0.64	0/2336
All	All	0.42	0/6900	0.64	1/9368 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	213	CYS	N-CA-C	6.16	127.63	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1696	0	1646	15	0
1	L	1696	0	1646	22	0
2	B	1674	0	1620	31	0
2	H	1674	0	1620	27	0
3	B	5	0	0	1	0
4	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	4	0	0	0	0
4	H	6	0	0	0	0
4	L	3	0	0	0	0
All	All	6760	0	6532	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:32:TYR:CD2	2:H:96:LYS:HA	2.18	0.77
1:L:21:ILE:HG23	1:L:102:THR:HG21	1.71	0.71
1:L:35:TRP:HB2	1:L:48:MET:HB2	1.74	0.69
1:A:21:ILE:HG23	1:A:102:THR:HG21	1.73	0.69
1:A:35:TRP:HB2	1:A:48:MET:HB2	1.75	0.67
2:H:119:PRO:HB3	2:H:145:TYR:HB3	1.79	0.65
2:H:66:ARG:HH11	2:H:66:ARG:HG3	1.62	0.64
2:B:119:PRO:HB3	2:B:145:TYR:HB3	1.81	0.63
1:L:212:GLU:HG2	1:L:212:GLU:O	2.01	0.61
2:B:40:ALA:HB3	2:B:43:GLN:CB	2.31	0.60
2:B:66:ARG:HH11	2:B:66:ARG:HG3	1.66	0.60
1:L:189:LYS:HB3	1:L:209:ASN:HD21	1.67	0.59
2:B:43:GLN:HG3	2:B:44:GLY:H	1.67	0.59
1:L:21:ILE:CG2	1:L:102:THR:HG21	2.33	0.58
2:B:96:LYS:O	2:B:100(A):TYR:HA	2.03	0.57
2:B:87:THR:HG23	2:B:110:THR:HA	1.87	0.57
1:A:189:LYS:HB3	1:A:209:ASN:HD21	1.69	0.57
2:B:28:THR:HA	3:B:213:SO4:O4	2.05	0.56
1:A:21:ILE:CG2	1:A:102:THR:HG21	2.34	0.56
2:B:51:ILE:O	2:B:52(A):PRO:HD3	2.06	0.55
2:B:40:ALA:HB3	2:B:43:GLN:HB2	1.89	0.54
1:L:27(B):LEU:HA	1:L:92:LEU:HD22	1.90	0.53
2:H:2:ILE:N	2:H:2:ILE:HD12	2.24	0.53
1:A:49:TYR:O	1:A:53:ASN:HB2	2.09	0.53
2:H:168:ALA:HA	2:H:178:LEU:HB3	1.91	0.52
2:H:51:ILE:O	2:H:52(A):PRO:HD3	2.09	0.52
1:A:27(B):LEU:HA	1:A:92:LEU:HD22	1.92	0.52
2:H:2:ILE:HD12	2:H:2:ILE:H	1.74	0.51
2:H:87:THR:HG23	2:H:110:THR:HA	1.92	0.51
2:B:168:ALA:HA	2:B:178:LEU:HB3	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:18:ARG:HG3	1:L:76:SER:HA	1.94	0.50
2:B:36:TRP:CE2	2:B:80:MET:HB2	2.47	0.50
1:A:21:ILE:HD13	1:A:86:TYR:HB2	1.94	0.49
1:A:91:HIS:CE1	2:B:100(A):TYR:HB2	2.46	0.49
1:A:18:ARG:HG3	1:A:76:SER:HA	1.95	0.49
2:B:139:GLY:HA2	2:B:154:TRP:CH2	2.47	0.49
2:H:139:GLY:HA2	2:H:154:TRP:CH2	2.48	0.49
1:L:49:TYR:O	1:L:53:ASN:HB2	2.12	0.49
2:H:163:VAL:HG22	2:H:182:VAL:HG22	1.94	0.49
1:L:21:ILE:HD13	1:L:86:TYR:HB2	1.95	0.48
2:H:39:GLN:HB2	2:H:45:LEU:HD13	1.94	0.48
2:H:159:LEU:HD21	2:H:182:VAL:HG11	1.96	0.48
2:B:163:VAL:HG22	2:B:182:VAL:HG22	1.96	0.47
2:B:39:GLN:HB2	2:B:45:LEU:HD13	1.97	0.47
2:H:62:LYS:HD2	2:H:62:LYS:N	2.29	0.47
2:H:32:TYR:CD1	2:H:32:TYR:N	2.83	0.47
2:H:32:TYR:HD2	2:H:96:LYS:HA	1.74	0.47
2:B:62:LYS:HD2	2:B:62:LYS:N	2.30	0.46
1:A:2:ILE:HB	1:A:90:GLN:NE2	2.30	0.46
2:B:32:TYR:N	2:B:32:TYR:CD1	2.83	0.46
1:L:210:ARG:HG3	1:L:210:ARG:HH11	1.79	0.46
1:L:188:HIS:O	1:L:210:ARG:NH1	2.49	0.46
2:H:36:TRP:CE2	2:H:80:MET:HB2	2.51	0.45
1:L:112:PRO:HB3	1:L:138:PHE:HB3	1.98	0.45
2:H:147:PRO:O	2:H:200:HIS:HE1	1.99	0.45
2:H:59:TYR:CE1	2:H:69:LEU:HG	2.52	0.45
2:B:147:PRO:O	2:B:200:HIS:HE1	1.99	0.45
1:L:123:GLN:HG2	1:L:128:THR:O	2.17	0.45
2:B:159:LEU:HD21	2:B:182:VAL:HG11	1.98	0.45
1:L:189:LYS:HB3	1:L:209:ASN:ND2	2.31	0.44
2:H:66:ARG:NH1	2:H:66:ARG:HG3	2.30	0.44
1:A:118:PRO:HD2	2:B:125:ALA:O	2.17	0.44
1:A:112:PRO:HB3	1:A:138:PHE:HB3	1.98	0.44
2:B:47:TRP:CZ2	2:B:49:GLY:HA2	2.53	0.44
1:L:189:LYS:HA	1:L:210:ARG:HB2	1.99	0.44
1:L:2:ILE:HB	1:L:90:GLN:NE2	2.33	0.44
2:H:105:GLN:HG3	2:H:105:GLN:H	1.58	0.44
1:A:118:PRO:HG3	1:A:213:CYS:SG	2.58	0.43
2:B:96:LYS:HE3	2:B:98:THR:O	2.18	0.43
1:A:189:LYS:HB3	1:A:209:ASN:ND2	2.32	0.43
2:B:40:ALA:HB3	2:B:43:GLN:HB3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ARG:NH1	1:A:82:ASP:OD1	2.52	0.42
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.54	0.42
2:B:37:MET:HE2	2:B:37:MET:HB2	1.79	0.42
2:B:178:LEU:HD12	2:B:178:LEU:C	2.40	0.42
1:L:161:SER:OG	2:H:167:PRO:HG2	2.20	0.42
2:H:1:GLU:O	2:H:1:GLU:HG2	2.19	0.42
2:H:105:GLN:HE21	2:H:105:GLN:HB2	1.64	0.41
1:L:158:SER:HA	1:L:177:THR:O	2.21	0.41
1:L:148:LYS:NZ	1:L:153:LEU:HD11	2.35	0.41
2:B:124:LEU:HD12	2:B:124:LEU:HA	1.92	0.41
2:B:66:ARG:NH1	2:B:66:ARG:HG3	2.34	0.41
2:B:97:THR:HG22	2:B:100(A):TYR:CE2	2.56	0.41
1:L:174:LEU:HD23	1:L:175:SER:N	2.36	0.41
2:B:6:GLN:HA	2:B:21:SER:O	2.21	0.41
2:B:200:HIS:HD2	2:B:203:SER:OG	2.04	0.40
2:H:35:ASN:O	2:H:92:CYS:HA	2.21	0.40
1:L:21:ILE:HD12	1:L:73:LEU:HD23	2.03	0.40
1:L:61:ARG:CZ	1:L:79:GLN:HG3	2.51	0.40
2:B:59:TYR:CE1	2:B:69:LEU:HG	2.57	0.40
2:H:6:GLN:HA	2:H:21:SER:O	2.22	0.40
2:H:95:GLU:O	2:H:97:THR:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/219 (99%)	210 (97%)	5 (2%)	2 (1%)	17	46
1	L	217/219 (99%)	208 (96%)	6 (3%)	3 (1%)	11	34
2	B	217/219 (99%)	204 (94%)	9 (4%)	4 (2%)	8	28
2	H	217/219 (99%)	204 (94%)	9 (4%)	4 (2%)	8	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	868/876 (99%)	826 (95%)	29 (3%)	13 (2%)	10 33

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	137	ASN
2	H	188	SER
1	A	137	ASN
2	B	188	SER
1	L	210	ARG
2	H	144	ASP
2	B	144	ASP
1	L	68	GLY
2	H	41	PRO
2	H	149	PRO
2	B	149	PRO
1	A	68	GLY
2	B	147	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	195/195 (100%)	188 (96%)	7 (4%)	35 69
1	L	195/195 (100%)	188 (96%)	7 (4%)	35 69
2	B	188/188 (100%)	179 (95%)	9 (5%)	25 58
2	H	188/188 (100%)	177 (94%)	11 (6%)	19 49
All	All	766/766 (100%)	732 (96%)	34 (4%)	28 61

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	28	ASN
1	L	72	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	90	GLN
1	L	100	GLN
1	L	105	GLU
1	L	141	ARG
1	L	210	ARG
2	H	1	GLU
2	H	32	TYR
2	H	62	LYS
2	H	82	LEU
2	H	105	GLN
2	H	115	SER
2	H	124	LEU
2	H	133	GLU
2	H	149	PRO
2	H	195	THR
2	H	197	ASN
1	A	28	ASN
1	A	72	THR
1	A	90	GLN
1	A	100	GLN
1	A	105	GLU
1	A	141	ARG
1	A	213	CYS
2	B	32	TYR
2	B	62	LYS
2	B	82	LEU
2	B	105	GLN
2	B	124	LEU
2	B	133	GLU
2	B	149	PRO
2	B	195	THR
2	B	197	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	28	ASN
1	L	90	GLN
1	L	209	ASN
2	H	164	HIS
2	H	200	HIS
1	A	28	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	90	GLN
1	A	91	HIS
1	A	159	GLN
1	A	209	ASN
2	B	171	GLN
2	B	200	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	213	-	4,4,4	0.84	0	6,6,6	0.49	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	213	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/219 (100%)	-0.35	1 (0%) 91 88	14, 34, 60, 96	0
1	L	219/219 (100%)	-0.18	4 (1%) 68 61	14, 37, 64, 100	0
2	B	219/219 (100%)	-0.20	8 (3%) 41 31	12, 29, 80, 100	0
2	H	219/219 (100%)	-0.19	7 (3%) 47 37	12, 31, 83, 100	0
All	All	876/876 (100%)	-0.23	20 (2%) 60 51	12, 33, 69, 100	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	131	THR	6.5
2	B	187	SER	4.3
2	H	128	SER	4.2
2	H	131	THR	3.8
2	B	132	SER	3.3
2	H	189	LEU	3.3
2	H	65	GLY	3.3
2	B	129	ARG	3.2
1	L	209	ASN	3.0
2	H	129	ARG	3.0
1	A	213	CYS	2.8
2	H	190	GLY	2.6
2	H	191	THR	2.5
2	B	130	SER	2.4
2	B	191	THR	2.4
2	B	190	GLY	2.2
2	B	128	SER	2.2
1	L	212	GLU	2.2
1	L	202	SER	2.1
1	L	164	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	B	213	5/5	0.97	0.13	42,43,44,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.