



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 6, 2022 – 04:28 pm GMT

PDB ID : 7ADI
Title : KirBac3.1 W46R: role of a highly conserved tryptophan at the membrane-water interface of Kir channel
Authors : Venien-Bryan, C.; Fagnen, C.; De Zorzi, R.; Bannwarth, L.; Oubella, I.; Haouz, A.
Deposited on : 2020-09-15
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.24
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	
1	B	301	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	K	A	1001	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inward rectifier potassium channel Kirbac3.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	2194	1403	390	391	10	0	3	0
1	B	265	2082	1337	368	367	10	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	ARG	TRP	engineered mutation	UNP D9N164
A	296	HIS	-	expression tag	UNP D9N164
A	297	HIS	-	expression tag	UNP D9N164
A	298	HIS	-	expression tag	UNP D9N164
A	299	HIS	-	expression tag	UNP D9N164
A	300	HIS	-	expression tag	UNP D9N164
A	301	HIS	-	expression tag	UNP D9N164
B	46	ARG	TRP	engineered mutation	UNP D9N164
B	296	HIS	-	expression tag	UNP D9N164
B	297	HIS	-	expression tag	UNP D9N164
B	298	HIS	-	expression tag	UNP D9N164
B	299	HIS	-	expression tag	UNP D9N164
B	300	HIS	-	expression tag	UNP D9N164
B	301	HIS	-	expression tag	UNP D9N164

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	K	0	0
			3	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	8	Total	O	0	0
			8	8		
4	B	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Inward rectifier potassium channel Kirbac3.1



- Molecule 1: Inward rectifier potassium channel Kirbac3.1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.77Å 113.98Å 89.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.02 – 2.80 48.02 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.02-2.80) 99.7 (48.02-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0266, BUSTER	Depositor
R, R_{free}	0.222 , 0.287 0.225 , 0.284	Depositor DCC
R_{free} test set	1372 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	89.3	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	4303	wwPDB-VP
Average B, all atoms (Å ²)	105.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2249	0.89	0/3054
1	B	0.71	0/2134	0.89	0/2897
All	All	0.70	0/4383	0.89	0/5951

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	141	GLY	Peptide
1	B	106	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2194	0	2148	41	0
1	B	2082	0	2044	36	0
2	A	3	0	0	0	0
3	A	1	0	0	0	0
4	A	8	0	0	0	0
4	B	15	0	0	0	0
All	All	4303	0	4192	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:THR:HG22	1:B:91:VAL:HG11	1.58	0.84
1:B:244:THR:HG22	1:B:255:HIS:CD2	2.18	0.78
1:A:257:ARG:NH2	1:B:170:GLN:OE1	2.18	0.77
1:B:56:LEU:O	1:B:60:THR:OG1	2.03	0.76
1:A:170:GLN:HE21	1:A:209:SER:HB3	1.55	0.71
1:A:244:THR:HG22	1:A:255[A]:HIS:CD2	2.26	0.69
1:B:44:VAL:HG22	1:B:45:SER:H	1.58	0.69
1:A:71:CYS:O	1:A:74:VAL:CG2	2.42	0.67
1:A:13:ARG:NH2	1:A:264:GLU:OE1	2.29	0.66
1:A:103:ILE:HG23	1:A:104:PRO:HD2	1.78	0.65
1:B:244:THR:CG2	1:B:255:HIS:CD2	2.80	0.64
1:A:224:ILE:HA	1:A:227:GLU:HG3	1.79	0.64
1:B:163:LEU:HD12	1:B:208:PHE:CZ	2.35	0.61
1:B:46:ARG:HB3	1:B:47:PRO:HD3	1.82	0.61
1:A:146:SER:H	1:A:291:HIS:HD2	1.48	0.61
1:B:146:SER:H	1:B:291:HIS:HD2	1.51	0.59
1:A:275:PHE:HA	1:A:284:ALA:O	2.03	0.57
1:B:138:PRO:HB2	1:B:250:PHE:CD2	2.40	0.56
1:A:103:ILE:CG2	1:A:104:PRO:HD2	2.35	0.55
1:A:79:ARG:O	1:A:82:SER:HB3	2.07	0.55
1:A:46:ARG:HB3	1:A:47:PRO:HD3	1.89	0.54
1:A:93:THR:HG21	1:A:113:VAL:HG22	1.91	0.53
1:A:255[A]:HIS:CD2	1:B:172:ILE:CD1	2.92	0.53
1:A:257:ARG:HD2	1:B:207:ILE:HD11	1.91	0.53
1:B:200:LEU:HD13	1:B:214:VAL:HG13	1.91	0.52
1:B:100:GLY:O	1:B:103:ILE:CG2	2.58	0.52
1:B:274:VAL:HG23	1:B:285:LEU:HD12	1.92	0.51
1:B:138:PRO:HB2	1:B:250:PHE:CE2	2.46	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:ALA:HB2	1:A:254:VAL:HB	1.93	0.50
1:A:274:VAL:HG13	1:A:285:LEU:HD12	1.94	0.50
1:B:75:ILE:O	1:B:78:ALA:HB3	2.13	0.49
1:A:51:THR:O	1:A:54:THR:OG1	2.30	0.48
1:A:162:ARG:CZ	1:A:271:PHE:HB3	2.43	0.48
1:A:224:ILE:HA	1:A:227:GLU:CG	2.43	0.48
1:B:148:MET:HE3	1:B:265:ILE:HG12	1.97	0.46
1:A:242:LEU:HD11	1:B:206:PRO:HB2	1.98	0.45
1:A:15:LEU:HD13	1:A:240:LEU:HD13	1.98	0.45
1:B:90:SER:OG	1:B:116:GLU:OE2	2.35	0.45
1:A:114:THR:HG22	1:B:91:VAL:HG21	2.00	0.44
1:A:136:THR:HB	1:A:252:GLN:HE22	1.83	0.44
1:A:210:LEU:HD13	1:A:275:PHE:HE1	1.83	0.44
1:A:143:LEU:O	1:A:163:LEU:HA	2.18	0.44
1:A:50:ILE:O	1:A:54:THR:HG23	2.19	0.43
1:A:146:SER:N	1:A:291:HIS:HD2	2.12	0.43
1:B:110:ASN:O	1:B:114:THR:HG23	2.17	0.43
1:A:74:VAL:HG12	1:A:105:ILE:O	2.18	0.43
1:A:102:LEU:HD12	1:A:102:LEU:HA	1.88	0.43
1:A:201:THR:HG22	1:A:217:PRO:CG	2.49	0.42
1:A:114:THR:CG2	1:B:91:VAL:HG21	2.49	0.42
1:B:178:LEU:HD13	1:B:241:VAL:HG22	2.02	0.42
1:B:229:ASP:OD1	1:B:233:ARG:NE	2.42	0.42
1:B:191:VAL:O	1:B:191:VAL:HG13	2.20	0.41
1:B:25:ARG:HG3	1:B:26:LEU:O	2.19	0.41
1:A:69:LEU:HD21	1:A:82:SER:O	2.20	0.41
1:B:105:ILE:O	1:B:106:GLY:O	2.38	0.41
1:A:240:LEU:HD21	1:B:207:ILE:HD12	2.01	0.41
1:B:247:HIS:ND1	1:B:249:ALA:HB3	2.35	0.41
1:B:146:SER:N	1:B:291:HIS:HD2	2.16	0.41
1:A:79:ARG:HA	1:A:80:PRO:HD3	1.92	0.41
1:A:228:THR:OG1	1:A:231:THR:HG23	2.21	0.41
1:B:275:PHE:HA	1:B:284:ALA:O	2.21	0.41
1:A:242:LEU:HD21	1:B:207:ILE:HG13	2.03	0.41
1:A:170:GLN:NE2	1:A:209:SER:HB3	2.31	0.40
1:A:255[A]:HIS:CD2	1:B:172:ILE:HD11	2.55	0.40
1:A:149:VAL:CG1	1:A:293:ILE:HD13	2.51	0.40
1:B:143:LEU:O	1:B:163:LEU:HA	2.21	0.40
1:B:274:VAL:HG23	1:B:285:LEU:CD1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/301 (91%)	246 (90%)	25 (9%)	2 (1%)	22	53
1	B	259/301 (86%)	228 (88%)	24 (9%)	7 (3%)	5	17
All	All	532/602 (88%)	474 (89%)	49 (9%)	9 (2%)	9	29

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	38	TYR
1	B	45	SER
1	B	106	GLY
1	A	283	ARG
1	B	99	TYR
1	B	41	LEU
1	B	129	SER
1	B	130	LEU
1	B	154	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/256 (92%)	222 (94%)	14 (6%)	19	49
1	B	221/256 (86%)	193 (87%)	28 (13%)	4	14
All	All	457/512 (89%)	415 (91%)	42 (9%)	9	27

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	SER
1	A	45	SER
1	A	74	VAL
1	A	76	GLU
1	A	147	ARG
1	A	163	LEU
1	A	201	THR
1	A	203	SER
1	A	209	SER
1	A	247	HIS
1	A	274	VAL
1	A	277	THR
1	A	282	ARG
1	A	295	GLN
1	B	25	ARG
1	B	42	LEU
1	B	45	SER
1	B	46	ARG
1	B	52	LEU
1	B	53	ILE
1	B	56	LEU
1	B	60	THR
1	B	99	TYR
1	B	135	PHE
1	B	139	THR
1	B	147	ARG
1	B	163	LEU
1	B	185	ILE
1	B	186	SER
1	B	192	PHE
1	B	202	ARG
1	B	204	ARG
1	B	209	SER
1	B	210	LEU
1	B	230	GLU
1	B	240	LEU
1	B	254	VAL
1	B	263	ASP
1	B	272	VAL
1	B	283	ARG
1	B	295	GLN
1	B	299	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	ASN
1	A	77	ASN
1	A	170	GLN
1	A	221	HIS
1	A	252	GLN
1	A	291	HIS
1	A	295	GLN
1	B	61	ASN
1	B	255	HIS
1	B	291	HIS
1	B	295	GLN
1	B	296	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/301 (91%)	0.32	19 (6%) 16 10	56, 100, 159, 193	0
1	B	265/301 (88%)	0.53	29 (10%) 5 3	58, 107, 167, 181	0
All	All	541/602 (89%)	0.42	48 (8%) 9 5	56, 103, 164, 193	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	186	SER	7.9
1	B	108	LEU	5.8
1	B	183	SER	5.5
1	B	188	GLU	5.3
1	B	187	GLN	5.0
1	B	298	HIS	4.5
1	B	191	VAL	4.5
1	B	192	PHE	4.1
1	B	292	GLU	4.0
1	B	185	ILE	4.0
1	B	83	PHE	3.9
1	B	190	MET	3.9
1	A	282	ARG	3.8
1	B	38	TYR	3.7
1	A	283	ARG	3.7
1	B	299	HIS	3.7
1	B	277	THR	3.7
1	B	71	CYS	3.7
1	B	276	THR	3.4
1	B	107	PRO	3.4
1	A	97	ILE	3.0
1	B	300	HIS	2.9
1	A	71	CYS	2.9
1	B	111	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	107	PRO	2.9
1	A	82	SER	2.8
1	A	111	THR	2.8
1	A	153	PHE	2.7
1	A	284	ALA	2.6
1	A	296	HIS	2.6
1	B	289	LYS	2.5
1	A	100	GLY	2.5
1	A	277	THR	2.5
1	A	152	ASP	2.4
1	B	52	LEU	2.4
1	A	98	GLY	2.3
1	A	275	PHE	2.3
1	B	252	GLN	2.3
1	A	108	LEU	2.2
1	B	282	ARG	2.2
1	B	184	GLU	2.2
1	A	83	PHE	2.2
1	B	49	PHE	2.1
1	B	189	GLY	2.1
1	B	235	SER	2.0
1	A	48	VAL	2.0
1	B	121	MET	2.0
1	A	109	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	K	A	1001	1/1	0.58	0.63	110,110,110,110	1
3	MG	A	1004	1/1	0.78	0.11	37,37,37,37	1
2	K	A	1003	1/1	0.86	0.40	106,106,106,106	1
2	K	A	1002	1/1	0.94	0.29	78,78,78,78	1

6.5 Other polymers [i](#)

There are no such residues in this entry.