



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 05:47 PM BST

PDB ID : 1AE8  
Title : HUMAN ALPHA-THROMBIN INHIBITION BY EOC-D-PHE-PRO-AZALYS-ONP  
Authors : De Simone, G.; Balliano, G.; Milla, P.; Gallina, C.; Giordano, C.; Tarricone, C.; Rizzi, M.; Bolognesi, M.; Ascenzi, P.  
Deposited on : 1997-03-06  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	L	36	58% (green), 25% (yellow), 17% (orange)
2	H	259	68% (green), 24% (yellow), 5% (orange), 3% (grey)
3	I	10	50% (green), 50% (yellow)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	H	400	X	-	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2638 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-THROMBIN (SMALL SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	36	287	177	48	61	1	60	0	0

- Molecule 2 is a protein called ALPHA-THROMBIN (LARGE SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	253	2053	1310	362	367	14	87	0	0

- Molecule 3 is a protein called HIRUGEN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	I	10	95	59	10	25	1	23	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	H	1	14	8	1	5	14	0

- Molecule 5 is 1-ETHOXYCARBONYL-D-PHE-PRO-2(4-AMINOBTYL)HYDRAZINE (three-letter code: AZL) (formula:  $C_{21}H_{33}N_5O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	H	1	30	21	5	4	1	0

- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	L	21	Total 21	O 21	0	0
6	H	137	Total 137	O 137	0	0
6	I	1	Total 1	O 1	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.71Å 72.55Å 73.20Å 90.00° 100.39° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 19.84 – 2.00	Depositor EDS
% Data completeness (in resolution range)	88.2 (20.00-2.00) 87.6 (19.84-2.00)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.26 (at 2.01Å)	Xtrriage
Refinement program	TNT 5E	Depositor
R, $R_{free}$	(Not available) , (Not available) 0.161 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.0	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 125.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2638	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AZL, NAG, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	L	1.52	2/290 (0.7%)	1.14	2/384 (0.5%)
2	H	0.93	0/2107	1.01	6/2846 (0.2%)
3	I	2.00	1/79 (1.3%)	0.73	0/103
All	All	1.06	3/2476 (0.1%)	1.02	8/3333 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	15	ARG	C-OXT	18.08	1.57	1.23
3	I	64	LEU	C-OXT	16.35	1.54	1.23
1	L	8	GLU	CD-OE1	6.23	1.32	1.25

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	15	ARG	NE-CZ-NH1	8.82	124.71	120.30
2	H	77(A)	ARG	NE-CZ-NH1	6.66	123.63	120.30
2	H	93	ARG	NE-CZ-NH1	6.17	123.38	120.30
2	H	206	ARG	NE-CZ-NH2	-5.87	117.37	120.30
2	H	148	TRP	N-CA-CB	-5.47	100.76	110.60
2	H	77(A)	ARG	NE-CZ-NH2	-5.25	117.67	120.30
2	H	101	ARG	NE-CZ-NH1	5.11	122.85	120.30
1	L	15	ARG	NE-CZ-NH2	-5.00	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	287	0	278	16	0
2	H	2053	0	2019	64	0
3	I	95	0	72	2	0
4	H	14	0	11	0	0
5	H	30	0	33	3	0
6	H	137	0	0	7	0
6	I	1	0	0	0	0
6	L	21	0	0	1	0
All	All	2638	0	2413	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:148:TRP:HE3	2:H:150:GLY:HA2	1.19	1.06
1:L:14(J):TYR:O	1:L:14(K):ILE:HG13	1.70	0.91
1:L:1(D):GLY:CA	2:H:123:LEU:H	1.95	0.79
2:H:148:TRP:CE3	2:H:150:GLY:HA2	2.12	0.78
1:L:1(F):GLY:HA2	2:H:235:LYS:CE	2.14	0.77
2:H:147:THR:HA	2:H:148:TRP:HD1	1.51	0.76
1:L:1(D):GLY:HA3	2:H:122:CYS:HA	1.68	0.74
2:H:147:THR:HA	2:H:148:TRP:CD1	2.26	0.71
2:H:173:ARG:HG3	6:H:451:HOH:O	1.94	0.68
2:H:191:CYS:HA	5:H:600:AZL:H22A	1.76	0.68
2:H:148:TRP:HE3	2:H:150:GLY:CA	2.04	0.67
1:L:14(J):TYR:C	1:L:14(K):ILE:HG13	2.14	0.67
2:H:147:THR:CA	2:H:148:TRP:HD1	2.10	0.65
6:H:504:HOH:O	3:I:56:PHE:HB2	1.97	0.65
2:H:237:TRP:O	2:H:241:VAL:HG12	1.98	0.64
2:H:73:ARG:HD2	6:H:498:HOH:O	2.00	0.62
2:H:36(A):SER:HA	2:H:37:PRO:C	2.21	0.61
2:H:70:LYS:HE3	2:H:72:SER:O	2.02	0.60
2:H:143:ASN:ND2	6:H:511:HOH:O	2.28	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:205:ASN:ND2	6:H:537:HOH:O	2.33	0.60
2:H:204(B):ASN:C	2:H:204(B):ASN:HD22	2.05	0.59
2:H:35:ARG:HB2	2:H:41:LEU:HD13	1.84	0.58
1:L:10:LYS:NZ	6:L:460:HOH:O	2.36	0.58
2:H:49:ASP:OD2	2:H:111:PRO:HB3	2.03	0.58
2:H:187:ARG:NH2	2:H:221:ASP:O	2.38	0.57
2:H:85:LEU:HD22	2:H:106:MET:HB3	1.87	0.56
2:H:85:LEU:HD13	2:H:106:MET:HE2	1.87	0.56
2:H:17:VAL:O	2:H:188:GLY:HA2	2.06	0.55
2:H:147:THR:C	2:H:148:TRP:HD1	2.10	0.54
2:H:151:GLN:HG2	6:H:480:HOH:O	2.06	0.53
1:L:14(A):LYS:HG2	2:H:23:GLU:OE2	2.09	0.53
2:H:71:HIS:CD2	2:H:154:VAL:CG2	2.93	0.52
1:L:15:ARG:NH2	2:H:204(A):PHE:O	2.42	0.52
2:H:136:GLY:HA3	2:H:199:PHE:CZ	2.44	0.52
2:H:85:LEU:HD13	2:H:106:MET:CE	2.40	0.51
2:H:230:HIS:CG	2:H:233:ARG:HG3	2.46	0.51
2:H:32:MET:HG3	2:H:40:LEU:CD1	2.41	0.50
2:H:49:ASP:O	2:H:111:PRO:HA	2.12	0.50
2:H:71:HIS:NE2	2:H:154:VAL:HG21	2.26	0.50
1:L:1(G):PHE:CG	1:L:1(F):GLY:N	2.79	0.50
5:H:600:AZL:C18	5:H:600:AZL:H122	2.42	0.49
1:L:14(J):TYR:C	1:L:14(K):ILE:CG1	2.82	0.48
2:H:60(B):PRO:N	2:H:60(C):PRO:CD	2.77	0.48
1:L:1(F):GLY:HA2	2:H:235:LYS:HE3	1.91	0.48
2:H:86:GLU:HB3	2:H:107:LYS:HG2	1.95	0.48
1:L:1(D):GLY:HA3	2:H:122:CYS:CA	2.40	0.48
2:H:125:ASP:OD1	2:H:128:THR:N	2.41	0.47
1:L:1(D):GLY:HA3	2:H:123:LEU:H	1.77	0.47
2:H:46:LEU:HD22	2:H:48:SER:O	2.14	0.47
1:L:15:ARG:CZ	2:H:204(A):PHE:O	2.63	0.47
2:H:85:LEU:CD1	2:H:106:MET:CE	2.94	0.46
2:H:85:LEU:HD11	2:H:106:MET:HE1	1.97	0.46
2:H:130:LEU:HD23	2:H:130:LEU:HA	1.77	0.46
2:H:35:ARG:O	2:H:38:GLN:HA	2.17	0.45
2:H:204(B):ASN:ND2	2:H:206:ARG:H	2.15	0.45
3:I:59:ILE:O	3:I:59:ILE:HG13	2.17	0.45
2:H:60(I):THR:HG23	2:H:61:GLU:N	2.31	0.45
2:H:80:GLU:O	2:H:81:LYS:HD2	2.18	0.44
2:H:71:HIS:NE2	2:H:154:VAL:CG2	2.81	0.43
2:H:94:TYR:CZ	2:H:96:TRP:HB3	2.53	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:99:LEU:HD11	5:H:600:AZL:H101	1.99	0.43
1:L:1(D):GLY:HA2	2:H:123:LEU:H	1.78	0.42
2:H:97(A):GLU:HB2	6:H:454:HOH:O	2.19	0.42
2:H:91:HIS:ND1	2:H:92:PRO:HD2	2.35	0.42
2:H:204(B):ASN:HD22	2:H:205:ASN:N	2.17	0.42
2:H:105:LEU:HD12	2:H:241:VAL:HG11	2.02	0.42
2:H:100:ASP:O	2:H:101:ARG:HB2	2.20	0.42
2:H:65:LEU:HD12	2:H:65:LEU:HA	1.79	0.42
2:H:36(A):SER:CA	2:H:37:PRO:C	2.88	0.41
2:H:60(I):THR:CG2	2:H:61:GLU:N	2.83	0.41
1:L:6:LEU:HA	1:L:6:LEU:HD12	1.90	0.41
2:H:60(A):TYR:C	2:H:60(C):PRO:HD2	2.41	0.40
2:H:238:ILE:O	2:H:241:VAL:HG13	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	34/36 (94%)	26 (76%)	6 (18%)	2 (6%)	1	0
2	H	249/259 (96%)	234 (94%)	15 (6%)	0	100	100
3	I	7/10 (70%)	7 (100%)	0	0	100	100
All	All	290/305 (95%)	267 (92%)	21 (7%)	2 (1%)	22	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	14(L)	ASP
1	L	1(G)	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	31/31 (100%)	23 (74%)	8 (26%)	0	0
2	H	221/225 (98%)	198 (90%)	23 (10%)	7	4
3	I	9/9 (100%)	8 (89%)	1 (11%)	6	3
All	All	261/265 (98%)	229 (88%)	32 (12%)	4	2

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	1(C)	GLU
1	L	6	LEU
1	L	14(A)	LYS
1	L	14(D)	ARG
1	L	14(F)	LEU
1	L	14(G)	LEU
1	L	14(K)	ILE
1	L	14(L)	ASP
2	H	41	LEU
2	H	46	LEU
2	H	50	ARG
2	H	60(I)	THR
2	H	61	GLU
2	H	65	LEU
2	H	87	LYS
2	H	107	LYS
2	H	109	LYS
2	H	126	ARG
2	H	127	GLU
2	H	148	TRP
2	H	154	VAL
2	H	173	ARG
2	H	182	CYS
2	H	187	ARG
2	H	192	GLU
2	H	195	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	H	204(B)	ASN
2	H	233	ARG
2	H	236	LYS
2	H	241	VAL
2	H	244	GLN
3	I	58	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	78	ASN
2	H	156	GLN
2	H	204(B)	ASN
2	H	239	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TYS	I	63	3	15,16,17	2.70	2 (13%)	18,22,24	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	I	63	3	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	63	TYS	OH-S	-9.39	1.44	1.58
3	I	63	TYS	OH-CZ	-3.73	1.36	1.42

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	H	400	2	14,14,15	1.09	1 (7%)	17,19,21	4.42	12 (70%)
5	AZL	H	600	-	31,31,31	1.30	1 (3%)	38,39,39	2.23	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	H	400	2	2/2/5/7	4/6/23/26	0/1/1/1
5	AZL	H	600	-	-	5/29/40/40	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	600	AZL	N7-N6	-5.78	1.37	1.41
4	H	400	NAG	C2-N2	2.70	1.50	1.46

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	400	NAG	O5-C5-C6	9.24	121.69	107.20
5	H	600	AZL	C24-O24-C23	8.76	132.64	116.04
5	H	600	AZL	O23-C23-N9	-7.42	112.69	124.85
4	H	400	NAG	O7-C7-N2	-6.40	110.18	121.95
4	H	400	NAG	C3-C4-C5	5.98	120.90	110.24
4	H	400	NAG	O3-C3-C2	5.88	121.63	109.47
4	H	400	NAG	O7-C7-C8	-5.65	111.56	122.06
5	H	600	AZL	O24-C23-N9	5.03	120.72	110.50
4	H	400	NAG	C2-N2-C7	4.45	129.25	122.90
4	H	400	NAG	C1-C2-N2	4.22	117.70	110.49
4	H	400	NAG	O4-C4-C5	3.86	118.88	109.30
4	H	400	NAG	O4-C4-C3	3.74	119.00	110.35
4	H	400	NAG	O6-C6-C5	3.38	122.87	111.29
4	H	400	NAG	O5-C1-C2	2.87	115.82	111.29
4	H	400	NAG	O3-C3-C4	2.60	116.37	110.35
5	H	600	AZL	C5-N6-N7	2.39	113.72	111.27
5	H	600	AZL	C10-C9-N8	2.02	106.03	103.03

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	H	400	NAG	C4
4	H	400	NAG	C2

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	400	NAG	C3-C2-N2-C7
4	H	400	NAG	C8-C7-N2-C2
4	H	400	NAG	O7-C7-N2-C2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
5	H	600	AZL	O23-C23-O24-C24
5	H	600	AZL	N9-C23-O24-C24
5	H	600	AZL	O8-C8-N7-N6
5	H	600	AZL	C3-C4-C5-N6
4	H	400	NAG	O5-C5-C6-O6
5	H	600	AZL	C2-C3-C4-C5


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	600	AZL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.