



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 04:52 am BST

PDB ID : 4AE9
Title : Structure and function of the Human Sperm-Specific Isoform of Protein Kinase A (PKA) Catalytic Subunit C alpha 2
Authors : Hereng, T.H.; Backe, P.H.; Kahmann, J.; Scheich, C.; Bjoras, M.; Skalhegg, B.S.; Rosendal, K.R.
Deposited on : 2012-01-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	 77% 14% •• 6%
1	B	343	 78% 14% • 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5801 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CAMP-DEPENDENT PROTEIN KINASE CATALYTIC SUB-UNIT ALPHA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	324	Total 2698	C 1756	N 452	O 478	P 3	S 9	0	3	0
1	B	324	Total 2739	C 1790	N 456	O 481	P 3	S 9	0	7	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	201	Total 201	O 201	0	0
2	B	163	Total 163	O 163	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CAMP-DEPENDENT PROTEIN KINASE CATALYTIC SUBUNIT ALPHA

Chain A: 



- Molecule 1: CAMP-DEPENDENT PROTEIN KINASE CATALYTIC SUBUNIT ALPHA

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.40Å 89.97Å 115.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.02 – 2.30 71.02 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.2 (71.02-2.30) 96.2 (71.02-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.83 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.198 , 0.264 0.195 , 0.262	Depositor DCC
R_{free} test set	1737 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	22.1	Xtrriage
Anisotropy	0.111	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5801	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.96	0/2745	0.88	5/3694 (0.1%)
1	B	0.94	0/2801	0.83	0/3768
All	All	0.95	0/5546	0.85	5/7462 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	LEU	CA-CB-CG	7.43	132.39	115.30
1	A	175	ASP	CB-CG-OD1	6.25	123.92	118.30
1	A	221[A]	TRP	CA-CB-CG	-5.84	102.60	113.70
1	A	221[B]	TRP	CA-CB-CG	-5.84	102.60	113.70
1	A	144	ARG	NE-CZ-NH2	-5.72	117.44	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2698	0	2704	52	0
1	B	2739	0	2752	56	0
2	A	201	0	0	7	0
2	B	163	0	0	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5801	0	5456	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:LEU:HD12	1:B:120[A]:MET:HE2	1.38	1.04
1:A:146:TYR:HD2	1:A:231:MET:CE	1.81	0.94
1:B:106:LEU:HD13	1:B:120[A]:MET:HE1	1.49	0.94
1:B:106:LEU:CD1	1:B:120[A]:MET:CE	2.49	0.90
1:A:241:ASP:HB2	2:A:2158:HOH:O	1.72	0.88
1:B:106:LEU:CD1	1:B:120[A]:MET:HE2	2.05	0.85
1:A:146:TYR:HD2	1:A:231:MET:HE1	1.44	0.81
1:B:106:LEU:HD13	1:B:120[A]:MET:CE	2.11	0.79
1:A:146:TYR:CD2	1:A:231:MET:HE1	2.18	0.78
1:A:39:HIS:H	1:A:42:GLN:HE21	1.32	0.78
1:B:106:LEU:HD12	1:B:120[A]:MET:CE	2.08	0.78
1:B:128:MET:SD	1:B:231:MET:CE	2.72	0.77
1:B:276:ASP:OD1	1:B:278:THR:HG22	1.85	0.76
1:B:34:ALA:H	1:B:96:GLN:HE22	1.31	0.76
1:A:118:MET:HB3	1:A:120[A]:MET:HE2	1.68	0.75
1:A:16:LYS:O	2:A:2001:HOH:O	2.05	0.73
1:B:276:ASP:CG	1:B:278:THR:HG22	2.08	0.73
1:B:128:MET:SD	1:B:231:MET:HE2	2.30	0.70
1:A:146:TYR:HD2	1:A:231:MET:HE2	1.56	0.70
1:A:221[A]:TRP:HZ3	2:A:2121:HOH:O	1.76	0.69
1:A:106:LEU:HD13	1:A:120[A]:MET:CE	2.23	0.68
1:B:131:HIS:CE1	1:B:135:ILE:HG13	2.28	0.68
1:B:128:MET:SD	1:B:231:MET:HE3	2.34	0.67
1:B:317:LYS:O	2:B:2149:HOH:O	2.13	0.67
1:B:221[A]:TRP:HZ3	2:B:2096:HOH:O	1.77	0.67
1:A:128:MET:HE3	1:A:231:MET:HG2	1.78	0.66
1:A:39:HIS:H	1:A:42:GLN:NE2	1.94	0.65
1:A:134:ARG:HG3	2:A:2110:HOH:O	1.97	0.65
1:A:62:HIS:CE1	1:A:64:GLU:HG3	2.33	0.64
1:B:103:LEU:HD22	1:B:185:PHE:HZ	1.63	0.62
1:A:128:MET:CE	1:A:231:MET:HG2	2.30	0.62
1:B:276:ASP:OD2	1:B:278:THR:HG22	2.01	0.61
1:A:146:TYR:CD2	1:A:231:MET:CE	2.71	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:34:ALA:H	1:B:96:GLN:NE2	1.98	0.59
1:B:128:MET:HE2	1:B:169:PRO:HA	1.84	0.58
1:B:96:GLN:HG2	1:B:96:GLN:O	2.02	0.58
1:A:39:HIS:N	1:A:42:GLN:HE21	2.01	0.58
1:B:133:ARG:HH11	1:B:133:ARG:HG2	1.69	0.57
1:B:253:GLY:HA2	2:B:2124:HOH:O	2.04	0.57
1:A:106:LEU:HD13	1:A:120[A]:MET:HE2	1.86	0.56
1:B:189:LYS:HG2	1:B:191:VAL:HG23	1.87	0.55
1:B:133:ARG:HD2	2:B:2084:HOH:O	2.07	0.54
1:B:295[B]:LYS:HG3	2:B:2141:HOH:O	2.08	0.53
1:B:92:LYS:HG2	1:B:350:PHE:CE2	2.44	0.53
1:B:92:LYS:HE3	2:B:2053:HOH:O	2.09	0.53
1:A:226:VAL:HG13	1:A:237:PRO:HD2	1.90	0.53
1:A:62:HIS:HE1	1:A:64:GLU:HG3	1.74	0.53
1:B:106:LEU:CD1	1:B:120[A]:MET:HE1	2.19	0.52
1:B:27:LEU:HD21	1:B:190:ARG:NH2	2.24	0.52
1:B:240:ALA:HB1	1:B:245:GLN:HB2	1.92	0.52
1:B:118:MET:HB3	1:B:120[A]:MET:CE	2.40	0.51
1:B:211:LEU:O	1:B:213:LYS:N	2.44	0.51
1:B:240:ALA:CB	1:B:245:GLN:HB2	2.40	0.51
1:A:133:ARG:HG3	1:A:133:ARG:NH1	2.25	0.51
1:B:124:PRO:HD2	1:B:176:GLN:OE1	2.11	0.50
1:A:176[A]:GLN:HE22	1:A:318:PHE:HA	1.76	0.50
1:B:131:HIS:CE1	1:B:135:ILE:CG1	2.94	0.50
1:B:221[A]:TRP:CZ3	2:B:2096:HOH:O	2.53	0.50
1:A:173:LEU:HD12	1:A:181:GLN:HB3	1.94	0.50
1:B:133:ARG:NH1	1:B:133:ARG:HG2	2.26	0.50
1:A:74:LEU:N	1:A:74:LEU:HD12	2.26	0.50
1:A:229:TYR:C	1:A:229:TYR:CD1	2.85	0.49
1:A:47:LYS:HE3	1:A:59:LEU:HD12	1.95	0.49
1:A:106:LEU:HD13	1:A:120[A]:MET:HE1	1.92	0.49
1:B:216:ASN:HB2	2:B:2107:HOH:O	2.12	0.49
1:A:284:LEU:HB2	2:A:2174:HOH:O	2.12	0.48
1:B:295[A]:LYS:HD3	2:B:2128:HOH:O	2.13	0.48
1:A:134:ARG:CZ	1:A:134:ARG:HB3	2.43	0.48
1:A:127:GLU:O	1:A:131:HIS:HD2	1.97	0.48
1:A:133:ARG:CG	1:A:133:ARG:HH11	2.28	0.47
1:A:133:ARG:HG3	1:A:133:ARG:HH11	1.77	0.47
1:A:139:SEP:O2P	1:A:142:HIS:HB2	2.14	0.47
1:A:103:LEU:HD22	1:A:185:PHE:HZ	1.79	0.47
1:B:118:MET:HB3	1:B:120[A]:MET:HE3	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:TYR:OH	1:A:321:PRO:HG2	2.15	0.46
1:A:75:ASP:HA	1:A:115:ASN:HD22	1.81	0.46
1:B:103:LEU:HD21	1:B:153:THR:HG23	1.98	0.46
1:B:26:PHE:CG	1:B:160:LEU:HD13	2.51	0.45
1:A:270:ARG:HH11	1:A:270:ARG:HG3	1.81	0.45
1:B:15:VAL:C	1:B:17:GLU:H	2.20	0.45
1:A:204:TYR:N	1:A:204:TYR:CD1	2.84	0.45
1:A:217:LYS:CE	2:A:2119:HOH:O	2.64	0.44
1:A:39:HIS:N	1:A:42:GLN:NE2	2.60	0.44
1:A:160:LEU:O	1:A:190:ARG:HD2	2.16	0.44
1:B:131:HIS:CE1	1:B:314:PHE:CZ	3.05	0.44
1:B:100:PHE:HB3	1:B:103:LEU:HG	2.00	0.44
1:A:103:LEU:HD22	1:A:185:PHE:CZ	2.53	0.44
1:A:176[A]:GLN:NE2	1:A:318:PHE:HA	2.34	0.43
1:A:239:PHE:O	1:A:240:ALA:HB2	2.18	0.43
1:B:242:GLN:HB2	1:B:245:GLN:HG3	1.99	0.43
1:B:15:VAL:O	1:B:17:GLU:N	2.52	0.43
1:A:257:PHE:HE2	1:A:270:ARG:HD3	1.85	0.42
1:A:74:LEU:HB2	1:A:116:LEU:HB2	2.01	0.42
1:B:106:LEU:HA	1:B:120[A]:MET:HE2	2.01	0.42
1:B:51:THR:HA	1:B:56:ARG:HA	2.02	0.42
1:A:266:LYS:O	1:A:270:ARG:HG2	2.20	0.42
1:B:53:SER:O	2:B:2029:HOH:O	2.22	0.41
1:B:17:GLU:OE1	1:B:17:GLU:HA	2.18	0.41
1:A:49:LEU:HA	1:A:49:LEU:HD23	1.86	0.41
1:B:154:PHE:CE2	1:B:220:ASP:HB3	2.55	0.41
1:A:146:TYR:CD2	1:A:231:MET:HE2	2.46	0.41
1:A:338:SEP:O2P	1:A:340:ASN:OD1	2.39	0.41
1:B:111:LYS:HE2	2:B:2014:HOH:O	2.20	0.41
1:A:127:GLU:O	1:A:131:HIS:CD2	2.73	0.41
1:A:93:ARG:HD2	2:A:2080:HOH:O	2.20	0.41
1:B:34:ALA:N	1:B:96:GLN:HE22	2.07	0.41
1:B:236:PRO:HB2	2:B:2111:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/343 (93%)	305 (95%)	15 (5%)	0	100	100
1	B	324/343 (94%)	308 (95%)	13 (4%)	3 (1%)	17	20
All	All	644/686 (94%)	613 (95%)	28 (4%)	3 (0%)	29	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	212	SER
1	B	16	LYS
1	B	211	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/300 (95%)	268 (94%)	18 (6%)	18	24
1	B	291/300 (97%)	280 (96%)	11 (4%)	33	47
All	All	577/600 (96%)	548 (95%)	29 (5%)	25	34

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LYS
1	A	21	LYS
1	A	56	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	59	LEU
1	A	127	GLU
1	A	133	ARG
1	A	134	ARG
1	A	137	ARG
1	A	173	LEU
1	A	190	ARG
1	A	192	LYS
1	A	213	LYS
1	A	221[A]	TRP
1	A	221[B]	TRP
1	A	229	TYR
1	A	263	SER
1	A	284	LEU
1	A	314	PHE
1	B	16	LYS
1	B	17	GLU
1	B	27	LEU
1	B	78	LYS
1	B	92	LYS
1	B	133	ARG
1	B	190	ARG
1	B	198	LEU
1	B	201	THR
1	B	241	ASP
1	B	285	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	42	GLN
1	A	115	ASN
1	A	131	HIS
1	A	260	HIS
1	A	271	ASN
1	A	340	ASN
1	B	68	HIS
1	B	84	GLN
1	B	96	GLN
1	B	115	ASN
1	B	340	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	B	139	1	8,9,10	2.40	4 (50%)	8,12,14	2.51	3 (37%)
1	SEP	A	139	1	8,9,10	2.12	3 (37%)	8,12,14	1.62	2 (25%)
1	SEP	A	338	1	8,9,10	1.33	1 (12%)	8,12,14	2.84	3 (37%)
1	TPO	A	197	1	8,10,11	0.88	0	10,14,16	1.12	0
1	TPO	B	197	-	5,9,11	1.53	1 (20%)	3,11,16	0.67	0
1	SEP	B	338	1	8,9,10	1.50	1 (12%)	8,12,14	1.70	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	139	1	-	0/5/8/10	-
1	SEP	A	139	1	-	3/5/8/10	-
1	SEP	A	338	1	-	3/5/8/10	-
1	TPO	A	197	1	-	0/9/11/13	-
1	TPO	B	197	-	-	2/6/10/13	-
1	SEP	B	338	1	-	3/5/8/10	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	139	SEP	P-O1P	4.29	1.64	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	139	SEP	P-O1P	4.15	1.63	1.50
1	B	197	TPO	OG1-CB	-3.30	1.41	1.45
1	B	139	SEP	P-O2P	3.01	1.66	1.54
1	B	338	SEP	P-O1P	2.91	1.59	1.50
1	B	139	SEP	P-OG	2.78	1.69	1.60
1	A	139	SEP	P-O2P	2.78	1.65	1.54
1	A	338	SEP	P-O1P	2.77	1.59	1.50
1	B	139	SEP	P-O3P	2.53	1.64	1.54
1	A	139	SEP	P-O3P	2.36	1.63	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	338	SEP	OG-CB-CA	6.49	114.46	108.14
1	B	139	SEP	OG-CB-CA	5.33	113.33	108.14
1	B	139	SEP	O3P-P-OG	3.39	115.76	106.73
1	A	139	SEP	OG-CB-CA	3.27	111.33	108.14
1	B	338	SEP	OG-CB-CA	3.15	111.21	108.14
1	A	338	SEP	O2P-P-OG	-2.97	98.82	106.73
1	B	139	SEP	O2P-P-OG	2.86	114.36	106.73
1	B	338	SEP	O3P-P-O2P	2.69	117.91	107.64
1	A	338	SEP	O3P-P-O2P	2.43	116.93	107.64
1	A	139	SEP	O3P-P-OG	2.23	112.66	106.73
1	B	338	SEP	P-OG-CB	-2.18	112.29	118.30

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	139	SEP	CB-OG-P-O2P
1	A	139	SEP	CB-OG-P-O3P
1	A	338	SEP	CA-CB-OG-P
1	A	338	SEP	CB-OG-P-O2P
1	B	197	TPO	O-C-CA-CB
1	B	338	SEP	CA-CB-OG-P
1	B	338	SEP	CB-OG-P-O2P
1	A	139	SEP	N-CA-CB-OG
1	A	338	SEP	N-CA-CB-OG
1	B	338	SEP	N-CA-CB-OG
1	B	197	TPO	CG2-CB-OG1-P

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	139	SEP	1	0
1	A	338	SEP	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/343 (93%)	-0.43	1 (0%) 94 96	7, 19, 41, 69	0
1	B	321/343 (93%)	-0.42	2 (0%) 89 92	6, 19, 44, 64	0
All	All	642/686 (93%)	-0.43	3 (0%) 91 94	6, 19, 43, 69	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	321	PRO	3.4
1	B	18[A]	PHE	3.3
1	B	31	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	B	139	10/11	0.81	0.18	18,20,47,48	0
1	SEP	A	139	10/11	0.81	0.17	27,29,45,47	0
1	TPO	B	197	10/12	0.96	0.12	23,26,28,29	0
1	SEP	A	338	10/11	0.97	0.09	17,22,24,26	0
1	SEP	B	338	10/11	0.98	0.10	13,13,18,19	0
1	TPO	A	197	11/12	0.99	0.08	5,13,15,15	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.