



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 05:46 pm GMT

PDB ID : 5AGX
Title : Bcl-2 alpha beta-1 LINEAR complex
Authors : Smith, B.J.; F Lee, E.; Checco, J.W.; Gellman, S.H.; Fairlie, W.D.
Deposited on : 2015-02-04
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2027 (2.26-2.22)
Clashscore	122126	2170 (2.26-2.22)
Ramachandran outliers	120053	2129 (2.26-2.22)
Sidechain outliers	120020	2130 (2.26-2.22)
RSRZ outliers	108989	1991 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	
2	B	166	
3	C	22	
3	D	22	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APOPTOSIS REGULATOR BCL-2, BCL-2-LIKE PROTEIN 1, APOPTOSIS REGULATOR BCL-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	140	1158	741	198	212	7	0	4	0

- Molecule 2 is a protein called APOPTOSIS REGULATOR BCL-2, BCL-2-LIKE PROTEIN 1, APOPTOSIS REGULATOR BCL-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	141	1169	749	203	211	6	0	3	0

- Molecule 3 is a protein called BCL-2-LIKE PROTEIN 11.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	22	181	123	27	31	0	0	0
3	D	22	187	126	30	31	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	ACE	-	expression tag	UNP O43521
D	-1	ACE	-	expression tag	UNP O43521
C	2	XCP	TRP	engineered mutation	UNP O43521
C	6	AJE	GLU	engineered mutation	UNP O43521
C	9	MH8	ARG	engineered mutation	UNP O43521
C	13	MH8	GLU	engineered mutation	UNP O43521
C	16	XCP	ALA	engineered mutation	UNP O43521
D	2	XCP	TRP	engineered mutation	UNP O43521
D	6	AJE	GLU	engineered mutation	UNP O43521
D	9	MH8	ARG	engineered mutation	UNP O43521

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	13	MH8	GLU	engineered mutation	UNP O43521
D	16	XCP	ALA	engineered mutation	UNP O43521

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	25	Total O 26 26	0	1
4	B	18	Total O 19 19	0	1
4	C	1	Total O 1 1	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.05Å 86.05Å 108.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.01 – 2.24 45.96 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.01-2.24) 99.9 (45.96-2.24)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.58 (at 2.24Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE: 1.9_1692)	Depositor
R, R_{free}	0.188 , 0.245 0.201 , 0.249	Depositor DCC
R_{free} test set	1013 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtrriage
Anisotropy	0.197	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2741	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XCP, AJE, ACE, MH8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1188	0.51	0/1610
2	B	0.36	0/1200	0.48	0/1625
3	C	0.43	0/125	0.50	0/161
3	D	0.33	0/131	0.48	0/168
All	All	0.38	0/2644	0.49	0/3564

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	6
3	D	0	6
All	All	0	12

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	1	ILE	Peptide
3	C	15	ASN	Peptide
3	C	16	XCP	Mainchain
3	C	2	XCP	Mainchain
3	C	5	GLN	Peptide
3	C	6	AJE	Mainchain
3	D	1	ILE	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
3	D	15	ASN	Peptide
3	D	16	XCP	Mainchain
3	D	2	XCP	Mainchain
3	D	5	GLN	Peptide
3	D	6	AJE	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1158	0	1071	6	0
2	B	1169	0	1088	8	0
3	C	181	0	172	3	0
3	D	187	0	183	3	0
4	A	26	0	0	0	0
4	B	19	0	0	0	0
4	C	1	0	0	0	0
All	All	2741	0	2514	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:ASP:OD1	1:A:106:ARG:NH1	2.14	0.81
1:A:122:THR:H	1:A:125:THR:HG22	1.48	0.79
2:B:18:TYR:OH	2:B:152:GLU:OE2	2.08	0.72
2:B:8:GLY:HA2	2:B:195:TRP:HB2	1.78	0.66
2:B:105:SER:OG	2:B:152:GLU:OE1	2.12	0.62
3:C:16:XCP:HD	3:D:13:MH8:HB12	1.81	0.62
1:A:18:TYR:OH	1:A:152:GLU:OE2	2.18	0.62
2:B:179:GLU:HG2	2:B:183:ARG:HH21	1.70	0.55
2:B:114:GLU:OE1	2:B:117:SER:OG	2.25	0.55
3:D:6:AJE:H31C	3:D:9:MH8:HZ2	1.91	0.52
1:A:120:HIS:O	1:A:125:THR:HG21	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:12:ARG:HH22	2:B:175:ILE:HD12	1.80	0.47
2:B:145:GLY:HA3	3:C:15:ASN:HB2	1.98	0.45
1:A:8:GLY:HA2	1:A:195:TRP:HB2	2.01	0.43
1:A:12:ARG:O	1:A:16:MET:HG2	2.19	0.43
2:B:99:GLN:O	2:B:103:ASP:HB2	2.19	0.42
3:C:9:MH8:HB11	3:D:16:XCP:HDA	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/166 (84%)	137 (98%)	2 (1%)	1 (1%)	24	22
2	B	140/166 (84%)	138 (99%)	2 (1%)	0	100	100
3	C	15/22 (68%)	15 (100%)	0	0	100	100
3	D	15/22 (68%)	15 (100%)	0	0	100	100
All	All	310/376 (82%)	305 (98%)	4 (1%)	1 (0%)	43	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/141 (84%)	116 (98%)	2 (2%)	63	71
2	B	119/141 (84%)	116 (98%)	3 (2%)	50	58
3	C	11/13 (85%)	11 (100%)	0	100	100
3	D	12/13 (92%)	12 (100%)	0	100	100
All	All	260/308 (84%)	255 (98%)	5 (2%)	58	67

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	THR
1	A	129	ARG
2	B	103	ASP
2	B	106	ARG
2	B	175	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	MH8	C	13	3	6,9,10	0.78	0	5,11,13	1.04	0
3	XCP	C	16	3	8,8,9	0.80	0	6,10,12	1.58	2 (33%)
3	XCP	C	2	3	8,8,9	0.88	1 (12%)	6,10,12	1.50	1 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AJE	C	6	3	12,15,16	0.90	1 (8%)	11,20,22	0.89	0
3	MH8	C	9	3	6,9,10	0.80	0	5,11,13	0.81	0
3	MH8	D	13	3	6,9,10	0.72	0	5,11,13	0.91	0
3	XCP	D	16	3	8,8,9	1.08	1 (12%)	6,10,12	1.52	2 (33%)
3	XCP	D	2	3	8,8,9	1.13	1 (12%)	6,10,12	1.46	1 (16%)
3	AJE	D	6	3	12,15,16	0.87	1 (8%)	11,20,22	1.02	1 (9%)
3	MH8	D	9	3	6,9,10	0.73	0	5,11,13	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MH8	C	13	3	-	0/7/9/12	0/0/0/0
3	XCP	C	16	3	-	0/1/12/14	0/1/1/1
3	XCP	C	2	3	-	0/1/12/14	0/1/1/1
3	AJE	C	6	3	-	0/8/23/25	0/1/1/1
3	MH8	C	9	3	-	0/7/9/12	0/0/0/0
3	MH8	D	13	3	-	0/7/9/12	0/0/0/0
3	XCP	D	16	3	-	0/1/12/14	0/1/1/1
3	XCP	D	2	3	-	0/1/12/14	0/1/1/1
3	AJE	D	6	3	-	0/8/23/25	0/1/1/1
3	MH8	D	9	3	-	0/7/9/12	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	XCP	CA-CB	-2.86	1.51	1.55
3	C	6	AJE	CA-CB	-2.69	1.51	1.55
3	D	16	XCP	CA-CB	-2.58	1.52	1.55
3	D	6	AJE	CA-CB	-2.42	1.52	1.55
3	C	2	XCP	CA-CB	-2.14	1.52	1.55

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	16	XCP	O-C-CA	-2.54	119.17	125.22
3	D	16	XCP	O-C-CA	-2.47	119.33	125.22
3	D	6	AJE	O-C-CA	-2.19	119.99	125.22
3	D	16	XCP	CE-CA-CB	2.21	105.25	100.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	16	XCP	CE-CA-CB	2.62	106.04	100.95
3	C	2	XCP	CE-CA-CB	2.67	106.14	100.95
3	D	2	XCP	CE-CA-CB	3.11	106.99	100.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	16	XCP	1	0
3	C	9	MH8	1	0
3	D	13	MH8	1	0
3	D	16	XCP	1	0
3	D	6	AJE	1	0
3	D	9	MH8	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	140/166 (84%)	-0.27	0 100 100	26, 42, 79, 112	0
2	B	141/166 (84%)	0.07	7 (4%) 29 29	28, 53, 90, 124	0
3	C	16/22 (72%)	-0.16	0 100 100	32, 36, 83, 90	0
3	D	16/22 (72%)	0.06	0 100 100	33, 42, 66, 75	0
All	All	313/376 (83%)	-0.09	7 (2%) 62 64	26, 46, 87, 124	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	32	ALA	4.6
2	B	31	ASP	3.3
2	B	12	ARG	2.6
2	B	16	MET	2.4
2	B	28	TYR	2.2
2	B	165	GLU	2.1
2	B	115	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	XCP	C	2	8/9	0.94	0.13	40,48,56,56	0
3	XCP	D	2	8/9	0.95	0.16	46,56,60,61	0
3	AJE	D	6	15/16	0.95	0.16	32,44,106,112	0
3	XCP	D	16	8/9	0.96	0.11	33,35,36,41	0
3	AJE	C	6	15/16	0.96	0.12	32,40,101,101	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MH8	D	9	10/11	0.97	0.13	22,38,40,48	0
3	MH8	C	9	10/11	0.97	0.13	23,30,39,49	0
3	XCP	C	16	8/9	0.97	0.12	25,28,36,45	0
3	MH8	C	13	10/11	0.97	0.16	28,34,42,51	0
3	MH8	D	13	10/11	0.98	0.11	29,34,54,54	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.