



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 08:47 am GMT

PDB ID : 4AIK
Title : Crystal structure of RovA from Yersinia in complex with an invasin promoter fragment
Authors : Quade, N.; Mendonca, C.; Herbst, K.; Heroven, A.K.; Heinz, D.W.; Dersch, P.
Deposited on : 2012-02-10
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2111 (1.86-1.86)
Clashscore	122126	2258 (1.86-1.86)
Ramachandran outliers	120053	2234 (1.86-1.86)
Sidechain outliers	120020	2234 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	151	66% 25% 8%
1	B	151	72% 22% 6%
2	C	21	43% 33% 24%
3	D	21	81% 19%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTIONAL REGULATOR SLYA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	139	1106	695	202	209	0	1	0
1	B	142	1136	713	206	217	0	2	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	144	LEU	-	expression tag	UNP B1JJ73
A	145	GLU	-	expression tag	UNP B1JJ73
A	146	HIS	-	expression tag	UNP B1JJ73
A	147	HIS	-	expression tag	UNP B1JJ73
A	148	HIS	-	expression tag	UNP B1JJ73
A	149	HIS	-	expression tag	UNP B1JJ73
A	150	HIS	-	expression tag	UNP B1JJ73
A	151	HIS	-	expression tag	UNP B1JJ73
A	81	SER	CYS	engineered mutation	UNP B1JJ73
A	108	SER	CYS	engineered mutation	UNP B1JJ73
B	144	LEU	-	expression tag	UNP B1JJ73
B	145	GLU	-	expression tag	UNP B1JJ73
B	146	HIS	-	expression tag	UNP B1JJ73
B	147	HIS	-	expression tag	UNP B1JJ73
B	148	HIS	-	expression tag	UNP B1JJ73
B	149	HIS	-	expression tag	UNP B1JJ73
B	150	HIS	-	expression tag	UNP B1JJ73
B	151	HIS	-	expression tag	UNP B1JJ73
B	81	SER	CYS	engineered mutation	UNP B1JJ73
B	108	SER	CYS	engineered mutation	UNP B1JJ73

- Molecule 2 is a DNA chain called ROVA PROMOTER FRAGMENT, 5'-D(*AP*TP*GP*A P*TP*AP*TP*TP *AP*TP**TP*TP*AP*TP*AP*TP*GP*AP*TP*AP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	21	Total	C	N	O	P	0	0	0
			430	210	75	125	20			

- Molecule 3 is a DNA chain called ROVA PROMOTER FRAGMENT, 5'-D(*TP*TP*TP*AP*TP*CP*AP*TP *AP*TP*AP*AP*AP*TP*AP*AP*TP*AP*TP*CP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	21	Total	C	N	O	P	0	0	0
			425	208	74	123	20			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	93	Total	O	0	0
			93	93		
4	B	66	Total	O	0	0
			66	66		
4	C	12	Total	O	0	0
			12	12		
4	D	13	Total	O	0	0
			13	13		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: TRANSCRIPTIONAL REGULATOR SLYA

Chain A: 



- Molecule 1: TRANSCRIPTIONAL REGULATOR SLYA

Chain B: 




- Molecule 2: ROVA PROMOTER FRAGMENT, 5'-D(*AP*TP*GP*AP*TP*AP*TP*TP *AP*TP**TP*TP*AP*TP*AP*TP*GP*AP*TP*AP*A)-3'

Chain C: 



- Molecule 3: ROVA PROMOTER FRAGMENT, 5'-D(*TP*TP*TP*AP*TP*CP*AP*TP *AP*TP*AP*AP*AP*TP*AP*AP*TP*AP*TP*CP*A)-3'

Chain D: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	88.68Å 88.68Å 67.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	76.81 – 1.85 26.67 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.0 (76.81-1.85) 98.5 (26.67-1.85)	Depositor EDS
R_{merge}	0.00	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.198 , 0.230 0.255 , 0.287	Depositor DCC
R_{free} test set	2510 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtrriage
Anisotropy	0.607	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 14.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtrriage
Estimated twinning fraction	0.008 for -h,-k,l 0.196 for h,-h-k,-l 0.017 for -k,-h,-l	Xtrriage
Reported twinning fraction	0.502 for H, K, L 0.498 for h,-h-k,-l	Depositor
Outliers	0 of 49952 reflections	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3281	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1121	0.64	0/1516
1	B	0.46	0/1154	0.60	0/1560
2	C	0.64	0/482	1.42	6/743 (0.8%)
3	D	0.66	0/476	1.27	1/732 (0.1%)
All	All	0.52	0/3233	0.92	7/4551 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	9	DA	P-O3'-C3'	8.23	129.58	119.70
3	D	11	DA	P-O3'-C3'	7.54	128.75	119.70
2	C	18	DA	P-O3'-C3'	7.36	128.53	119.70
2	C	3	DG	P-O3'-C3'	6.66	127.70	119.70
2	C	6	DA	P-O3'-C3'	5.60	126.42	119.70
2	C	5	DT	O4'-C1'-N1	5.54	111.88	108.00
2	C	20	DA	P-O3'-C3'	5.02	125.72	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1106	0	1172	31	0
1	B	1136	0	1202	33	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	430	0	243	9	0
3	D	425	0	242	2	0
4	A	93	0	0	3	0
4	B	66	0	0	4	0
4	C	12	0	0	0	0
4	D	13	0	0	0	0
All	All	3281	0	2859	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:ASP:O	1:A:135:LYS:HG2	1.66	0.95
1:B:65:ARG:HG2	1:B:65:ARG:HH11	1.43	0.84
1:A:140:LEU:O	1:A:141:GLN:HB3	1.76	0.84
1:A:102:GLN:HG2	4:A:2072:HOH:O	1.79	0.82
1:A:129:LEU:HD21	1:B:122:GLU:HB3	1.62	0.81
1:B:27:LEU:HB3	1:B:29:LEU:HD13	1.68	0.74
2:C:18:DA:H2''	2:C:19:DT:H5''	1.75	0.68
1:B:27:LEU:HD21	4:B:2052:HOH:O	1.93	0.68
2:C:16:DT:H2''	2:C:17:DG:C8	2.29	0.67
1:B:20[B]:ILE:HG22	1:B:110:THR:HG21	1.77	0.66
1:A:118:ILE:HD13	1:B:133:LEU:HD11	1.77	0.66
1:A:53:ALA:HB1	1:A:58:ILE:O	1.95	0.66
1:B:65:ARG:CG	1:B:65:ARG:HH11	2.08	0.65
1:A:15:VAL:HG12	1:B:137:ILE:CG2	2.29	0.63
1:A:31:GLN:O	1:A:35:VAL:HG23	2.03	0.59
1:A:60:GLN:HG3	1:A:61:PRO:HD3	1.84	0.58
1:A:102:GLN:NE2	4:A:2071:HOH:O	2.37	0.58
1:A:15:VAL:HG12	1:B:137:ILE:HG22	1.86	0.57
1:B:35:VAL:HG12	1:B:56:ILE:HG23	1.87	0.57
1:A:125:VAL:HG13	1:A:129:LEU:HD13	1.87	0.56
1:A:96:SER:HA	1:A:99:ILE:HD12	1.86	0.56
1:B:22:HIS:O	1:B:25:LYS:HB2	2.06	0.56
1:A:118:ILE:HG22	1:B:132:LYS:HD2	1.88	0.55
1:A:118:ILE:HG22	1:B:132:LYS:CD	2.38	0.54
2:C:20:DA:H61	3:D:3:DT:H3	1.55	0.53
1:A:111:ARG:NH2	4:A:2041:HOH:O	2.22	0.53
1:B:31:GLN:NE2	4:B:2010:HOH:O	2.42	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:9:DA:H1'	2:C:10:DT:C5'	2.40	0.52
2:C:9:DA:H1'	2:C:10:DT:H5'	1.92	0.52
1:B:92:LEU:HD22	1:B:100:ILE:HD11	1.91	0.50
1:B:20[B]:ILE:HD11	1:B:34:TRP:CZ3	2.47	0.49
1:A:9:LEU:HG	1:B:130:ILE:HD11	1.95	0.48
3:D:20:DC:H2''	3:D:21:DA:N7	2.29	0.48
1:A:122:GLU:HB3	1:B:129:LEU:HD21	1.96	0.47
1:A:15:VAL:HG12	1:B:137:ILE:HG21	1.96	0.47
1:A:69:GLN:O	1:A:72:GLU:HB2	2.14	0.47
1:A:119:SER:O	1:A:123[B]:ILE:HG23	2.14	0.47
1:A:140:LEU:O	1:A:141:GLN:CB	2.53	0.46
2:C:2:DT:H2''	2:C:3:DG:C8	2.50	0.46
1:B:17:ARG:HD2	1:B:31:GLN:OE1	2.15	0.46
1:B:94:GLU:CD	1:B:94:GLU:H	2.19	0.46
1:B:20[B]:ILE:HD13	1:B:107:ILE:CG1	2.46	0.45
1:A:63:LEU:O	1:A:67:LEU:HG	2.17	0.45
1:B:20[B]:ILE:HD13	1:B:107:ILE:HG12	1.98	0.44
1:A:69:GLN:O	1:A:73:LYS:HG2	2.17	0.44
1:B:140:LEU:HD23	1:B:143:LYS:HE3	2.00	0.44
2:C:9:DA:H2''	2:C:10:DT:OP2	2.17	0.44
1:B:47:GLN:HG2	1:B:51:GLN:HB2	1.99	0.44
1:A:42:ARG:NH1	1:B:4:THR:HG21	2.34	0.43
2:C:6:DA:H2'	2:C:7:DT:H71	2.00	0.43
1:B:27:LEU:CD2	4:B:2052:HOH:O	2.57	0.43
1:A:59:GLU:HB2	1:A:61:PRO:HD2	2.01	0.42
1:A:73:LYS:HB2	1:A:75:LEU:HG	2.02	0.42
1:A:130:ILE:HG21	1:B:8:ASP:HB3	2.01	0.42
1:B:52:LEU:HD23	1:B:63:LEU:HD21	2.02	0.41
1:A:126:LEU:HD11	1:B:9:LEU:HD21	2.02	0.41
1:B:65:ARG:NH1	1:B:65:ARG:CG	2.74	0.41
1:A:8:ASP:HB3	1:B:130:ILE:HG21	2.03	0.40
2:C:19:DT:H2''	2:C:20:DA:C8	2.56	0.40
1:B:102:GLN:NE2	4:B:2052:HOH:O	2.38	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/151 (91%)	136 (99%)	2 (1%)	0	100	100
1	B	142/151 (94%)	140 (99%)	2 (1%)	0	100	100
All	All	280/302 (93%)	276 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/137 (92%)	116 (92%)	10 (8%)	13	3
1	B	130/137 (95%)	127 (98%)	3 (2%)	53	37
All	All	256/274 (93%)	243 (95%)	13 (5%)	26	10

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	29	LEU
1	A	59	GLU
1	A	60	GLN
1	A	65	ARG
1	A	83	ASN
1	A	86	ARG
1	A	97	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	121	ASP
1	A	141	GLN
1	B	65	ARG
1	B	85	ARG
1	B	97	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	A	102	GLN
1	A	136	ASN
1	B	31	GLN
1	B	41	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.