



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 21, 2021 – 06:05 PM BST

PDB ID : 7AKP  
Title : Crystal structure of E. coli RNA helicase HrpA-D305A  
Authors : Grass, L.M.; Wollenhaupt, J.; Barthel, T.; Loll, B.; Wahl, M.C.  
Deposited on : 2020-10-01  
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.20  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.20


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	787	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent RNA helicase HrpA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	723	5800	3654	1063	1062	21	0	0	0


There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP A0A024L2B5
A	-2	ALA	-	expression tag	UNP A0A024L2B5
A	-1	MET	-	expression tag	UNP A0A024L2B5
A	0	ALA	-	expression tag	UNP A0A024L2B5
A	305	ALA	ASP	engineered mutation	UNP A0A024L2B5

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-dependent RNA helicase HrpA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.89Å 114.90Å 94.61Å 90.00° 101.94° 90.00°	Depositor
Resolution (Å)	48.81 – 2.59 48.81 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.81-2.59) 98.4 (48.81-2.59)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.96 (at 2.58Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122, PHENIX 1.19.1_4122	Depositor
R, $R_{free}$	0.211 , 0.239 0.217 , 0.238	Depositor DCC
$R_{free}$ test set	1285 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.0	Xtrriage
Anisotropy	0.544	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.23$	Xtrriage
Estimated twinning fraction	0.258 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5800	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/5902	0.53	0/7969

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5800	0	5907	106	0
All	All	5800	0	5907	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:LEU:HB3	1:A:522:THR:HG23	1.62	0.81
1:A:273:ASP:HB2	1:A:276:GLN:HB2	1.65	0.77
1:A:682:ARG:HG3	1:A:683:ILE:HG13	1.67	0.76
1:A:101:GLU:O	1:A:103:GLY:N	2.16	0.76
1:A:425:ASP:O	1:A:430:ARG:NH1	2.19	0.75

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:LYS:HE2	1:A:228:THR:HB	1.71	0.70
1:A:417:PHE:HA	1:A:420:ARG:HD2	1.76	0.67
1:A:278:ILE:HD11	1:A:295:ILE:HG21	1.78	0.66
1:A:440:MET:HA	1:A:445:LEU:HD12	1.79	0.65
1:A:275:LEU:HD13	1:A:306:THR:HG22	1.79	0.65
1:A:186:ARG:O	1:A:212:TYR:OH	2.14	0.64
1:A:307:ALA:O	1:A:311:ASN:ND2	2.29	0.64
1:A:338:SER:O	1:A:340:SER:N	2.31	0.64
1:A:501:ARG:HH22	1:A:632:SER:HB3	1.63	0.64
1:A:94:GLN:HA	1:A:224:LYS:HD3	1.81	0.62
1:A:605:VAL:HG13	1:A:610:ILE:HB	1.81	0.62
1:A:184:GLN:N	1:A:184:GLN:OE1	2.34	0.61
1:A:495:GLN:HG3	1:A:608:LEU:HD21	1.84	0.60
1:A:98:VAL:HG12	1:A:106:LYS:HD2	1.86	0.58
1:A:552:GLU:HA	1:A:660:LEU:HD21	1.86	0.57
1:A:64:ARG:NH1	1:A:120:GLY:O	2.38	0.57
1:A:351:ALA:O	1:A:399:ARG:NH2	2.36	0.57
1:A:493:LEU:HD21	1:A:500:PRO:HA	1.86	0.56
1:A:554:ASP:OD2	1:A:658:SER:OG	2.21	0.56
1:A:512:HIS:HB3	1:A:621:GLU:HB3	1.88	0.55
1:A:7:LEU:HD12	1:A:48:GLN:HA	1.90	0.54
1:A:213:LEU:O	1:A:217:LEU:HG	2.08	0.54
1:A:428:ILE:HG21	1:A:455:GLU:HB2	1.90	0.53
1:A:315:LEU:HB2	1:A:318:THR:HB	1.91	0.53
1:A:370:ARG:NH2	1:A:596:ASP:OD2	2.42	0.52
1:A:667:TRP:CD1	1:A:689:PRO:HG3	2.44	0.52
1:A:613:ASN:ND2	1:A:615:GLU:O	2.43	0.52
1:A:671:ALA:HB3	1:A:683:ILE:HB	1.92	0.52
1:A:467:ARG:HE	1:A:648:ARG:HH21	1.58	0.52
1:A:580:LEU:HA	1:A:583:THR:HG23	1.92	0.52
1:A:327:LEU:HB2	1:A:332:GLN:HG3	1.92	0.52
1:A:157:LYS:HB3	1:A:171:LYS:HZ3	1.76	0.51
1:A:182:ILE:HD13	1:A:188:LEU:HD21	1.92	0.51
1:A:95:VAL:HG12	1:A:225:ILE:HB	1.93	0.50
1:A:362:TYR:CD1	1:A:407:ILE:HB	2.47	0.50
1:A:501:ARG:NH1	1:A:629:GLY:O	2.45	0.50
1:A:241:PHE:O	1:A:243:ASN:N	2.46	0.49
1:A:195:ILE:HG12	1:A:226:ILE:HB	1.94	0.49
1:A:278:ILE:HD11	1:A:295:ILE:HD13	1.93	0.49
1:A:717:MET:HE3	1:A:734:LYS:HD3	1.94	0.49
1:A:171:LYS:HD3	1:A:191:TYR:HE1	1.77	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:289:SER:HB3	1:A:362:TYR:OH	2.12	0.49
1:A:366:PRO:HA	1:A:411:LEU:HB2	1.95	0.49
1:A:274:GLN:H	1:A:274:GLN:CD	2.16	0.48
1:A:333:ASN:HA	1:A:336:PHE:HD2	1.78	0.48
1:A:133:LEU:HD12	1:A:136:ARG:HD3	1.95	0.47
1:A:702:ARG:HH21	1:A:733:ARG:HH22	1.62	0.47
1:A:289:SER:OG	1:A:290:HIS:N	2.47	0.47
1:A:299:GLY:O	1:A:303:ILE:HG13	2.15	0.47
1:A:202:ARG:HH22	1:A:455:GLU:HG2	1.80	0.47
1:A:22:ARG:NH1	1:A:190:GLN:HA	2.31	0.46
1:A:723:THR:HA	1:A:727:LEU:O	2.14	0.46
1:A:688:ASP:HB2	1:A:691:TRP:CE2	2.50	0.46
1:A:519:MET:O	1:A:523:SER:OG	2.29	0.45
1:A:79:VAL:HG22	1:A:249:VAL:HG21	1.98	0.45
1:A:295:ILE:HG12	1:A:364:ILE:HB	1.99	0.45
1:A:195:ILE:HA	1:A:226:ILE:O	2.16	0.45
1:A:364:ILE:HG23	1:A:411:LEU:HD11	1.99	0.44
1:A:622:ILE:O	1:A:626:LEU:HG	2.17	0.44
1:A:530:PRO:HA	1:A:590:ARG:HH11	1.82	0.44
1:A:127:HIS:HB3	1:A:172:LEU:HD22	1.98	0.44
1:A:187:LEU:HA	1:A:216:LEU:HD11	2.00	0.44
1:A:298:SER:HB2	1:A:370:ARG:HD2	2.00	0.43
1:A:124:LEU:HB2	1:A:191:TYR:CD1	2.53	0.43
1:A:502:LEU:HA	1:A:505:MET:HE3	2.00	0.43
1:A:334:ARG:HA	1:A:337:GLN:HG2	2.00	0.43
1:A:400:CYS:HB3	1:A:406:GLY:HA3	1.99	0.43
1:A:489:LEU:HD21	1:A:504:ARG:NE	2.33	0.43
1:A:543:GLU:OE1	1:A:546:ARG:NH2	2.51	0.43
1:A:294:LEU:HB2	1:A:360:ILE:HG12	1.99	0.43
1:A:324:TYR:HB2	1:A:327:LEU:HG	2.00	0.43
1:A:708:HIS:O	1:A:716:VAL:HA	2.18	0.43
1:A:112:LYS:HE3	1:A:146:LEU:HG	2.01	0.43
1:A:257:GLU:HB3	1:A:407:ILE:HG12	2.01	0.43
1:A:698:HIS:CD2	1:A:699:LEU:HG	2.54	0.42
1:A:153:CYS:O	1:A:169:MET:N	2.51	0.42
1:A:78:PRO:HB3	1:A:252:ARG:HD2	2.00	0.42
1:A:491:ARG:O	1:A:495:GLN:HG2	2.18	0.42
1:A:95:VAL:HG11	1:A:241:PHE:CD2	2.54	0.42
1:A:475:ILE:HA	1:A:486:LEU:HA	2.01	0.42
1:A:19:LEU:CD2	1:A:23:ASP:HB2	2.50	0.42
1:A:199:ALA:HA	1:A:206:ILE:HG21	2.01	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:623:HIS:CD2	1:A:626:LEU:HD12	2.54	0.42
1:A:275:LEU:O	1:A:278:ILE:HG22	2.19	0.42
1:A:297:MET:HE1	1:A:306:THR:HG21	2.01	0.42
1:A:323:LEU:HD12	1:A:327:LEU:HD12	2.02	0.42
1:A:487:THR:HG22	1:A:489:LEU:H	1.85	0.41
1:A:259:ARG:HD3	1:A:261:ARG:NH1	2.36	0.41
1:A:478:ASP:OD2	1:A:485:LYS:HD2	2.21	0.41
1:A:95:VAL:O	1:A:245:PRO:HD2	2.20	0.41
1:A:240:HIS:CD2	1:A:454:VAL:HG22	2.55	0.41
1:A:365:ASP:OD1	1:A:396:ARG:NH1	2.52	0.41
1:A:217:LEU:HD11	1:A:241:PHE:CD2	2.56	0.41
1:A:19:LEU:HD21	1:A:23:ASP:HB2	2.02	0.41
1:A:320:ILE:HG23	1:A:346:LEU:HD13	2.03	0.40
1:A:499:ASP:O	1:A:503:ALA:N	2.36	0.40
1:A:310:LEU:HB2	1:A:320:ILE:HG12	2.03	0.40
1:A:68:ARG:NH1	1:A:118:GLY:HA2	2.36	0.40
1:A:154:ILE:HG13	1:A:170:VAL:HB	2.03	0.40
1:A:197:ASP:OD1	1:A:198:GLU:N	2.54	0.40
1:A:670:VAL:HG12	1:A:684:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	713/787 (91%)	686 (96%)	22 (3%)	5 (1%)	22 43

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	102	THR
1	A	339	HIS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	314	ASN
1	A	683	ILE
1	A	232	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	626/680 (92%)	605 (97%)	21 (3%)	37 63

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	ASP
1	A	19	LEU
1	A	23	ASP
1	A	28	SER
1	A	38	LYS
1	A	145	GLU
1	A	161	SER
1	A	233	ASP
1	A	247	ILE
1	A	274	GLN
1	A	278	ILE
1	A	290	HIS
1	A	360	ILE
1	A	463	GLN
1	A	493	LEU
1	A	523	SER
1	A	583	THR
1	A	642	GLN
1	A	702	ARG
1	A	704	TYR
1	A	746	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are

no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	723/787 (91%)	0.48	48 (6%) <b>18</b> <b>13</b>	39, 65, 112, 163	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	LEU	11.0
1	A	102	THR	5.1
1	A	660	LEU	4.8
1	A	291	GLY	4.5
1	A	160	PHE	4.3
1	A	402	ARG	3.8
1	A	360	ILE	3.7
1	A	362	TYR	3.6
1	A	403	VAL	3.5
1	A	25	LEU	3.4
1	A	47	PHE	3.1
1	A	46	ILE	3.1
1	A	27	PHE	3.0
1	A	24	ARG	2.9
1	A	232	ILE	2.8
1	A	642	GLN	2.7
1	A	673	LEU	2.7
1	A	59	GLY	2.6
1	A	716	VAL	2.5
1	A	33	GLY	2.5
1	A	45	ALA	2.5
1	A	654	ILE	2.4
1	A	745	CYS	2.4
1	A	749	PHE	2.4
1	A	744	LEU	2.4
1	A	40	PRO	2.3
1	A	549	HIS	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	172	LEU	2.3
1	A	112	LYS	2.2
1	A	252	ARG	2.2
1	A	361	LYS	2.2
1	A	61	VAL	2.2
1	A	718	ALA	2.2
1	A	135	ALA	2.2
1	A	293	ILE	2.2
1	A	408	CYS	2.2
1	A	43	GLN	2.2
1	A	62	LEU	2.2
1	A	601	LEU	2.2
1	A	577	PHE	2.2
1	A	12	LEU	2.1
1	A	383	LEU	2.1
1	A	377	ARG	2.1
1	A	32	HIS	2.1
1	A	407	ILE	2.1
1	A	22	ARG	2.1
1	A	13	GLN	2.1
1	A	9	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.