

# Full wwPDB X-ray Structure Validation Report (i)

#### Aug 7, 2023 – 06:11 pm BST

PDB ID : 8AKE

Title: Human Sirt6 in complex with ADP-ribose and fragment nicotinamide

Authors: You, W.; Steegborn, C.

Deposited on : 2022-07-29

Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.34

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.34

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	302	80%	12%	• 7%			
1	В	302	80%	11%	• 8%			



## 2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 4586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-6.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	A 282	Total	С	N	О	S	0	0	0
	202	2202	1386	404	401	11	0	U		
1	D	277	Total	С	N	О	S	0	0	0
1	1 B	211	2158	1358	396	393	11	0	U	

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	GLY	-	expression tag	UNP Q8N6T7
A	8	ILE	-	expression tag	UNP Q8N6T7
A	9	ASP	-	expression tag	UNP Q8N6T7
A	10	PRO	-	expression tag	UNP Q8N6T7
A	11	PHE	-	expression tag	UNP Q8N6T7
A	12	THR	-	expression tag	UNP Q8N6T7
В	7	GLY	_	expression tag	UNP Q8N6T7
В	8	ILE	-	expression tag	UNP Q8N6T7
В	9	ASP	_	expression tag	UNP Q8N6T7
В	10	PRO	-	expression tag	UNP Q8N6T7
В	11	PHE	-	expression tag	UNP Q8N6T7
В	12	THR	-	expression tag	UNP Q8N6T7

• Molecule 2 is  $[(2R,3S,4R,5R)-5-(6-AMINOPURIN-9-YL)-3,4-DIHYDROXY-OXOLAN-2-YL]METHYL [HYDROXY-[[(2R,3S,4R,5S)-3,4,5-TRIHYDROXYOXOLAN-2-YL]ME THOXY]PHOSPHORYL] HYDROGEN PHOSPHATE (three-letter code: AR6) (formula: <math>C_{15}H_{23}N_5O_{14}P_2$ ).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Ω Λ	Λ 1	Total	С	N	О	Р	0	0
$\begin{array}{ c c c c c }\hline Z & A & A \\\hline \end{array}$	1	36	15	5	14	2	U	0	
9	) D	1	Total	С	N	О	Р	0	0
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	1	36	15	5	14	2	U	U	

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	В	1	Total Zn 1 1	0	0


• Molecule 4 is NICOTINAMIDE (three-letter code: NCA) (formula:  $C_6H_6N_2O$ ) (labeled as "Ligand of Interest" by depositor).





M	[ol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	4	A	1	Total 9		N 2	O 1	0	0
4	4	В	1	Total 9		N 2	O 1	0	0


 $\bullet$  Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $\mathrm{C_2H_6O_2}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0


 $\bullet \ \ Molecule \ 6 \ is \ DI(HYDROXYETHYL)ETHER \ (three-letter \ code: \ PEG) \ (formula: \ C_4H_{10}O_3).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 7 4 3	0	0
6	В	1	Total C O 7 4 3	0	0

 $\bullet$  Molecule 7 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O S 5 4 1	0	0
7	A	1	Total O S 5 4 1	0	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O S 5 4 1	0	0
7	В	1	Total O S 5 4 1	0	0

#### • Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	53	Total O 53 53	0	0
8	В	43	Total O 43 43	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NAD-dependent protein deacetylase sirtuin-6





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants	91.11Å 91.11Å 143.43Å	Donositon
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	45.55 - 1.82	Depositor
rtesolution (A)	45.55 - 1.82	EDS
% Data completeness	99.7 (45.55-1.82)	Depositor
(in resolution range)	99.7 (45.55-1.82)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.03 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
D D.	0.177 , 0.224	Depositor
$R, R_{free}$	0.182 , $0.223$	DCC
$R_{free}$ test set	2101 reflections $(3.49\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.5	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37 , 21.9	EDS
L-test for twinning <sup>2</sup>	$< L >=0.39, < L^2>=0.22$	Xtriage
Estimated twinning fraction	0.466 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.516 for H, K, L	Depositor
Reported twinning fraction	0.484  for -K, -H, -L	Depositor
Outliers	0 of 60246 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4586	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: AR6, EDO, SO4, NCA, PEG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond lengths		Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.77	0/2251	1.03	$13/3053 \ (0.4\%)$	
1	В	0.74	0/2205	0.98	$12/2991 \ (0.4\%)$	
All	All	0.76	0/4456	1.00	25/6044~(0.4%)	

There are no bond length outliers.

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
1	A	253	ARG	NE-CZ-NH1	-7.22	116.69	120.30
1	A	220	ARG	NE-CZ-NH2	7.20	123.90	120.30
1	A	39	ARG	NE-CZ-NH2	7.08	123.84	120.30
1	A	39	ARG	NE-CZ-NH1	-6.66	116.97	120.30
1	A	264	ARG	NE-CZ-NH1	6.52	123.56	120.30
1	В	232	ARG	NE-CZ-NH2	-6.45	117.07	120.30
1	В	253	ARG	NE-CZ-NH2	-6.40	117.10	120.30
1	A	264	ARG	NE-CZ-NH2	-6.35	117.13	120.30
1	A	76	ARG	NE-CZ-NH2	-6.23	117.19	120.30
1	В	280	ARG	NE-CZ-NH2	6.21	123.41	120.30
1	A	76	ARG	NE-CZ-NH1	6.18	123.39	120.30
1	В	253	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	A	220	ARG	NE-CZ-NH1	-6.14	117.23	120.30
1	A	253	ARG	NE-CZ-NH2	5.99	123.30	120.30
1	A	103	ARG	NE-CZ-NH2	5.92	123.26	120.30
1	В	232	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	В	280	ARG	NE-CZ-NH1	-5.71	117.45	120.30
1	В	39	ARG	NE-CZ-NH2	-5.61	117.50	120.30
1	В	76	ARG	NE-CZ-NH2	5.53	123.07	120.30
1	A	232	ARG	NE-CZ-NH1	-5.47	117.56	120.30
1	В	76	ARG	NE-CZ-NH1	-5.44	117.58	120.30
1	В	39	ARG	NE-CZ-NH1	5.42	123.01	120.30



Continued from previous page...

Mol	Chain	Res	Type	Atoms	${f Z}$	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	В	103	ARG	NE-CZ-NH1	5.30	122.95	120.30
1	A	280	ARG	NE-CZ-NH1	5.29	122.94	120.30
1	В	220	ARG	NE-CZ-NH1	5.11	122.85	120.30

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2202	0	2215	22	1
1	В	2158	0	2171	26	0
2	A	36	0	21	1	0
2	В	36	0	21	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	A	9	0	6	0	0
4	В	9	0	6	0	0
5	A	4	0	6	0	0
6	A	7	0	10	1	0
6	В	7	0	10	0	0
7	A	15	0	0	0	0
7	В	5	0	0	1	0
8	A	53	0	0	3	0
8	В	43	0	0	7	0
All	All	4586	0	4466	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)	
1:B:286:LEU:HD23	8:B:542:HOH:O	1.74	0.87	
1:B:286:LEU:HA	8:B:542:HOH:O	1.86	0.74	



 $Continued\ from\ previous\ page...$ 

Continued from prei		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:154:VAL:CG1	1:A:162:THR:HG21	2.28	0.63
1:A:59:SER:OG	1:A:90:ARG:O	2.14	0.62
1:B:50:HIS:HA	1:B:111:VAL:O	2.01	0.61
1:A:102:GLU:OE2	1:A:128:LYS:NZ	2.27	0.60
1:B:154:VAL:CG1	1:B:162:THR:HG21	2.33	0.58
1:A:39:ARG:HD3	8:A:516:HOH:O	2.03	0.58
1:B:104:VAL:HB	1:B:291:ARG:HB3	1.90	0.53
1:B:57:THR:HA	1:B:61:ILE:O	2.09	0.51
1:A:104:VAL:HB	1:A:291:ARG:HB3	1.93	0.51
1:B:293:PRO:O	8:B:501:HOH:O	2.19	0.51
1:B:253:ARG:NH1	7:B:405:SO4:S	2.84	0.51
1:B:111:VAL:CG1	1:B:132:LEU:HD11	2.41	0.50
1:A:154:VAL:CG1	1:A:162:THR:CG2	2.92	0.48
1:B:59:SER:C	8:B:509:HOH:O	2.52	0.48
1:B:110:LEU:HD21	1:B:119:HIS:CE1	2.49	0.48
1:A:50:HIS:HA	1:A:111:VAL:O	2.14	0.47
1:A:53:ALA:HB1	1:A:63:ASP:HA	1.96	0.47
1:B:25:ASP:OD1	1:B:255:HIS:HB2	2.15	0.46
1:A:111:VAL:CG1	1:A:132:LEU:HD11	2.44	0.46
1:A:72:THR:O	1:A:76:ARG:HG2	2.15	0.46
1:A:87:GLU:CD	8:A:524:HOH:O	2.53	0.46
1:A:154:VAL:HG11	1:A:162:THR:HG21	1.97	0.46
1:B:61:ILE:HG13	8:B:509:HOH:O	2.16	0.45
1:A:247:ASP:O	1:A:253:ARG:HD3	2.16	0.45
1:A:161:ALA:HB2	1:A:180:GLU:OE2	2.16	0.45
1:A:63:ASP:OD2	2:A:401:AR6:H8	2.17	0.44
1:A:207:ALA:O	1:A:229:THR:HG23	2.17	0.44
1:B:59:SER:OG	1:B:90:ARG:O	2.25	0.44
6:A:405:PEG:H21	1:B:82:PHE:CD2	2.53	0.44
1:A:280:ARG:N	8:A:506:HOH:O	2.51	0.43
1:B:59:SER:CB	8:B:509:HOH:O	2.66	0.43
1:A:57:THR:HA	1:A:61:ILE:O	2.19	0.43
1:A:90:ARG:HA	1:A:91:PRO:HD3	1.92	0.43
1:B:154:VAL:HG11	1:B:162:THR:HG21	1.99	0.42
1:A:26:PRO:HD2	1:A:29:GLU:OE2	2.20	0.42
1:A:23:ILE:O	1:A:255:HIS:HB3	2.20	0.41
1:A:84:THR:HG21	1:A:89:ALA:HB2	2.01	0.41
1:B:76:ARG:HB2	1:B:78:LEU:HD12	2.03	0.41
1:B:104:VAL:HG21	1:B:271:LEU:HD11	2.01	0.41
1:B:110:LEU:HD21	1:B:119:HIS:ND1	2.35	0.41
1:B:35:TRP:CE2	1:B:268:HIS:CE1	3.08	0.41



I 'omtamalod	tmom	mmonia	maaa
Continued	110111	DIEUIUU	DUJUE
00,000,000	.,	p	p = 9 =

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:23:ILE:O	1:B:255:HIS:HB3	2.21	0.41
1:B:235:ARG:HD2	1:B:251:ASP:OD2	2.19	0.41
1:B:262:MET:O	1:B:266:MET:HG2	2.21	0.41
1:B:154:VAL:CG1	1:B:162:THR:CG2	2.99	0.41
1:B:253:ARG:NH2	8:B:506:HOH:O	2.53	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{aligned}$	Clash overlap (Å)
1:A:11:PHE:O	1:A:17:LYS:NZ[2_565]	1.94	0.26

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	278/302 (92%)	272 (98%)	6 (2%)	0	100	100
1	В	273/302 (90%)	267 (98%)	6 (2%)	0	100	100
All	All	551/604 (91%)	539 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentile	
1	A	239/256~(93%)	236 (99%)	3 (1%)	69	61
1	В	234/256 (91%)	233 (100%)	1 (0%)	91	89
All	All	473/512 (92%)	469 (99%)	4 (1%)	81	77

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	THR
1	A	103	ARG
1	A	244	THR
1	В	222	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	224	ASN
1	A	268	HIS
1	В	249	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res Link Bond lengths			Bond angles				
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	PEG	В	401	-	6,6,6	0.13	0	5,5,5	0.08	0
2	AR6	A	401	-	34,39,39	0.73	0	40,60,60	0.94	3 (7%)
7	SO4	A	407	-	4,4,4	0.39	0	6,6,6	0.15	0
7	SO4	В	405	-	4,4,4	0.41	0	6,6,6	0.07	0
4	NCA	В	404	-	9,9,9	0.21	0	11,11,11	0.39	0
7	SO4	A	406	-	4,4,4	0.54	0	6,6,6	0.12	0
5	EDO	A	404	_	3,3,3	0.21	0	2,2,2	0.38	0
4	NCA	A	403	-	9,9,9	0.20	0	11,11,11	0.63	0
2	AR6	В	402	_	34,39,39	0.62	0	40,60,60	0.80	0
7	SO4	A	408	-	4,4,4	0.37	0	6,6,6	0.11	0
6	PEG	A	405	_	6,6,6	0.26	0	5,5,5	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	В	401	-	-	3/4/4/4	-
2	AR6	A	401	-	-	1/18/54/54	0/4/4/4
4	NCA	В	404	-	-	0/4/4/4	0/1/1/1
5	EDO	A	404	-	-	1/1/1/1	-
4	NCA	A	403	-	-	0/4/4/4	0/1/1/1
2	AR6	В	402	-	-	3/18/54/54	0/4/4/4
6	PEG	A	405	-	-	3/4/4/4	_

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	401	AR6	O1A-PA-O2A	2.33	123.77	112.24
2	A	401	AR6	C2D-C3D-C4D	-2.27	98.24	102.64
2	A	401	AR6	C5-C6-N6	2.11	123.56	120.35

There are no chirality outliers.

All (11) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	404	EDO	O1-C1-C2-O2
6	A	405	PEG	O2-C3-C4-O4
2	В	402	AR6	O4'-C4'-C5'-O5'
6	В	401	PEG	O1-C1-C2-O2
6	A	405	PEG	C4-C3-O2-C2
6	В	401	PEG	C4-C3-O2-C2
6	A	405	PEG	O1-C1-C2-O2
6	В	401	PEG	O2-C3-C4-O4
2	A	401	AR6	O4'-C4'-C5'-O5'
2	В	402	AR6	C5D-O5D-PB-O2B
2	В	402	AR6	C3'-C4'-C5'-O5'


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	AR6	1	0
7	В	405	SO4	1	0
6	A	405	PEG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$282/302 \ (93\%)$	-0.22	1 (0%) 92 91	21, 33, 51, 65	0
1	В	277/302 (91%)	-0.10	10 (3%) 42 37	24, 37, 58, 74	0
All	All	559/604 (92%)	-0.16	11 (1%) 65 61	21, 35, 56, 74	0

All (11) RSRZ outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	RSRZ
1	A	169	ALA	6.0
1	В	169	ALA	4.3
1	В	168	VAL	3.0
1	В	188	TRP	3.0
1	В	178	ARG	2.8
1	В	233	GLY	2.7
1	В	232	ARG	2.6
1	В	277	ASP	2.5
1	В	76	ARG	2.3
1	В	297	LEU	2.2
1	В	15	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
6	PEG	A	405	7/7	0.90	0.11	39,40,46,52	0
4	NCA	В	404	9/9	0.94	0.09	34,37,39,43	0
4	NCA	A	403	9/9	0.94	0.10	26,27,32,32	0
6	PEG	В	401	7/7	0.94	0.08	34,39,40,43	0
5	EDO	A	404	4/4	0.95	0.06	44,44,44,45	0
3	ZN	В	403	1/1	0.98	0.04	62,62,62,62	0
2	AR6	A	401	36/36	0.98	0.08	16,28,32,35	0
2	AR6	В	402	36/36	0.98	0.07	25,34,41,48	0
7	SO4	A	408	5/5	0.98	0.10	45,46,50,50	0
7	SO4	A	406	5/5	0.99	0.07	36,39,42,43	0
7	SO4	A	407	5/5	0.99	0.12	45,45,45,50	0
3	ZN	A	402	1/1	0.99	0.05	48,48,48,48	0
7	SO4	В	405	5/5	0.99	0.11	59,62,65,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














#### Electron density around AR6 A 401:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around AR6 B 402:

 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









## 6.5 Other polymers (i)

There are no such residues in this entry.

