



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 26, 2023 – 04:30 AM EDT

PDB ID : 1ALQ
Title : CIRCULARLY PERMUTED BETA-LACTAMASE FROM STAPHYLOCOCCUS AUREUS PC1
Authors : Pieper, U.; Herzberg, O.
Deposited on : 1997-06-02
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	266	 72% 20% 5% •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2254 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CP254 BETA-LACTAMASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	259	2033	1296	341	393	3	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 1 3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	212	Total O 212 212	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: CP254 BETA-LACTAMASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	53.90Å 94.20Å 138.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80	Depositor
% Data completeness (in resolution range)	88.8 (30.00-1.80)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	SHELXL-97, X-PLOR	Depositor
R, R_{free}	0.196 , 0.256	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2254	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/2062	1.77	36/2769 (1.3%)

There are no bond length outliers.

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	244	ARG	NE-CZ-NH1	12.96	126.78	120.30
1	A	151	ARG	NE-CZ-NH2	-12.44	114.08	120.30
1	A	227	LYS	C-N-CA	10.94	149.06	121.70
1	A	244	ARG	NE-CZ-NH2	-10.78	114.91	120.30
1	A	226	PRO	C-N-CA	10.68	148.39	121.70
1	A	151	ARG	NE-CZ-NH1	10.59	125.59	120.30
1	A	141	GLU	OE1-CD-OE2	9.78	135.03	123.30
1	A	171	TYR	CB-CG-CD2	-9.49	115.31	121.00
1	A	65	ARG	NE-CZ-NH2	8.86	124.73	120.30
1	A	272	ASP	CB-CG-OD2	8.68	126.11	118.30
1	A	65	ARG	CD-NE-CZ	8.59	135.63	123.60
1	A	101	ASP	CB-CG-OD1	-8.39	110.75	118.30
1	A	202	GLU	OE1-CD-OE2	-8.36	113.27	123.30
1	A	164	ARG	NE-CZ-NH2	7.59	124.09	120.30
1	A	129	TYR	CB-CG-CD1	-7.41	116.55	121.00
1	A	145	ILE	C-N-CA	7.30	139.96	121.70
1	A	229	TYR	CG-CD2-CE2	-6.84	115.83	121.30
1	A	105	TYR	CB-CG-CD2	6.79	125.08	121.00
1	A	227	LYS	N-CA-CB	-6.43	99.03	110.60
1	A	229	TYR	CZ-CE2-CD2	6.36	125.53	119.80
1	A	82	GLU	OE1-CD-OE2	6.26	130.82	123.30
1	A	297	ALA	C-N-CA	6.26	137.34	121.70
1	A	64	LYS	CA-CB-CG	5.90	126.38	113.40
1	A	34	ASN	OD1-CG-ND2	5.87	135.40	121.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	34	ASN	CA-CB-CG	-5.87	100.50	113.40
1	A	68	TYR	CB-CG-CD1	5.85	124.51	121.00
1	A	34	ASN	CB-CG-OD1	-5.53	110.54	121.60
1	A	111	LYS	CB-CG-CD	5.51	125.93	111.60
1	A	65	ARG	NH1-CZ-NH2	-5.50	113.35	119.40
1	A	241	TYR	CB-CG-CD1	-5.45	117.73	121.00
1	A	141	GLU	CG-CD-OE2	-5.40	107.50	118.30
1	A	40	TYR	CB-CG-CD1	5.37	124.22	121.00
1	A	32	GLU	OE1-CD-OE2	5.31	129.67	123.30
1	A	129	TYR	CB-CG-CD2	5.28	124.17	121.00
1	A	218	ASP	O-C-N	-5.06	114.61	122.70
1	A	34	ASN	CB-CA-C	-5.02	100.36	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2033	0	2111	31	0
2	A	5	0	0	0	0
3	A	4	0	0	0	0
4	A	212	0	0	11	0
All	All	2254	0	2111	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:LYS:HD3	1:A:218:ASP:OD2	1.93	0.68
1:A:291:ALA:HB2	4:A:464:HOH:O	1.93	0.68
1:A:215:LYS:HD2	4:A:415:HOH:O	2.00	0.59
1:A:38:LYS:HE2	4:A:443:HOH:O	2.04	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:GLN:O	1:A:154:GLU:HG3	2.07	0.55
1:A:267:LYS:HE3	1:A:41:ASN:O	2.08	0.54
1:A:151:ARG:O	1:A:155:LEU:HG	2.08	0.54
1:A:258:PRO:HB3	1:A:251:TYR:CE2	2.43	0.53
1:A:120:LYS:O	1:A:124:GLU:HG3	2.09	0.53
1:A:222:LYS:HG3	4:A:310:HOH:O	2.08	0.53
1:A:174:PRO:HD3	1:A:241:TYR:CE2	2.43	0.53
1:A:147:LYS:HE2	4:A:445:HOH:O	2.09	0.52
1:A:205:LYS:HE2	4:A:493:HOH:O	2.10	0.52
1:A:229:TYR:CG	1:A:252:PRO:HA	2.45	0.52
1:A:288:LYS:HD2	1:A:298:ALA:HB2	1.98	0.45
1:A:34:ASN:ND2	1:A:34:ASN:H	2.14	0.45
1:A:195:ALA:O	1:A:204:LYS:NZ	2.49	0.45
1:A:228:ASP:OD1	1:A:228:ASP:N	2.49	0.45
1:A:269:ASN:HB2	1:A:272:ASP:CG	2.37	0.45
1:A:253:LYS:HD3	4:A:499:HOH:O	2.17	0.45
1:A:147:LYS:NZ	4:A:371:HOH:O	2.50	0.44
1:A:229:TYR:CD2	1:A:252:PRO:HA	2.53	0.43
1:A:198:LYS:HB2	4:A:440:HOH:O	2.19	0.43
1:A:257:GLU:HB3	1:A:51:THR:OG1	2.19	0.43
1:A:250:VAL:O	1:A:252:PRO:HD3	2.18	0.43
1:A:229:TYR:CD1	1:A:252:PRO:HA	2.54	0.42
1:A:277:LYS:HD2	4:A:419:HOH:O	2.20	0.42
1:A:150:GLN:O	1:A:150:GLN:HG2	2.17	0.42
1:A:52:LYS:HD3	4:A:429:HOH:O	2.20	0.41
1:A:269:ASN:HB2	1:A:272:ASP:OD1	2.20	0.41
1:A:48:ALA:O	1:A:56:GLU:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	255/266 (96%)	249 (98%)	5 (2%)	1 (0%)	34 21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	225/231 (97%)	208 (92%)	17 (8%)	13 4

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	257	GLU
1	A	277	LYS
1	A	31	LYS
1	A	35	ASP
1	A	38	LYS
1	A	83	GLN
1	A	111	LYS
1	A	137	LYS
1	A	145	ILE
1	A	146	LYS
1	A	149	LYS
1	A	153	LYS
1	A	201	LYS
1	A	212	LEU
1	A	215	LYS
1	A	216	SER
1	A	228	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	92	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CO3	A	302	-	2,3,3	0.85	0	2,3,3	0.69	0
2	SO4	A	301	-	4,4,4	0.53	0	6,6,6	0.22	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.