



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 23, 2020 – 02:09 pm GMT

PDB ID : 7APF
Title : Crystal structure of JAK3 in complex with FM601 (compound 10a)
Authors : Chaikuad, A.; Forster, M.; Gehringer, M.; Laufer, S.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2020-10-16
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	 7% 93% 5% ..
1	B	294	 9% 90% 6% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5072 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase JAK3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	290	2335	1483	415	422	15	0	3	0
1	B	285	2293	1460	403	415	15	0	3	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	810	SER	-	expression tag	UNP P52333
A	811	MET	-	expression tag	UNP P52333
A	949	ALA	ASP	conflict	UNP P52333
A	1040	SER	CYS	conflict	UNP P52333
A	1048	SER	CYS	conflict	UNP P52333
B	810	SER	-	expression tag	UNP P52333
B	811	MET	-	expression tag	UNP P52333
B	949	ALA	ASP	conflict	UNP P52333
B	1040	SER	CYS	conflict	UNP P52333
B	1048	SER	CYS	conflict	UNP P52333

- Molecule 2 is 3-[3-(propanoylamino)phenyl]-1 {H}-pyrrolo[2,3-b]pyridine-5-carboxamide (three-letter code: RQZ) (formula: C₁₇H₁₆N₄O₂) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	46	34	8	4	0	1
2	B	1	46	34	8	4	0	1

- Molecule 3 is 1-phenylurea (three-letter code: PHU) (formula: $C_7H_8N_2O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	10	7	2	1	0	0
3	B	1	10	7	2	1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	169	Total O 169 169	0	0
5	B	131	Total O 131 131	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK3



- Molecule 1: Tyrosine-protein kinase JAK3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.67Å 62.50Å 67.87Å 90.00° 101.32° 90.00°	Depositor
Resolution (Å)	45.56 – 1.95 45.56 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.3 (45.56-1.95) 97.3 (45.56-1.95)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.203 , 0.255 0.212 , 0.257	Depositor DCC
R_{free} test set	1868 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5072	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5204e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RQZ, EDO, PHU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/2398	0.70	0/3240
1	B	0.67	0/2354	0.69	0/3181
All	All	0.69	0/4752	0.70	0/6421

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	3
All	All	0	8

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1036	ARG	Sidechain
1	A	1085	ARG	Sidechain
1	A	820	ARG	Sidechain
1	A	870	ARG	Sidechain
1	A	918	ARG	Sidechain
1	B	1036	ARG	Sidechain
1	B	1085	ARG	Sidechain
1	B	953	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2335	0	2319	12	0
1	B	2293	0	2270	14	0
2	A	46	0	0	3	0
2	B	46	0	0	1	0
3	A	10	0	8	0	0
3	B	10	0	8	0	0
4	A	20	0	30	0	0
4	B	12	0	18	0	0
5	A	169	0	0	8	0
5	B	131	0	0	4	0
All	All	5072	0	4653	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1201[B]:RQZ:O	5:B:1301:HOH:O	2.02	0.77
2:A:1201[B]:RQZ:O	5:A:1301:HOH:O	2.06	0.72
1:A:870:ARG:HD2	5:A:1400:HOH:O	1.91	0.69
2:A:1201[A]:RQZ:O	5:A:1302:HOH:O	2.12	0.66
1:B:1083[B]:GLN:NE2	1:B:1083[B]:GLN:HA	2.11	0.65
1:A:1048:SER:OG	5:A:1303:HOH:O	2.14	0.64
1:B:891:TYR:CE1	1:B:898:LEU:HD13	2.40	0.57
1:B:948:ARG:HD3	5:B:1302:HOH:O	2.08	0.54
1:A:918:ARG:NH2	5:A:1316:HOH:O	2.43	0.51
1:B:1083[B]:GLN:HE21	1:B:1083[B]:GLN:HA	1.76	0.50
1:B:953:ARG:NH1	5:B:1312:HOH:O	2.46	0.48
1:B:891:TYR:HE1	1:B:898:LEU:HD13	1.77	0.48
1:B:910:LEU:HD21	1:B:1020:LEU:HD21	1.95	0.48
1:B:906:PRO:HD2	5:B:1315:HOH:O	2.15	0.47
1:A:827:GLN:NE2	1:A:830:LYS:HB2	2.30	0.46
1:B:1081:SER:OG	1:B:1084:ASP:OD2	2.34	0.46
1:A:862:PRO:HA	1:A:865:GLN:OE1	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:910:LEU:HD21	1:A:1020:LEU:HD21	1.99	0.45
1:A:982:VAL:HB	1:B:851:LEU:CD1	2.46	0.45
1:B:972:LYS:HG3	1:B:981:TYR:CZ	2.52	0.44
1:A:918:ARG:HD2	5:A:1316:HOH:O	2.18	0.43
1:A:953:ARG:NH1	5:A:1327:HOH:O	2.51	0.42
1:A:918:ARG:HG3	1:A:1023:TYR:CG	2.54	0.42
1:B:924:SER:OG	1:B:1102:SER:HB3	2.19	0.41
2:A:1201[A]:RQZ:C12	2:A:1201[A]:RQZ:O	2.68	0.41
1:B:862:PRO:HA	1:B:865:GLN:OE1	2.21	0.41
1:B:1033:GLU:CD	1:B:1036:ARG:HH11	2.24	0.41
1:A:881:ASP:HA	5:A:1331:HOH:O	2.21	0.40
1:A:926:LEU:HD11	1:A:1023:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/294 (99%)	284 (98%)	7 (2%)	0	100	100
1	B	284/294 (97%)	280 (99%)	3 (1%)	1 (0%)	34	22
All	All	575/588 (98%)	564 (98%)	10 (2%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	967	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/256 (99%)	253 (100%)	1 (0%)	91	90
1	B	249/256 (97%)	249 (100%)	0	100	100
All	All	503/512 (98%)	502 (100%)	1 (0%)	93	93

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	863	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	1204	-	3,3,3	0.42	0	2,2,2	0.31	0
2	RQZ	A	1201[B]	1	24,25,25	0.81	1 (4%)	27,35,35	0.76	0
4	EDO	A	1205	-	3,3,3	0.28	0	2,2,2	0.34	0
4	EDO	B	1205	-	3,3,3	0.32	0	2,2,2	0.56	0
2	RQZ	A	1201[A]	1	24,25,25	0.83	1 (4%)	27,35,35	0.72	0
4	EDO	A	1206	-	3,3,3	0.53	0	2,2,2	0.11	0
4	EDO	A	1203	-	3,3,3	0.61	0	2,2,2	0.19	0
4	EDO	B	1204	-	3,3,3	0.54	0	2,2,2	0.16	0
2	RQZ	B	1201[A]	1	24,25,25	0.82	0	27,35,35	0.61	0
3	PHU	B	1202	-	10,10,10	0.33	0	12,12,12	0.41	0
4	EDO	A	1207	-	3,3,3	0.36	0	2,2,2	0.43	0
4	EDO	B	1203	-	3,3,3	0.40	0	2,2,2	0.32	0
3	PHU	A	1202	-	10,10,10	0.43	0	12,12,12	0.30	0
2	RQZ	B	1201[B]	1	24,25,25	0.79	0	27,35,35	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1204	-	-	1/1/1/1	-
2	RQZ	A	1201[B]	1	-	2/14/14/14	0/3/3/3
4	EDO	A	1205	-	-	1/1/1/1	-
4	EDO	B	1205	-	-	0/1/1/1	-
2	RQZ	A	1201[A]	1	-	2/14/14/14	0/3/3/3
4	EDO	A	1206	-	-	0/1/1/1	-
4	EDO	A	1203	-	-	1/1/1/1	-
4	EDO	B	1204	-	-	1/1/1/1	-
2	RQZ	B	1201[A]	1	-	0/14/14/14	0/3/3/3
3	PHU	B	1202	-	-	2/4/4/4	0/1/1/1
4	EDO	A	1207	-	-	0/1/1/1	-
4	EDO	B	1203	-	-	0/1/1/1	-
3	PHU	A	1202	-	-	0/4/4/4	0/1/1/1
2	RQZ	B	1201[B]	1	-	2/14/14/14	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201[A]	RQZ	C3-N1	-2.50	1.33	1.37
2	A	1201[B]	RQZ	C3-N1	-2.38	1.33	1.37

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	1201[A]	RQZ	C11-C10-C9-N2
4	A	1204	EDO	O1-C1-C2-O2
2	A	1201[A]	RQZ	C11-C10-C9-O
2	A	1201[B]	RQZ	C4-C5-C6-C14
2	B	1201[B]	RQZ	C4-C5-C6-C14
4	A	1203	EDO	O1-C1-C2-O2
3	B	1202	PHU	C4-C2-N1-C1
4	B	1204	EDO	O1-C1-C2-O2
3	B	1202	PHU	C3-C2-N1-C1
4	A	1205	EDO	O1-C1-C2-O2
2	A	1201[B]	RQZ	C4-C5-C6-C7
2	B	1201[B]	RQZ	C4-C5-C6-C7


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1201[B]	RQZ	1	0
2	A	1201[A]	RQZ	2	0
2	B	1201[B]	RQZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/294 (98%)	0.44	21 (7%) 15 23	15, 27, 58, 97	0
1	B	285/294 (96%)	0.58	26 (9%) 9 15	17, 32, 66, 98	0
All	All	575/588 (97%)	0.51	47 (8%) 11 18	15, 29, 64, 98	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	985	GLU	7.8
1	B	987	GLY	7.6
1	A	833	PHE	7.3
1	B	833	PHE	6.0
1	A	859	HIS	5.8
1	A	814	PRO	5.5
1	A	832	ASN	5.5
1	B	1102	SER	5.1
1	B	832	ASN	4.7
1	B	891	TYR	4.6
1	B	814	PRO	4.5
1	B	862	PRO	4.2
1	A	834	GLY	4.1
1	A	860	SER	4.0
1	A	916	ARG	3.8
1	A	831	GLY	3.7
1	B	1041	GLU	3.4
1	A	858	GLN	3.4
1	A	862	PRO	3.3
1	B	988	GLN	3.1
1	B	1101	GLY	3.1
1	A	820	ARG	3.0
1	B	845	GLY	2.9
1	B	1043	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	848	THR	2.8
1	A	987	GLY	2.8
1	A	864	GLN	2.7
1	B	1040	SER	2.7
1	B	816	ILE	2.7
1	A	845	GLY	2.7
1	B	1048	SER	2.6
1	B	989	SER	2.6
1	A	815	THR	2.6
1	B	1042	ARG	2.5
1	B	1097	MET	2.5
1	A	861	GLY	2.4
1	B	866	ARG	2.4
1	B	990	PRO	2.4
1	A	844	LEU	2.3
1	B	986	PRO	2.2
1	B	1100	SER	2.2
1	B	861	GLY	2.2
1	A	977	ASP	2.1
1	B	897	SER	2.1
1	A	843	PRO	2.1
1	B	844	LEU	2.0
1	A	988	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	B	1205	4/4	0.82	0.24	55,55,56,60	0
4	EDO	A	1204	4/4	0.85	0.18	50,51,51,54	0
4	EDO	A	1205	4/4	0.86	0.18	33,33,36,38	0
3	PHU	B	1202	10/10	0.86	0.14	26,28,29,30	0
4	EDO	A	1206	4/4	0.88	0.11	32,33,33,36	0
4	EDO	A	1207	4/4	0.89	0.13	45,46,49,50	0
4	EDO	B	1204	4/4	0.92	0.20	28,28,29,30	0
2	RQZ	A	1201[B]	23/23	0.94	0.12	20,21,24,25	23
2	RQZ	A	1201[A]	23/23	0.94	0.12	23,25,28,28	23
2	RQZ	B	1201[A]	23/23	0.95	0.11	14,16,17,17	23
3	PHU	A	1202	10/10	0.95	0.11	17,19,20,21	0
2	RQZ	B	1201[B]	23/23	0.95	0.11	15,16,18,20	23
4	EDO	A	1203	4/4	0.97	0.08	25,26,26,27	0
4	EDO	B	1203	4/4	0.97	0.11	27,28,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around RQZ A 1201 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around RQZ B 1201 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.