



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2024 – 03:18 PM EDT

PDB ID : 1ARP  
Title : Crystal structure of the fungal peroxidase from *Arthromyces ramosus* at 1.9 angstroms resolution: structural comparisons with the lignin and cytochrome C peroxidases  
Authors : Kunishima, N.; Fukuyama, K.  
Deposited on : 1993-06-18  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	 74% 19%
2	B	2	 50% 50%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2784 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2465	1537	421	492	15	0	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
4	A	1	43	34	1	4	4	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	246	Total	O	0	0
			246	246		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PEROXIDASE

Chain A: 



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.57Å 74.57Å 117.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 1.90 7.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (7.00-1.90) 71.5 (7.00-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.174 , (Not available) 0.164 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.9	Xtrriage
Anisotropy	0.170	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 72.8	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2784	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.81	0/2521	1.59	31/3436 (0.9%)

There are no bond length outliers.

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	48	ARG	NE-CZ-NH2	-18.93	110.84	120.30
1	A	48	ARG	NE-CZ-NH1	15.53	128.06	120.30
1	A	141	ARG	NE-CZ-NH1	14.06	127.33	120.30
1	A	272	ARG	NE-CZ-NH1	11.32	125.96	120.30
1	A	287	ARG	NE-CZ-NH1	8.89	124.75	120.30
1	A	218	LEU	CA-CB-CG	8.60	135.08	115.30
1	A	287	ARG	NE-CZ-NH2	-8.39	116.10	120.30
1	A	272	ARG	NE-CZ-NH2	-8.07	116.27	120.30
1	A	210	THR	N-CA-CB	-7.93	95.24	110.30
1	A	166	ARG	NE-CZ-NH1	7.49	124.05	120.30
1	A	259	TRP	CD1-CG-CD2	7.25	112.10	106.30
1	A	26	TRP	CD1-CG-CD2	7.21	112.07	106.30
1	A	243	MET	CG-SD-CE	-7.16	88.75	100.20
1	A	26	TRP	CE2-CD2-CG	-7.14	101.58	107.30
1	A	244	ARG	NE-CZ-NH2	-7.12	116.74	120.30
1	A	259	TRP	CE2-CD2-CG	-6.94	101.75	107.30
1	A	26	TRP	CG-CD2-CE3	6.74	139.97	133.90
1	A	242	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	A	126	MET	CA-CB-CG	-6.38	102.45	113.30
1	A	141	ARG	NE-CZ-NH2	-6.22	117.19	120.30
1	A	271	GLN	CG-CD-NE2	5.85	130.75	116.70
1	A	221	THR	N-CA-CB	-5.69	99.49	110.30
1	A	325	GLU	CA-CB-CG	5.57	125.66	113.40
1	A	242	ARG	NE-CZ-NH2	-5.54	117.53	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	251	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	169	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	141	ARG	NH1-CZ-NH2	-5.39	113.47	119.40
1	A	10	VAL	CA-C-N	-5.30	105.54	117.20
1	A	202	ASP	CA-CB-CG	5.28	125.02	113.40
1	A	26	TRP	CB-CG-CD1	-5.19	120.25	127.00
1	A	244	ARG	NE-CZ-NH1	5.10	122.85	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2465	0	2375	36	0
2	B	28	0	25	0	0
3	A	2	0	0	0	0
4	A	43	0	30	2	0
5	A	246	0	0	5	0
All	All	2784	0	2430	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:PRO:HB3	1:A:243:MET:HE1	1.82	0.62
1:A:217:LEU:HD13	1:A:250:ALA:HB1	1.82	0.61
1:A:219:LYS:HG3	1:A:317:ASP:O	2.02	0.59
1:A:86:ILE:HG12	1:A:147:PRO:HB3	1.85	0.59
1:A:54:VAL:HG23	1:A:118:ILE:HG22	1.86	0.57
1:A:138:LEU:HD12	1:A:292:ASP:HA	1.87	0.57
1:A:86:ILE:HG13	1:A:87:GLU:N	2.21	0.55
1:A:236:PRO:HD3	1:A:242:ARG:NH2	2.24	0.53

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:LEU:HD13	1:A:276:ALA:HB1	1.91	0.52
1:A:301:VAL:HG12	5:A:482:HOH:O	2.08	0.52
1:A:318:ILE:HD11	1:A:320:VAL:HG22	1.91	0.52
1:A:12:CYS:HB3	5:A:350:HOH:O	2.10	0.50
1:A:308:VAL:HG12	1:A:331:ALA:HB3	1.93	0.49
1:A:34:GLN:HE21	1:A:34:GLN:HA	1.78	0.49
1:A:48:ARG:HD3	1:A:187:ALA:HA	1.95	0.48
1:A:215:GLU:HA	1:A:218:LEU:CD2	2.43	0.48
1:A:156:PRO:HB3	1:A:243:MET:CE	2.44	0.47
1:A:210:THR:HG22	5:A:530:HOH:O	2.14	0.46
1:A:26:TRP:HA	1:A:29:VAL:CG1	2.45	0.46
1:A:85:ASN:H	1:A:85:ASN:HD22	1.63	0.46
1:A:215:GLU:HA	1:A:218:LEU:HD22	1.97	0.46
1:A:221:THR:HG22	1:A:222:THR:OG1	2.17	0.44
1:A:48:ARG:HD2	4:A:345:HEM:O2D	2.19	0.43
1:A:51:LEU:HB2	4:A:345:HEM:HMD3	2.01	0.43
1:A:192:LEU:HG	1:A:230:PHE:O	2.19	0.43
1:A:297:ILE:HA	1:A:298:PRO:HD2	1.93	0.42
1:A:309:ILE:HA	1:A:310:PRO:HD3	1.79	0.42
1:A:280:MET:HA	1:A:283:LEU:HD13	2.02	0.42
1:A:221:THR:HG21	5:A:538:HOH:O	2.20	0.41
1:A:20:ASN:HB2	5:A:563:HOH:O	2.21	0.41
1:A:32:ASP:OD2	1:A:110:HIS:HE1	2.04	0.41
1:A:168:GLY:HA2	1:A:172:PHE:O	2.21	0.41
1:A:238:PRO:HB2	1:A:339:SER:HA	2.02	0.40
1:A:51:LEU:O	1:A:54:VAL:HG12	2.22	0.40
1:A:53:ILE:HG13	1:A:96:LEU:HD22	2.04	0.40
1:A:61:PHE:HB2	1:A:75:GLY:HA2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/344 (97%)	325 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/273 (99%)	242 (89%)	29 (11%)	5	2

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	VAL
1	A	11	THR
1	A	13	PRO
1	A	29	VAL
1	A	34	GLN
1	A	50	ILE
1	A	63	PRO
1	A	65	LEU
1	A	85	ASN
1	A	86	ILE
1	A	138	LEU
1	A	146	GLN
1	A	181	LEU
1	A	186	LEU
1	A	192	LEU
1	A	193	ASN
1	A	201	LEU
1	A	210	THR
1	A	217	LEU
1	A	218	LEU
1	A	221	THR
1	A	227	SER
1	A	234	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	237	PHE
1	A	248	LEU
1	A	271	GLN
1	A	302	SER
1	A	318	ILE
1	A	329	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	85	ASN
1	A	110	HIS
1	A	128	ASN
1	A	193	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	B	1	2,1	14,14,15	0.63	0	17,19,21	1.37	3 (17%)
2	NAG	B	2	2	14,14,15	0.61	0	17,19,21	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-C2-N2	3.35	115.71	110.43
2	B	1	NAG	O5-C1-C2	-2.42	107.55	111.29
2	B	1	NAG	C6-C5-C4	-2.34	107.26	113.02

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	HEM	A	345	5,1	42,50,50	1.50	10 (23%)	46,82,82	1.16	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	A	345	5,1	-	4/12/54/54	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	345	HEM	C3C-CAC	-3.55	1.39	1.47
4	A	345	HEM	C3C-C4C	3.29	1.46	1.41
4	A	345	HEM	O2D-CGD	-3.01	1.20	1.30
4	A	345	HEM	CBB-CAB	3.01	1.44	1.30
4	A	345	HEM	C3C-C2C	-2.65	1.36	1.40
4	A	345	HEM	O2A-CGA	-2.54	1.22	1.30
4	A	345	HEM	CAB-C3B	-2.44	1.40	1.47
4	A	345	HEM	CBC-CAC	2.43	1.44	1.29
4	A	345	HEM	CHB-C1B	2.39	1.40	1.34
4	A	345	HEM	C4D-C3D	-2.17	1.41	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	345	HEM	C4B-CHC-C1C	2.81	126.27	122.56
4	A	345	HEM	C3B-C4B-NB	2.54	111.29	109.47
4	A	345	HEM	O2D-CGD-CBD	2.35	121.44	114.00
4	A	345	HEM	CMA-C3A-C4A	-2.33	125.04	128.46
4	A	345	HEM	C4C-CHD-C1D	2.06	125.28	122.56

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	345	HEM	C2B-C3B-CAB-CBB
4	A	345	HEM	C4B-C3B-CAB-CBB
4	A	345	HEM	CAA-CBA-CGA-O1A
4	A	345	HEM	CAA-CBA-CGA-O2A

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	345	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	336/344 (97%)	-0.83	0 <b>100</b> <b>100</b>	7, 14, 25, 31	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	2	14/15	0.92	0.05	13,18,21,21	0
2	NAG	B	1	14/15	0.94	0.05	15,17,22,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CA	A	346	1/1	0.98	0.03	12,12,12,12	0
4	HEM	A	345	43/43	0.98	0.04	8,11,14,19	0
3	CA	A	347	1/1	0.99	0.05	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.