



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2024 – 05:37 PM EST

PDB ID : 3ASP  
Title : Crystal structure of P domain from Norovirus Funabashi258 stain in the complex with A-antigen  
Authors : Kubota, T.; Kumagai, A.; Itoh, H.; Furukawa, S.; Narimatsu, H.; Wakita, T.; Ishii, K.; Takeda, N.; Someya, Y.; Shirato, H.  
Deposited on : 2010-12-17  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	 7% 91% 5%
1	B	326	 8% 88% 5% 7%
2	C	4	 25% 75%
2	D	4	 50% 50%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	311	Total	C	N	O	S	0	2	0
			2392	1530	403	450	9			
1	B	302	Total	C	N	O	S	0	2	0
			2325	1490	391	435	9			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	216	GLY	-	expression tag	UNP Q8JW44
A	217	PRO	-	expression tag	UNP Q8JW44
A	218	LEU	-	expression tag	UNP Q8JW44
A	219	GLY	-	expression tag	UNP Q8JW44
A	220	SER	-	expression tag	UNP Q8JW44
B	216	GLY	-	expression tag	UNP Q8JW44
B	217	PRO	-	expression tag	UNP Q8JW44
B	218	LEU	-	expression tag	UNP Q8JW44
B	219	GLY	-	expression tag	UNP Q8JW44
B	220	SER	-	expression tag	UNP Q8JW44

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	4	Total	C	N	O	0	0	0
			50	28	2	20			
2	D	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Na 1	0	0
3	B	1	Total 1	Na 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	205	Total 205	O 205	0	0
4	B	202	Total 202	O 202	0	0



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.62Å 74.62Å 107.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.23 – 1.60 35.23 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.23-1.60) 99.9 (35.23-1.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.22 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.182 , 0.201 0.182 , 0.201	Depositor DCC
$R_{free}$ test set	4406 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtriage
Anisotropy	0.005	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 51.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.006 for -h,-k,l 0.028 for h,-h-k,-l 0.016 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5226	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A2G, FUC, NA, GAL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2471	0.52	0/3383
1	B	0.36	0/2404	0.53	0/3292
All	All	0.36	0/4875	0.53	0/6675

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2392	0	2322	7	0
1	B	2325	0	2254	11	0
2	C	50	0	44	0	0
2	D	50	0	44	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	205	0	0	0	0
4	B	202	0	0	1	0
All	All	5226	0	4664	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:524:ARG:HG2	1:B:524:ARG:HH11	1.31	0.95
1:B:254:MET:HG2	1:B:507:PRO:HB2	1.77	0.65
1:B:440:ASN:HD22	1:B:446:ASN:ND2	1.96	0.62
1:B:529:LYS:HD3	4:B:2004:HOH:O	2.06	0.55
1:A:440:ASN:HD22	1:A:446:ASN:ND2	2.05	0.55
1:B:524:ARG:HG2	1:B:524:ARG:NH1	2.09	0.54
1:A:470:ASP:OD2	1:A:524:ARG:HG2	2.09	0.52
1:A:287:GLN:HG3	1:B:250:LEU:HD11	1.95	0.48
1:A:273:ILE:HG12	1:A:492:PRO:HA	1.96	0.47
1:B:273:ILE:HG12	1:B:492:PRO:HA	1.97	0.47
1:A:315:PHE:HA	1:A:363:HIS:CE1	2.53	0.43
1:A:248:PRO:O	1:B:290:ARG:NH2	2.50	0.43
1:A:363:HIS:CE1	1:A:414:ALA:H	2.37	0.42
1:B:524:ARG:NH1	1:B:524:ARG:CG	2.77	0.42
1:B:400:LEU:HD13	1:B:448:VAL:HG22	2.01	0.42
1:B:373:PHE:CE2	1:B:375:HIS:HB2	2.55	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/326 (95%)	309 (99%)	2 (1%)	0	100	100
1	B	302/326 (93%)	301 (100%)	1 (0%)	0	100	100
All	All	613/652 (94%)	610 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/277 (96%)	264 (99%)	3 (1%)	73	57
1	B	260/277 (94%)	257 (99%)	3 (1%)	71	54
All	All	527/554 (95%)	521 (99%)	6 (1%)	73	57

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	317	SER
1	A	523	SER
1	A	524	ARG
1	B	317	SER
1	B	483	ARG
1	B	524	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	296	ASN
1	A	354	GLN
1	A	374	ASN
1	A	415	ASN
1	A	446	ASN
1	B	296	ASN
1	B	297	GLN
1	B	415	ASN
1	B	446	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	1	2	15,15,15	0.42	0	21,21,21	1.05	1 (4%)
2	GAL	C	2	2	11,11,12	0.51	0	15,15,17	1.24	3 (20%)
2	FUC	C	3	2	10,10,11	0.61	0	14,14,16	0.56	0
2	A2G	C	4	2	14,14,15	0.23	0	17,19,21	0.78	1 (5%)
2	NAG	D	1	2	15,15,15	0.42	0	21,21,21	0.63	0
2	GAL	D	2	2	11,11,12	0.58	0	15,15,17	1.30	3 (20%)
2	FUC	D	3	2	10,10,11	0.62	0	14,14,16	0.51	0
2	A2G	D	4	2	14,14,15	0.25	0	17,19,21	0.94	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2	-	1/6/26/26	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	FUC	C	3	2	-	-	0/1/1/1
2	A2G	C	4	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	2	-	0/6/26/26	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	FUC	D	3	2	-	-	0/1/1/1
2	A2G	D	4	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4	A2G	C1-O5-C5	3.08	116.37	112.19
2	D	2	GAL	C1-C2-C3	2.72	113.01	109.67
2	C	2	GAL	O3-C3-C2	-2.64	104.93	109.99
2	D	2	GAL	O3-C3-C2	-2.57	105.07	109.99
2	C	1	NAG	C3-C4-C5	2.33	114.40	110.24
2	C	2	GAL	C1-C2-C3	2.33	112.53	109.67
2	C	4	A2G	C1-O5-C5	2.13	115.07	112.19
2	C	2	GAL	O5-C5-C6	2.02	110.37	107.20
2	D	2	GAL	C1-O5-C5	2.00	114.90	112.19

There are no chirality outliers.


All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	4	A2G	O5-C5-C6-O6
2	D	4	A2G	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/326 (95%)	0.26	24 (7%) <b>13</b> <b>11</b>	10, 15, 29, 41	0
1	B	302/326 (92%)	0.28	26 (8%) <b>10</b> <b>9</b>	10, 16, 30, 38	0
All	All	613/652 (94%)	0.27	50 (8%) <b>11</b> <b>10</b>	10, 16, 30, 41	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	315	PHE	17.4
1	A	315	PHE	15.7
1	A	410	LEU	13.6
1	B	410	LEU	11.4
1	B	314	ALA	10.1
1	B	409	SER	9.9
1	A	409	SER	9.6
1	A	412	GLN	8.2
1	A	314	ALA	8.2
1	A	503	VAL	7.9
1	A	408	SER	7.4
1	A	411	SER	6.9
1	A	414	ALA	6.8
1	A	225	ILE	6.8
1	A	316	ASP	6.8
1	B	412	GLN	6.6
1	B	503	VAL	6.1
1	B	525	PHE	5.6
1	B	316	ASP	5.3
1	B	408	SER	5.3
1	B	504	GLY	5.0
1	B	425	PHE	4.9
1	A	407	GLY	4.8
1	A	504	GLY	4.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	230	ARG	4.2
1	B	411	SER	4.2
1	A	413	ALA	3.9
1	B	414	ALA	3.8
1	B	313	MET	3.5
1	A	415	ASN	3.4
1	B	505	ALA	3.4
1	A	224	THR	3.2
1	A	317	SER	3.1
1	A	534	ALA	3.0
1	B	317	SER	3.0
1	B	413	ALA	3.0
1	B	483	ARG	2.9
1	B	480	ASP	2.8
1	B	501	ASN	2.7
1	A	261	SER	2.6
1	A	358	GLN	2.6
1	B	231	ALA	2.6
1	B	502	GLY	2.5
1	A	288	LEU	2.5
1	A	313	MET	2.4
1	A	227	GLN	2.4
1	A	483	ARG	2.3
1	B	415	ASN	2.3
1	B	524	ARG	2.2
1	B	407	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	1	15/15	0.62	0.28	31,36,38,39	0
2	NAG	D	1	15/15	0.69	0.31	32,36,37,38	0


*Continued on next page...*




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GAL	D	2	11/12	0.78	0.21	22,27,30,32	0
2	FUC	C	3	10/11	0.84	0.14	24,26,27,27	0
2	GAL	C	2	11/12	0.90	0.09	19,24,27,28	0
2	FUC	D	3	10/11	0.92	0.14	25,28,28,29	0
2	A2G	C	4	14/15	0.95	0.07	14,15,17,20	0
2	A2G	D	4	14/15	0.95	0.08	14,16,20,24	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	A	701	1/1	0.95	0.11	33,33,33,33	0
3	NA	B	701	1/1	0.98	0.09	24,24,24,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.