



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 16, 2021 – 02:07 pm BST

PDB ID : 7AT5
Title : Structure of protein kinase ck2 catalytic subunit (csnk2a1 gene product) in complex with the bivalent inhibitor KN2
Authors : Lindenblatt, D.; Applegate, V.; Nickelsen, A.; Klussmann, M.; Neundorf, I.; Goetz, C.; Jose, J.; Niefind, K.
Deposited on : 2020-10-29
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	349	 91% 5%
1	B	349	 90% 5% 5%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6207 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2829	1810	498	508	13	0	3	0
1	B	331	2832	1811	501	508	12	0	4	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP P68400
A	-12	GLY	-	expression tag	UNP P68400
A	-11	SER	-	expression tag	UNP P68400
A	-10	SER	-	expression tag	UNP P68400
A	-9	HIS	-	expression tag	UNP P68400
A	-8	HIS	-	expression tag	UNP P68400
A	-7	HIS	-	expression tag	UNP P68400
A	-6	HIS	-	expression tag	UNP P68400
A	-5	HIS	-	expression tag	UNP P68400
A	-4	HIS	-	expression tag	UNP P68400
A	-3	SER	-	expression tag	UNP P68400
A	-2	GLN	-	expression tag	UNP P68400
A	-1	ASP	-	expression tag	UNP P68400
A	0	PRO	-	expression tag	UNP P68400
B	-13	MET	-	initiating methionine	UNP P68400
B	-12	GLY	-	expression tag	UNP P68400
B	-11	SER	-	expression tag	UNP P68400
B	-10	SER	-	expression tag	UNP P68400
B	-9	HIS	-	expression tag	UNP P68400
B	-8	HIS	-	expression tag	UNP P68400
B	-7	HIS	-	expression tag	UNP P68400
B	-6	HIS	-	expression tag	UNP P68400
B	-5	HIS	-	expression tag	UNP P68400
B	-4	HIS	-	expression tag	UNP P68400
B	-3	SER	-	expression tag	UNP P68400

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLN	-	expression tag	UNP P68400
B	-1	ASP	-	expression tag	UNP P68400
B	0	PRO	-	expression tag	UNP P68400

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is {N}'-[2-(3,4-dichlorophenyl)ethyl]- {N}-[4-[4,5,6,7-tetrakis(bromanyl)benzimidazol-1-yl]butyl]butanediamide (three-letter code: RXE) (formula: C₂₃H₂₂Br₄Cl₂N₄O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	Br	C	Cl	N			O
3	A	1	35	4	23	2	4	2	0	0
3	B	1	35	4	23	2	4	2	0	0

- Molecule 4 is 2-(3,4-dichlorophenyl)ethanamine (three-letter code: 42J) (formula: $C_8H_9Cl_2N$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	Cl	N		
4	A	1	11	8	2	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	Cl	N		
4	B	1	11	8	2	1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	B	1	4	2	2	0	0
5	B	1	4	2	2	0	0
5	B	1	4	2	2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	210	211	211	0	1
6	B	209	211	211	0	2

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: Casein kinase II subunit alpha

Chain A:  91% 5%



- Molecule 1: Casein kinase II subunit alpha

Chain B:  90% 5% 5%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	127.59Å 127.59Å 124.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	73.06 – 1.77	Depositor
% Data completeness (in resolution range)	74.9 (73.06-1.77)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.174 , 0.195	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6207	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 42J, EDO, RXE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2904	0.49	0/3925
1	B	0.31	0/2907	0.49	0/3930
All	All	0.31	0/5811	0.49	0/7855

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2829	0	2769	7	0
1	B	2832	0	2773	11	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
3	A	35	0	0	0	0
3	B	35	0	0	1	0
4	A	11	0	9	0	0
4	B	11	0	9	0	0
5	B	12	0	18	1	0
6	A	211	0	0	0	0
6	B	211	0	0	0	0
All	All	6207	0	5578	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:229:LYS:HE3	5:B:401:EDO:H11	1.76	0.68
1:A:78:ILE:HD13	1:A:109:PRO:HG2	1.86	0.58
1:B:121:PHE:HD1	1:B:124:LEU:HD23	1.72	0.54
1:A:134:ARG:HG2	1:A:323:TYR:CZ	2.46	0.51
1:A:261:TYR:OH	1:A:310:GLN:HG3	2.11	0.50
1:B:121:PHE:HE1	1:B:164:ILE:HB	1.78	0.49
1:B:134:ARG:HG2	1:B:323:TYR:CZ	2.48	0.48
1:A:159:PRO:HG3	1:A:221:MET:HG3	1.94	0.48
1:B:286:HIS:CE1	1:B:288:GLU:HB2	2.49	0.47
1:A:74:LYS:HA	1:A:74:LYS:HD3	1.72	0.46
1:A:286:HIS:CE1	1:A:288:GLU:HB2	2.52	0.45
1:B:279:LYS:HB3	1:B:283:ARG:HD3	1.99	0.45
1:B:266:ASP:OD1	1:B:268:ARG:HG2	2.19	0.42
1:B:159:PRO:HD3	1:B:221:MET:HG2	2.02	0.42
1:B:221:MET:O	1:B:225[B]:MET:HG2	2.21	0.41
1:B:116:VAL:O	3:B:405:RXE:BR2	2.95	0.40
1:B:240:ASP:O	1:B:244:ARG:HG2	2.21	0.40
1:A:49:LYS:O	1:A:71:LYS:HD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/349 (95%)	326 (98%)	7 (2%)	0	100	100
1	B	333/349 (95%)	325 (98%)	8 (2%)	0	100	100
All	All	666/698 (95%)	651 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/320 (96%)	307 (100%)	1 (0%)	92	90
1	B	308/320 (96%)	306 (99%)	2 (1%)	86	82
All	All	616/640 (96%)	613 (100%)	3 (0%)	92	86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	THR
1	B	225[A]	MET
1	B	225[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	403	-	4,4,4	0.16	0	6,6,6	0.06	0
4	42J	B	406	-	10,11,11	0.15	0	13,14,14	0.35	0
5	EDO	B	401	-	3,3,3	0.50	0	2,2,2	0.28	0
4	42J	A	403	-	10,11,11	0.14	0	13,14,14	0.26	0
3	RXE	A	402	-	33,37,37	1.89	10 (30%)	40,51,51	1.15	4 (10%)
3	RXE	B	405	-	33,37,37	1.88	10 (30%)	40,51,51	1.09	3 (7%)
5	EDO	B	404	-	3,3,3	0.48	0	2,2,2	0.37	0
2	SO4	A	404	-	4,4,4	0.14	0	6,6,6	0.07	0
2	SO4	B	402	-	4,4,4	0.12	0	6,6,6	0.16	0
5	EDO	B	407	-	3,3,3	0.49	0	2,2,2	0.26	0
2	SO4	A	401	-	4,4,4	0.15	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	42J	B	406	-	-	0/3/3/3	0/1/1/1
5	EDO	B	401	-	-	0/1/1/1	-
4	42J	A	403	-	-	0/3/3/3	0/1/1/1
3	RXE	A	402	-	-	3/19/19/19	0/3/3/3
3	RXE	B	405	-	-	6/19/19/19	0/3/3/3
5	EDO	B	404	-	-	0/1/1/1	-
5	EDO	B	407	-	-	0/1/1/1	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	405	RXE	C12-N11	5.58	1.46	1.33
3	A	402	RXE	C12-N11	5.49	1.45	1.33
3	B	405	RXE	C16-N18	5.27	1.45	1.33
3	A	402	RXE	C16-N18	5.25	1.45	1.33
3	A	402	RXE	BR2-C30	3.32	1.96	1.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	405	RXE	BR2-C30	3.20	1.96	1.89
3	B	405	RXE	BR3-C32	2.91	1.95	1.89
3	A	402	RXE	BR3-C32	2.88	1.95	1.89
3	B	405	RXE	BR1-C28	2.84	1.96	1.89
3	A	402	RXE	BR1-C28	2.83	1.96	1.89
3	A	402	RXE	BR4-C34	2.30	1.94	1.89
3	A	402	RXE	C02-CL1	2.26	1.79	1.73
3	B	405	RXE	BR4-C34	2.23	1.94	1.89
3	B	405	RXE	O13-C12	-2.22	1.18	1.23
3	B	405	RXE	C07-CL2	2.19	1.78	1.73
3	A	402	RXE	O17-C16	-2.19	1.18	1.23
3	A	402	RXE	C07-CL2	2.18	1.78	1.73
3	B	405	RXE	C02-CL1	2.16	1.78	1.73
3	B	405	RXE	O17-C16	-2.12	1.19	1.23
3	A	402	RXE	O13-C12	-2.09	1.19	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	RXE	C19-N18-C16	-2.73	117.77	122.84
3	A	402	RXE	C15-C16-N18	2.65	120.88	116.42
3	B	405	RXE	C30-C28-C27	-2.15	119.06	121.33
3	A	402	RXE	C07-C06-C05	-2.14	119.00	120.46
3	B	405	RXE	C15-C16-N18	2.14	120.02	116.42
3	B	405	RXE	C19-N18-C16	-2.10	118.94	122.84
3	A	402	RXE	O17-C16-N18	-2.00	119.23	123.01

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	402	RXE	N11-C12-C14-C15
3	B	405	RXE	O13-C12-C14-C15
3	B	405	RXE	N11-C12-C14-C15
3	A	402	RXE	O13-C12-C14-C15
3	B	405	RXE	C12-C14-C15-C16
3	B	405	RXE	C19-C20-C21-C22
3	A	402	RXE	C19-C20-C21-C22
3	B	405	RXE	C21-C22-N23-C24
3	B	405	RXE	C09-C10-N11-C12


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	401	EDO	1	0
3	B	405	RXE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.