



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2021 – 04:12 PM EDT

PDB ID : 2AUH  
Title : Crystal structure of the Grb14 BPS region in complex with the insulin receptor tyrosine kinase  
Authors : Depetris, R.S.; Hu, J.; Gimpelevich, I.; Holt, L.J.; Daly, R.J.; Hubbard, S.R.  
Deposited on : 2005-08-27  
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
2	B	59	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2639 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Insulin receptor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	300	2363	1494	401	446	3	19	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1158	PTR	TYR	modified residue	UNP P06213
A	1162	PTR	TYR	modified residue	UNP P06213
A	1163	PTR	TYR	modified residue	UNP P06213
A	1251	ASN	LYS	variant	UNP P06213
A	981	SER	CYS	engineered mutation	UNP P06213

- Molecule 2 is a protein called Growth factor receptor-bound protein 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	37	274	170	47	56	1	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	411	SER	CYS	engineered mutation	UNP Q14449


- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Insulin receptor



- Molecule 2: Growth factor receptor-bound protein 14



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.18Å 127.18Å 65.92Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.27 – 3.20 36.71 – 3.20	Depositor EDS
% Data completeness (in resolution range)	97.2 (29.27-3.20) 97.2 (36.71-3.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.82 (at 3.18Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.223 , 0.254 0.207 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.9	Xtrriage
Anisotropy	0.529	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 25.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.054 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2639	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/2365	0.66	0/3201
2	B	0.42	0/277	0.68	0/375
All	All	0.40	0/2642	0.66	0/3576

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2363	0	2276	103	0
2	B	274	0	258	8	0
3	A	2	0	0	0	0
All	All	2639	0	2534	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1228:LEU:H	1:A:1228:LEU:HD12	1.26	1.00
1:A:1018:ILE:HD11	1:A:1025:THR:HB	1.49	0.92
1:A:1214:SER:HB3	1:A:1217:GLN:HG3	1.56	0.88
1:A:1007:PHE:HB2	1:A:1040:GLU:HG2	1.54	0.88
1:A:1063:LEU:HB2	1:A:1075:VAL:HG12	1.59	0.85
1:A:999:LEU:HG	1:A:1013:GLY:HA2	1.58	0.85
1:A:1215:ASN:O	1:A:1218:VAL:HG12	1.82	0.80
1:A:1007:PHE:CB	1:A:1040:GLU:HG2	2.18	0.73
1:A:1003:GLY:O	1:A:1009:MET:HG3	1.91	0.71
1:A:1228:LEU:HD12	1:A:1228:LEU:N	2.03	0.70
1:A:1028:ALA:HB2	1:A:1078:LEU:HA	1.74	0.69
1:A:1018:ILE:HG13	1:A:1019:ILE:H	1.57	0.67
1:A:1050:VAL:HG12	1:A:1054:PHE:CE2	2.31	0.65
1:A:1018:ILE:HG13	1:A:1019:ILE:N	2.12	0.64
1:A:1057:HIS:O	1:A:1147:LYS:HE2	1.98	0.63
1:A:1079:MET:HE1	1:A:1141:ALA:HB2	1.80	0.63
1:A:1228:LEU:H	1:A:1228:LEU:CD1	2.07	0.63
2:B:389:ILE:HD12	2:B:398:VAL:HG21	1.80	0.62
1:A:1026:ARG:HH11	1:A:1026:ARG:CB	2.14	0.61
1:A:1038:LEU:HD21	2:B:404:LEU:HD23	1.83	0.60
1:A:1046:ASN:O	1:A:1050:VAL:HG23	2.02	0.60
1:A:1059:VAL:O	1:A:1060:VAL:C	2.41	0.59
1:A:1050:VAL:HG12	1:A:1054:PHE:CZ	2.39	0.57
1:A:1271:PHE:HB3	1:A:1272:PRO:HD3	1.86	0.57
1:A:1108:GLU:O	1:A:1112:MET:HG3	2.05	0.57
1:A:1089:ARG:NE	1:A:1092:ARG:NH2	2.53	0.56
1:A:1174:ARG:HH11	1:A:1174:ARG:HG2	1.70	0.56
2:B:389:ILE:HD12	2:B:398:VAL:CG2	2.35	0.56
1:A:1111:GLN:O	1:A:1115:GLU:HG3	2.06	0.56
1:A:1170:LEU:O	1:A:1171:LEU:HD23	2.06	0.56
1:A:1026:ARG:HH11	1:A:1026:ARG:HB2	1.70	0.56
1:A:1072:THR:O	1:A:1073:LEU:HD23	2.06	0.55
1:A:1179:GLU:OE2	1:A:1253:ARG:NH1	2.40	0.55
1:A:1106:LEU:O	1:A:1110:ILE:HG12	2.06	0.55
1:A:1125:ALA:C	1:A:1127:LYS:H	2.10	0.54
1:A:1047:GLU:O	1:A:1051:MET:HG3	2.07	0.54
1:A:1110:ILE:HG23	1:A:1263:LEU:HD22	1.90	0.53
1:A:1160:THR:O	1:A:1161:ASP:HB2	2.07	0.53
1:A:1032:VAL:O	1:A:1032:VAL:HG13	2.09	0.52
1:A:1162:PTR:CD1	1:A:1162:PTR:C	2.88	0.52
1:A:1263:LEU:O	1:A:1265:ASP:N	2.43	0.51
1:A:1210:TYR:CZ	1:A:1228:LEU:HD11	2.46	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1066:VAL:O	1:A:1072:THR:HG23	2.11	0.51
2:B:408:LYS:O	2:B:409:LYS:CB	2.59	0.51
1:A:1209:PRO:HB3	1:A:1228:LEU:HD21	1.93	0.51
1:A:1136:ARG:CZ	2:B:375:SER:OG	2.58	0.51
1:A:989:TRP:O	1:A:990:GLU:C	2.50	0.50
1:A:1105:THR:HG23	1:A:1108:GLU:HG3	1.93	0.50
1:A:1051:MET:HA	1:A:1054:PHE:CD2	2.47	0.50
1:A:1130:HIS:O	1:A:1131:ARG:HB2	2.12	0.50
1:A:1065:VAL:HG12	1:A:1066:VAL:N	2.28	0.49
1:A:1174:ARG:HG2	1:A:1174:ARG:NH1	2.27	0.48
2:B:386:SER:O	2:B:387:ARG:HB2	2.14	0.48
1:A:1194:SER:O	1:A:1195:PHE:C	2.50	0.48
1:A:1007:PHE:CD2	1:A:1008:GLY:N	2.83	0.47
1:A:1105:THR:HG22	1:A:1108:GLU:OE1	2.14	0.47
1:A:1063:LEU:HB2	1:A:1075:VAL:CG1	2.38	0.46
1:A:1213:LEU:HD22	1:A:1217:GLN:HB3	1.96	0.46
1:A:1261:ASN:O	1:A:1264:LYS:HG2	2.14	0.46
1:A:1032:VAL:HG12	1:A:1072:THR:HB	1.97	0.46
1:A:1173:VAL:HG13	1:A:1181:LEU:HD21	1.98	0.46
1:A:1105:THR:CG2	1:A:1108:GLU:HG3	2.45	0.46
1:A:1089:ARG:NE	1:A:1092:ARG:HH21	2.14	0.45
1:A:1079:MET:HG3	1:A:1139:MET:CB	2.45	0.45
1:A:1091:LEU:HD13	1:A:1104:PRO:HD3	1.98	0.45
1:A:1061:ARG:H	1:A:1077:GLU:HG2	1.80	0.45
1:A:1079:MET:HG3	1:A:1139:MET:HB3	1.99	0.45
1:A:1089:ARG:O	1:A:1092:ARG:HG3	2.17	0.45
1:A:1038:LEU:HD12	1:A:1038:LEU:O	2.17	0.44
1:A:995:LYS:HD3	1:A:1016:ARG:O	2.18	0.44
1:A:1089:ARG:CD	1:A:1092:ARG:HH21	2.30	0.43
1:A:1109:MET:CE	1:A:1235:PRO:HG2	2.49	0.43
1:A:1091:LEU:HD22	1:A:1102:PRO:O	2.18	0.43
1:A:1132:ASP:HB2	1:A:1153:MET:HB2	2.01	0.43
1:A:1109:MET:HE3	1:A:1235:PRO:HG2	2.00	0.43
1:A:987:ASP:O	1:A:989:TRP:N	2.52	0.42
1:A:1080:ALA:HB3	1:A:1142:HIS:HA	2.01	0.42
1:A:1210:TYR:CE1	1:A:1228:LEU:HG	2.55	0.42
1:A:1058:HIS:HA	1:A:1147:LYS:HG2	2.01	0.42
1:A:1163:PTR:C	1:A:1163:PTR:CD1	2.97	0.42
1:A:1192:MET:O	1:A:1195:PHE:HB3	2.20	0.42
1:A:1057:HIS:CG	1:A:1274:VAL:HB	2.55	0.42
1:A:1127:LYS:CG	1:A:1158:PTR:HA	2.49	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1049:SER:HA	1:A:1052:LYS:HD2	2.01	0.41
1:A:1101:ARG:NH2	1:A:1108:GLU:OE1	2.51	0.41
1:A:990:GLU:OE2	1:A:1066:VAL:HA	2.20	0.41
1:A:1108:GLU:O	1:A:1111:GLN:HB3	2.21	0.41
1:A:1218:VAL:HG13	1:A:1219:LEU:N	2.34	0.41
1:A:1219:LEU:C	1:A:1219:LEU:HD23	2.40	0.41
1:A:1192:MET:HG2	1:A:1254:PRO:O	2.21	0.41
1:A:1271:PHE:O	1:A:1275:SER:HB3	2.21	0.41
2:B:373:GLU:HA	2:B:373:GLU:OE1	2.20	0.41
1:A:1214:SER:O	1:A:1215:ASN:C	2.58	0.41
1:A:1032:VAL:O	1:A:1032:VAL:CG1	2.68	0.41
1:A:1041:ARG:O	1:A:1044:PHE:HB3	2.21	0.40
1:A:1065:VAL:O	1:A:1066:VAL:HG23	2.20	0.40
1:A:1170:LEU:HA	2:B:377:VAL:O	2.21	0.40
1:A:1218:VAL:CG1	1:A:1219:LEU:N	2.83	0.40
1:A:1262:LEU:HA	1:A:1262:LEU:HD12	1.86	0.40
1:A:1079:MET:HE3	1:A:1141:ALA:N	2.37	0.40
1:A:1175:TRP:CD1	1:A:1197:VAL:HG11	2.57	0.40
1:A:1269:PRO:C	1:A:1271:PHE:H	2.24	0.40
1:A:1089:ARG:HD3	1:A:1092:ARG:NE	2.37	0.40
1:A:1200:TRP:CH2	1:A:1207:GLU:HB2	2.56	0.40
1:A:1249:ASN:HA	1:A:1250:PRO:HD2	1.81	0.40
1:A:1157:ILE:O	1:A:1158:PTR:C	2.68	0.40
1:A:1179:GLU:HG2	1:A:1180:SER:N	2.36	0.40
1:A:1186:PHE:CD1	1:A:1186:PHE:N	2.90	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	295/306 (96%)	249 (84%)	36 (12%)	10 (3%)	<b>3</b> <b>24</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	B	35/59 (59%)	25 (71%)	7 (20%)	3 (9%)	1 4
All	All	330/365 (90%)	274 (83%)	43 (13%)	13 (4%)	3 22

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	990	GLU
1	A	1060	VAL
1	A	1264	LYS
1	A	988	GLU
1	A	1020	LYS
2	B	374	ASN
1	A	1023	ALA
2	B	387	ARG
2	B	399	ALA
1	A	986	PRO
1	A	1250	PRO
1	A	1270	SER
1	A	1066	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	249/265 (94%)	246 (99%)	3 (1%)	71 88
2	B	28/50 (56%)	28 (100%)	0	100 100
All	All	277/315 (88%)	274 (99%)	3 (1%)	73 88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1007	PHE
1	A	1026	ARG
1	A	1107	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1057	HIS
1	A	1070	GLN
1	A	1124	ASN
1	A	1233	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	PTR	A	1163	1	15,16,17	0.96	0	19,22,24	1.14	2 (10%)
1	PTR	A	1162	1	15,16,17	1.23	1 (6%)	19,22,24	0.80	0
1	PTR	A	1158	1	15,16,17	1.54	1 (6%)	19,22,24	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1163	1	-	1/10/11/13	0/1/1/1
1	PTR	A	1162	1	-	0/10/11/13	0/1/1/1
1	PTR	A	1158	1	-	2/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1158	PTR	P-OH	5.40	1.67	1.59
1	A	1162	PTR	P-OH	3.69	1.65	1.59

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1163	PTR	P-OH-CZ	3.27	134.24	123.75
1	A	1163	PTR	OH-P-O1P	-2.38	100.33	109.31

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1163	PTR	CZ-OH-P-O3P
1	A	1158	PTR	CA-CB-CG-CD1
1	A	1158	PTR	CA-CB-CG-CD2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1163	PTR	1	0
1	A	1162	PTR	1	0
1	A	1158	PTR	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	297/306 (97%)	-0.39	0 100 100	13, 40, 86, 101	0
2	B	37/59 (62%)	-0.40	0 100 100	27, 42, 68, 69	0
All	All	334/365 (91%)	-0.39	0 100 100	13, 40, 86, 101	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PTR	A	1158	16/17	0.83	0.23	66,85,99,100	0
1	PTR	A	1162	16/17	0.95	0.14	43,54,59,60	0
1	PTR	A	1163	16/17	0.98	0.13	40,42,43,43	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	A	2	1/1	0.98	0.22	50,50,50,50	0
3	CA	A	1	1/1	0.99	0.36	55,55,55,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.