

Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2024 - 04:07 PM EDT

PDB ID : 9AU1

Title : SARS-CoV-2 XBB.1.5 RBD bound to the VIR-7229 and the S309 Fab frag-

ments

Authors: Rietz, T.; Park, Y.J.; Errico, J.; Czudnochowski, N.; Nix, J.C.; Corti, D.; Snell,

G.; Marco, A.D.; Pinto, D.; Cameroni, E.; Seattle Structural Genomics Center for Infectious Disease (SSGCID); Veesler, D.; Structural Genomics Consortium

(SGC)

Deposited on : 2024-02-27

Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.20.1

EDS : 3.0

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11


Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	164625	5670 (2.44-2.40)
Clashscore	180529	6299 (2.44-2.40)
Ramachandran outliers	177936	6232 (2.44-2.40)
Sidechain outliers	177891	6233 (2.44-2.40)
RSRZ outliers	164620	5670 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	M	230	91%	9%
2	N	212	89%	11%
3	Н	225	91%	7% •
4	L	216	83%	14% •



Continued from previous page...

Mol	Chain	Length	Quality of chain		
			4%		
5	\mathbf{R}	250	65%	12%	23%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	K	N	301	-	=	-	X



2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 8190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called S309 Fab heavy chain.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	М	230	Total	С	N	О	S	0	0	0
1	1V1	230	1714	1082	286	338	8	0	U	U

• Molecule 2 is a protein called S309 Fab light chain.

\mathbf{Mol}	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	N	212	Total 1606	C 998	N 271	O 332	S 5	0	0	0

• Molecule 3 is a protein called VIR-7229 Fab heavy chain.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
3	Н	219	Total 1668	C 1071	N 271	O 320	S 6	0	0	0

• Molecule 4 is a protein called VIR-7229 Fab light chain.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
4	L	212	Total 1589	C 998	N 262	O 324	S 5	0	0	0

• Molecule 5 is a protein called SARS-CoV-2 XBB.1.5 RBD.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
5	R	193	Total 1501	C 960	N 254	O 279	S 8	0	1	0

• Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	N	1	Total K 1 1	0	0




 \bullet Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	Н	1	Total C O 4 2 2	0	0
7	R	1	Total C O 4 2 2	0	0


 \bullet Molecule 8 is ACETIC ACID (three-letter code: ACY) (formula: $\mathrm{C_2H_4O_2}).$



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	L	1	Total 4	C 2	O 2	0	0




• Molecule 9 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	L	1	Total	С	N	O	S	0	0
	_	_	15	8	2	4	1		

• Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf		
10	R	1	Total 14	C 8	N 1	O 5	0	0



• Molecule 11 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	M	25	Total O 25 25	0	0
11	N	15	Total O 15 15	0	0
11	Н	12	Total O 12 12	0	0
11	L	9	Total O 9 9	0	0
11	R	9	Total O 9 9	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: S309 Fab heavy chain









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	73.73Å 112.53Å 170.31Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.98 - 2.41	Depositor
Resolution (A)	44.98 - 2.41	EDS
% Data completeness	99.9 (44.98-2.41)	Depositor
(in resolution range)	99.9 (44.98-2.41)	EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.25 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
D D.	0.227 , 0.257	Depositor
R, R_{free}	0.225 , 0.256	DCC
R_{free} test set	2760 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	60.6	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.28 , 29.0	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8190	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: K, ACY, EPE, PCA, EDO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5
1	M	0.45	0/1757	0.61	0/2395
2	N	0.47	0/1637	0.58	0/2224
3	Н	0.44	0/1712	0.57	0/2336
4	L	0.42	0/1624	0.55	0/2220
5	R	0.41	0/1547	0.56	0/2110
All	All	0.44	0/8277	0.57	0/11285

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	1714	0	1658	12	0
2	N	1606	0	1554	12	0
3	Н	1668	0	1626	9	0
4	L	1589	0	1520	18	0
5	R	1501	0	1405	15	0
6	N	1	0	0	0	0
7	Н	4	0	6	0	0
7	R	4	0	6	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	L	4	0	3	0	0
9	L	15	0	17	0	0
10	R	14	0	13	1	0
11	Η	12	0	0	0	1
11	L	9	0	0	0	0
11	M	25	0	0	1	1
11	N	15	0	0	0	0
11	R	9	0	0	0	0
All	All	8190	0	7808	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:19:THR:HG22	4:L:76:ILE:HG23	1.72	0.72
5:R:399:SER:HB3	5:R:511:VAL:HG12	1.75	0.68
5:R:384:PRO:HA	5:R:387:LEU:HD12	1.76	0.67
4:L:136:LEU:HD12	4:L:182:LEU:HD23	1.76	0.67
4:L:85:GLU:HG3	4:L:102:EEC:HD25	1.77	0.65
3:H:168:LEU:HD21	3:H:191:VAL:HG21	1.83	0.60
2:N:18:ARG:HG3	2:N:77:SER:HA	1.87	0.56
4:L:185:THR:HB	4:L:188:GLN:HG2	1.88	0.56
2:N:170:ASP:OD1	2:N:172:THR:OG1	2.18	0.54
1:M:131:LYS:HE3	1:M:158:ASP:O	2.16	0.54
1:M:131:LYS:NZ	11:M:301:HOH:O	2.40	0.53
4:L:26:SER:O	4:L:31:ALA:HB2	2.40	0.53
4:L:47:LYS:HD3	4:L:49:MET:SD	2.48	0.53
5:R:342:PHE:HB3	10:R:602:NAG:H82	1.93	0.50
3:H:104:LEU:HD22	5:R:459:SER:HA	1.93	0.50
5:R:342:PHE:CE1	5:R:511:VAL:HG21	2.46	0.50
4:L:137:VAL:HG12	4:L:139:LEU:HD13	1.96	0.30
4:L:136:LEU:HB2 3:H:34:MET:HB3	4:L:182:LEU:HB3 3:H:79:LEU:HD22	1.94	0.48
		1.95	0.48
5:R:425:LEU:HD21	5:R:512:VAL:HG11	1.95	0.48
2:N:140:TYR:CD1	2:N:141:PRO:HA	2.49	0.48
2:N:140:TYR:CG	2:N:141:PRO:HA	2.49	0.47
1:M:67:ARG:NH2	1:M:90:ASP:OD2	2.47	0.47
2:N:16:GLY:HA2	2:N:78:ARG:HG3	1.97	0.47
3:H:174:THR:HG23	3:H:189:SER:HB2	1.97	0.47



 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	Interatomic	Clash
	Atom-2	$\operatorname{distance}\ (ext{\AA})$	overlap (Å)
5:R:354:ASN:O	5:R:398:ASP:HA	2.15	0.47
4:L:23:THR:HG22	4:L:72:THR:HG23	1.97	0.46
3:H:97:ALA:HB3	3:H:110:LEU:HD23	1.98	0.46
2:N:183:LYS:HE2	2:N:187:GLU:OE2	2.15	0.46
5:R:526:GLY:O	5:R:528:LYS:N	2.48	0.46
2:N:122:ASP:O	2:N:126:LYS:HG3	2.15	0.46
5:R:438:SER:HB3	5:R:509:ARG:HG3	1.98	0.45
1:M:71:THR:OG1	1:M:80:TYR:HB2	2.17	0.45
5:R:503:VAL:HA	5:R:506:GLN:HG2	1.97	0.45
1:M:105:TRP:CD2	5:R:337:PRO:HG3	2.51	0.45
1:M:111:ILE:HG12	2:N:32:THR:HB	1.98	0.45
4:L:184:LEU:HD21	4:L:195:TYR:CZ	2.51	0.44
1:M:23:LYS:NZ	1:M:77:THR:OG1	2.47	0.44
5:R:398:ASP:O	5:R:511:VAL:HA	2.18	0.44
5:R:338:PHE:O	5:R:341:VAL:HG12	2.17	0.44
1:M:91:THR:HG23	1:M:124:THR:HA	2.00	0.44
2:N:163:VAL:HG22	2:N:175:LEU:HD12	2.00	0.43
3:H:223:LYS:NZ	4:L:126:SER:H	2.16	0.43
2:N:55:ARG:HG2	2:N:59:ILE:HB	2.01	0.43
4:L:124:PRO:HD3	4:L:136:LEU:CD2	2.49	0.43
4:L:184:LEU:HD23	4:L:189:TRP:HB2	2.01	0.43
4:L:185:THR:HG22	4:L:187:GLU:H	1.83	0.43
5:R:506:GLN:HB2	5:R:507:PRO:HD2	2.01	0.43
1:M:48:MET:HG2	1:M:64:PHE:CE2	2.54	0.42
4:L:142:ASP:O	4:L:175:LYS:HD3	2.19	0.42
1:M:32:TYR:CZ	1:M:98:ARG:NH2	2.87	0.42
1:M:67:ARG:HG2	1:M:85:ARG:CZ	2.49	0.42
2:N:186:TYR:HA	2:N:192:TYR:OH	2.19	0.42
3:H:104:LEU:HD23	5:R:460:LYS:HG3	2.02	0.41
4:L:137:VAL:HG13	4:L:181:TYR:CE2	2.55	0.41
2:N:62:ARG:NH1	2:N:83:ASP:OD1	2.52	0.41
3:H:12:VAL:HG21	3:H:86:LEU:HD13	2.02	0.41
3:H:144:THR:HG23	3:H:192:THR:HG23	2.02	0.41
4:L:37:TRP:CE2	4:L:75:LEU:HB2	2.56	0.41
1:M:23:LYS:NZ	1:M:76:THR:O	2.45	0.41
4:L:184:LEU:HG	4:L:188:GLN:HG3	2.02	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
11:M:322:HOH:O	11:H:412:HOH:O[3_555]	2.06	0.14

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	M	228/230 (99%)	219 (96%)	9 (4%)	0	100 100
2	N	210/212 (99%)	204 (97%)	6 (3%)	0	100 100
3	Н	215/225~(96%)	206 (96%)	9 (4%)	0	100 100
4	L	210/216 (97%)	202 (96%)	7 (3%)	1 (0%)	25 36
5	R	190/250 (76%)	177 (93%)	10 (5%)	3 (2%)	8 10
All	All	1053/1133 (93%)	1008 (96%)	41 (4%)	4 (0%)	30 42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	R	482	GLY
5	R	519	HIS
4	L	53	VAL
5	R	527	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Percentiles	
1	M	188/192 (98%)	186 (99%)	2 (1%)	70 84	



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Rotameric Outliers	
2	N	181/183 (99%)	177 (98%)	4 (2%)	47 65
3	Н	183/192 (95%)	182 (100%)	1 (0%)	86 94
4	L	177/181 (98%)	173 (98%)	4 (2%)	45 64
5	R	160/218 (73%)	154 (96%)	6 (4%)	28 45
All	All	889/966 (92%)	872 (98%)	17 (2%)	52 70

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Μ	211	ASN
1	M	229	SER
2	N	34	LEU
2	N	78	ARG
2	N	90	GLN
2	N	93	ASP
3	Н	50	TYR
4	L	28	ASP
4	L	108	THR
4	L	167	THR
4	L	179	SER
5	R	333	THR
5	R	393	THR
5	R	458	LYS
5	R	510	VAL
5	R	516	GLU
5	R	517	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
4	L	188	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			В	ond ang	gles
					Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	PCA	L	1	4	7,8,9	2.50	1 (14%)	9,10,12	1.33	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PCA	L	1	4	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
4	L	1	PCA	CD-N	5.69	1.48	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
4	L	1	PCA	CB-CG-CD	2.89	108.88	104.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Вс	Bond lengths			Bond angles		
MIOI	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
9	EPE	L	302	-	15,15,15	1.05	1 (6%)	19,20,20	1.85	6 (31%)	
8	ACY	L	301	-	3,3,3	1.22	0	3,3,3	1.50	0	
10	NAG	R	602	5	14,14,15	0.54	0	17,19,21	0.63	1 (5%)	
7	EDO	R	601	-	3,3,3	0.46	0	2,2,2	0.30	0	
7	EDO	Н	301	-	3,3,3	0.62	0	2,2,2	0.06	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	R	602	5	-	0/6/23/26	0/1/1/1
9	EPE	L	302	-	-	6/9/19/19	0/1/1/1
7	EDO	Н	301	-	-	1/1/1/1	-
7	EDO	R	601	_	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
9	L	302	EPE	C10-S	3.48	1.82	1.77

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathbf{Ideal}(^o)$
9	L	302	EPE	C5-N4-C3	3.99	117.43	108.84
9	L	302	EPE	C7-N4-C5	3.37	120.23	111.24
9	L	302	EPE	O3S-S-C10	3.27	112.40	106.00
9	L	302	EPE	O1S-S-C10	2.81	110.97	106.73
9	L	302	EPE	C7-N4-C3	2.23	117.19	111.24



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
10	R	602	NAG	C1-O5-C5	2.08	114.97	112.19
9	L	302	EPE	O3S-S-O1S	-2.06	106.24	111.40

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	L	302	EPE	C9-C10-S-O1S
9	L	302	EPE	C9-C10-S-O3S
9	L	302	EPE	C9-C10-S-O2S
9	L	302	EPE	S-C10-C9-N1
9	L	302	EPE	C10-C9-N1-C2
9	L	302	EPE	C10-C9-N1-C6
7	Н	301	EDO	O1-C1-C2-O2
7	R	601	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	R	602	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2		$OWAB(Å^2)$	Q<0.9	
1	M	$230/230\ (100\%)$	-0.10	1 (0%)	89	87	54, 69, 89, 138	0
2	N	212/212 (100%)	-0.11	1 (0%)	87	86	56, 72, 93, 123	0
3	Н	219/225~(97%)	-0.01	2 (0%)	81	79	63, 73, 114, 136	0
4	L	211/216 (97%)	-0.01	1 (0%)	87	86	66, 79, 97, 125	0
5	R	193/250 (77%)	0.29	10 (5%)	34	31	51, 76, 120, 139	1 (0%)
All	All	1065/1133 (93%)	0.01	15 (1%)	73	71	51, 73, 104, 139	1 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	R	374	PHE	4.3
5	R	369	TYR	3.9
5	R	370	ASN	3.8
5	R	368	ILE	3.4
5	R	528	LYS	3.2
2	N	32	THR	3.1
3	Н	136	SER	2.9
5	R	375	PHE	2.9
5	R	436	TRP	2.6
5	R	367	VAL	2.5
1	M	230	CYS	2.5
5	R	376	ALA	2.4
5	R	377	PHE	2.2
3	Н	123	ASN	2.1
4	L	3	VAL	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	PCA	L	1	8/9	0.69	0.19	105,111,116,117	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
6	K	N	301	1/1	0.37	0.46	205,205,205,205	0
8	ACY	L	301	4/4	0.79	0.15	88,90,90,91	0
10	NAG	R	602	14/15	0.79	0.17	85,90,98,105	14
7	EDO	R	601	4/4	0.80	0.23	93,93,94,96	0
7	EDO	Н	301	4/4	0.85	0.15	63,64,65,66	0
9	EPE	L	302	15/15	0.93	0.12	72,75,82,90	0

6.5 Other polymers (i)

There are no such residues in this entry.

