



Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 04:16 pm BST

PDB ID : 3AXF
Title : Perrhenate binding to A11C/R153C ModA mutant
Authors : He, C.; Aryal, B.P.; Brugarolas, P.
Deposited on : 2011-04-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	
1	B	237	
1	C	237	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Molybdate-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	231	Total 1804	C 1153	N 302	O 345	S 4	0	8	0
1	B	231	Total 1742	C 1111	N 291	O 336	S 4	0	1	0
1	C	231	Total 1785	C 1136	N 299	O 345	S 5	0	7	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP P37329
A	-2	SER	-	EXPRESSION TAG	UNP P37329
A	-1	HIS	-	EXPRESSION TAG	UNP P37329
A	0	MET	-	EXPRESSION TAG	UNP P37329
A	11	CYS	ALA	ENGINEERED MUTATION	UNP P37329
A	153	CYS	ARG	ENGINEERED MUTATION	UNP P37329
B	-3	GLY	-	EXPRESSION TAG	UNP P37329
B	-2	SER	-	EXPRESSION TAG	UNP P37329
B	-1	HIS	-	EXPRESSION TAG	UNP P37329
B	0	MET	-	EXPRESSION TAG	UNP P37329
B	11	CYS	ALA	ENGINEERED MUTATION	UNP P37329
B	153	CYS	ARG	ENGINEERED MUTATION	UNP P37329
C	-3	GLY	-	EXPRESSION TAG	UNP P37329
C	-2	SER	-	EXPRESSION TAG	UNP P37329
C	-1	HIS	-	EXPRESSION TAG	UNP P37329
C	0	MET	-	EXPRESSION TAG	UNP P37329
C	11	CYS	ALA	ENGINEERED MUTATION	UNP P37329
C	153	CYS	ARG	ENGINEERED MUTATION	UNP P37329

- Molecule 2 is PERRHENATE (three-letter code: REO) (formula: O₄Re).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	Re	0	0
			5	4	1		
2	B	1	Total	O	Re	0	0
			5	4	1		
2	C	1	Total	O	Re	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	56	Total	O	0	0
			56	56		
3	B	90	Total	O	0	0
			90	90		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Molybdate-binding periplasmic protein



- Molecule 1: Molybdate-binding periplasmic protein



- Molecule 1: Molybdate-binding periplasmic protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.75Å 77.46Å 163.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 40.76 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.00) 100.0 (40.76-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.204 , 0.258 0.207 , 0.259	Depositor DCC
R_{free} test set	2322 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtrriage
Anisotropy	0.035	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5492	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: REO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.22	8/1845 (0.4%)	0.95	3/2497 (0.1%)
1	B	1.28	5/1779 (0.3%)	1.00	5/2414 (0.2%)
1	C	0.79	0/1826	0.78	0/2479
All	All	1.12	13/5450 (0.2%)	0.91	8/7390 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	11	CYS	CB-SG	-22.15	1.44	1.82
1	B	188	GLU	CG-CD	7.11	1.62	1.51
1	A	96[A]	LYS	CE-NZ	6.58	1.65	1.49
1	A	96[B]	LYS	CE-NZ	6.58	1.65	1.49
1	B	37	ALA	CA-CB	6.33	1.65	1.52
1	B	188	GLU	CD-OE2	6.15	1.32	1.25
1	A	198	VAL	CB-CG1	5.74	1.64	1.52
1	A	55	PHE	CE2-CZ	5.46	1.47	1.37
1	B	11	CYS	CB-SG	-5.43	1.73	1.81
1	A	37	ALA	CA-CB	5.35	1.63	1.52
1	A	202	GLU	CD-OE2	-5.25	1.19	1.25
1	A	75	ALA	CA-CB	5.07	1.63	1.52
1	B	84	SER	CA-CB	5.04	1.60	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	115	LEU	CB-CG-CD2	7.02	122.93	111.00
1	A	44	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	B	101	ASP	CB-CG-OD1	5.63	123.36	118.30
1	A	11	CYS	CB-CA-C	-5.41	99.58	110.40
1	B	217	LYS	CD-CE-NZ	5.38	124.07	111.70
1	A	114	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	B	109	LEU	CB-CG-CD1	5.10	119.67	111.00
1	B	151	ASP	CB-CG-OD1	5.04	122.84	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	232	ILE	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1804	0	1820	25	0
1	B	1742	0	1755	10	0
1	C	1785	0	1792	37	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	56	0	0	1	0
3	B	90	0	0	2	0
All	All	5492	0	5367	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:204:HIS:O	1:C:205[B]:ASN:HB2	1.74	0.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:53:ASP:OD1	1:B:204:HIS:HD2	1.61	0.82
1:B:74:THR:HG21	3:B:482:HOH:O	1.80	0.80
1:A:121:GLU:O	1:A:122[B]:HIS:CG	2.38	0.76
1:C:206:ASN:OD1	1:C:209:VAL:HG23	1.84	0.76
1:C:227:ARG:HH11	1:C:227:ARG:HB3	1.53	0.73
1:A:53:ASP:OD1	1:A:204:HIS:HD2	1.71	0.73
1:C:204:HIS:O	1:C:205[B]:ASN:CB	2.37	0.73
1:C:8:PHE:CD1	1:C:52:ALA:HB2	2.24	0.71
1:A:193:LYS:HE2	1:A:233[B]:LYS:HE3	1.73	0.69
1:A:156:LEU:HD11	1:A:180:VAL:HG21	1.74	0.69
1:A:204:HIS:HE1	3:A:449:HOH:O	1.78	0.67
1:C:105:ASN:O	1:C:109:LEU:HG	1.97	0.65
1:C:11[B]:CYS:SG	1:C:37:ALA:HA	2.37	0.64
1:B:132:ALA:HA	1:B:190:SER:O	1.98	0.63
1:A:94:VAL:O	1:A:96[A]:LYS:HD3	1.98	0.63
1:C:6:THR:HB	1:C:52:ALA:HA	1.81	0.62
1:B:53:ASP:OD1	1:B:204:HIS:CD2	2.50	0.61
1:A:24:PHE:CE1	1:A:28:LYS:HG3	2.37	0.60
1:A:93:SER:O	1:A:181:LYS:NZ	2.27	0.59
1:B:231:THR:HG23	3:B:463:HOH:O	2.03	0.58
1:C:86:VAL:O	1:C:168:ILE:HA	2.08	0.54
1:B:96:LYS:HE2	1:C:189:ASP:HB3	1.89	0.54
1:C:88:VAL:HG11	1:C:174:ALA:HB2	1.88	0.54
1:A:178:LYS:HD3	1:A:178:LYS:H	1.74	0.52
1:B:106:TRP:HB3	1:B:142:LEU:HD11	1.91	0.52
1:A:91[B]:LYS:HG3	1:A:92:ALA:N	2.23	0.52
1:B:114:ARG:O	1:B:165:PRO:HD2	2.11	0.50
1:C:115:LEU:HD22	1:C:167:GLY:HA2	1.94	0.49
1:C:14:THR:HA	1:C:36:PHE:CE1	2.47	0.49
1:A:193:LYS:HE2	1:A:233[B]:LYS:CE	2.40	0.49
1:C:11[B]:CYS:SG	1:C:37:ALA:CA	3.00	0.49
1:C:11[B]:CYS:HB3	1:C:153:CYS:SG	2.52	0.48
1:B:210:LYS:HD2	1:B:210:LYS:HA	1.64	0.48
1:C:121:GLU:HA	1:C:121:GLU:OE2	2.14	0.47
1:A:178:LYS:CD	1:A:178:LYS:N	2.78	0.47
1:C:100:ILE:HG21	1:C:186:PHE:CD2	2.51	0.46
1:C:32:VAL:HG12	1:C:33:VAL:N	2.30	0.46
1:B:24:PHE:CE1	1:B:28:LYS:HG3	2.50	0.46
1:A:210:LYS:HD2	1:A:210:LYS:HA	1.73	0.46
1:A:15:ASN:HB3	1:A:228:TYR:CE1	2.50	0.46
1:C:225:PHE:O	1:C:230:PHE:HD2	1.99	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:LEU:HD13	1:A:142:LEU:HD23	1.98	0.45
1:C:119[B]:ASP:OD1	1:C:148:PRO:HB2	2.16	0.45
1:C:172:SER:HB2	1:C:228:TYR:O	2.17	0.45
1:A:53:ASP:OD1	1:A:204:HIS:CD2	2.61	0.44
1:C:13:LEU:O	1:C:17:MET:HB2	2.16	0.44
1:C:83:ASN:HD21	1:C:172:SER:CB	2.29	0.44
1:A:121:GLU:O	1:A:122[B]:HIS:CD2	2.71	0.44
1:C:132:ALA:HA	1:C:190:SER:O	2.18	0.44
1:C:83:ASN:O	1:C:194:VAL:HB	2.17	0.44
1:C:227:ARG:NH1	1:C:227:ARG:HB3	2.28	0.43
1:C:5:ILE:O	1:C:33:VAL:HG23	2.18	0.43
1:C:28:LYS:HD2	1:C:28:LYS:O	2.19	0.43
1:C:143:SER:N	1:C:144:PRO:HD2	2.34	0.42
1:A:193:LYS:CE	1:A:233[B]:LYS:HE3	2.44	0.42
1:C:52:ALA:O	1:C:201:VAL:HG21	2.20	0.42
1:C:8:PHE:HB2	1:C:46:ILE:HD11	2.02	0.42
1:C:93:SER:O	1:C:181:LYS:NZ	2.49	0.42
1:A:23:GLN:HG2	1:A:215:TYR:OH	2.19	0.41
1:C:9:ALA:HB1	1:C:13:LEU:HD12	2.02	0.41
1:C:83:ASN:HB3	1:C:194:VAL:HG11	2.01	0.41
1:A:121:GLU:C	1:A:122[B]:HIS:CG	2.93	0.41
1:C:155:ALA:O	1:C:158:LEU:HB2	2.21	0.41
1:A:178:LYS:HD3	1:A:178:LYS:N	2.35	0.41
1:C:37:ALA:O	1:C:38:SER:C	2.58	0.41
1:A:40:SER:HB3	1:A:62:TRP:CE2	2.56	0.40
1:C:34[B]:SER:HB2	1:C:36:PHE:CZ	2.56	0.40
1:A:79:THR:HG22	1:A:233[A]:LYS:HG3	2.04	0.40
1:A:110:LEU:HD23	1:A:110:LEU:HA	1.95	0.40
1:A:9:ALA:O	1:A:36:PHE:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/237 (100%)	229 (97%)	7 (3%)	0	100	100
1	B	230/237 (97%)	228 (99%)	2 (1%)	0	100	100
1	C	236/237 (100%)	214 (91%)	17 (7%)	5 (2%)	7	2
All	All	702/711 (99%)	671 (96%)	26 (4%)	5 (1%)	25	16

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	205[A]	ASN
1	C	205[B]	ASN
1	C	10	ALA
1	C	97	ASP
1	C	51	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/187 (102%)	172 (90%)	18 (10%)	8	5
1	B	183/187 (98%)	174 (95%)	9 (5%)	25	21
1	C	189/187 (101%)	164 (87%)	25 (13%)	4	2
All	All	562/561 (100%)	510 (91%)	52 (9%)	10	5

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LYS
1	A	26	LYS
1	A	54[A]	LEU
1	A	54[B]	LEU
1	A	74	THR
1	A	77	ARG
1	A	96[A]	LYS
1	A	96[B]	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	97	ASP
1	A	115	LEU
1	A	135	LYS
1	A	142	LEU
1	A	158	LEU
1	A	178	LYS
1	A	210	LYS
1	A	223	GLU
1	A	233[A]	LYS
1	A	233[B]	LYS
1	B	54	LEU
1	B	74	THR
1	B	91	LYS
1	B	96	LYS
1	B	109	LEU
1	B	115	LEU
1	B	142	LEU
1	B	158	LEU
1	B	210	LYS
1	C	6	THR
1	C	11[A]	CYS
1	C	11[B]	CYS
1	C	15	ASN
1	C	28	LYS
1	C	30	VAL
1	C	31	ASP
1	C	33	VAL
1	C	41	THR
1	C	45	GLN
1	C	54[A]	LEU
1	C	54[B]	LEU
1	C	57	SER
1	C	91	LYS
1	C	96	LYS
1	C	99	THR
1	C	115	LEU
1	C	135	LYS
1	C	158	LEU
1	C	178	LYS
1	C	193	LYS
1	C	206	ASN
1	C	210	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	226	LYS
1	C	227	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	204	HIS
1	B	15	ASN
1	B	204	HIS
1	C	23	GLN
1	C	45	GLN
1	C	95	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	REO	C	301	-	3,4,4	2.98	3 (100%)	-		
2	REO	A	301	-	3,4,4	3.56	2 (66%)	-		
2	REO	B	301	-	3,4,4	2.36	2 (66%)	-		

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	REO	RE-O13	4.72	1.91	1.70
2	A	301	REO	RE-O14	3.60	1.86	1.70
2	B	301	REO	RE-O13	3.54	1.86	1.70
2	C	301	REO	RE-O14	3.22	1.84	1.70
2	C	301	REO	RE-O12	2.96	1.83	1.70
2	C	301	REO	RE-O13	2.73	1.82	1.70
2	B	301	REO	RE-O12	2.00	1.79	1.70

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/237 (97%)	0.19	6 (2%) 56 54	19, 36, 57, 67	1 (0%)
1	B	231/237 (97%)	-0.02	1 (0%) 92 92	18, 26, 40, 52	0
1	C	231/237 (97%)	1.81	84 (36%) 0 0	45, 67, 93, 117	0
All	All	693/711 (97%)	0.66	91 (13%) 3 3	18, 39, 81, 117	1 (0%)

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	207	ALA	7.2
1	C	30	VAL	7.0
1	C	99	THR	7.0
1	C	51	PRO	6.5
1	C	183	VAL	6.4
1	C	98	PHE	6.0
1	C	32	VAL	5.6
1	C	97	ASP	5.5
1	C	100	ILE	5.3
1	C	175	VAL	5.1
1	C	122[A]	HIS	5.0
1	C	5	ILE	4.9
1	C	20	ILE	4.9
1	C	48	ALA	4.8
1	C	87	VAL	4.6
1	C	106	TRP	4.5
1	C	184	ALA	4.4
1	C	104	THR	4.3
1	A	233[A]	LYS	4.2
1	C	203	GLY	4.0
1	C	213	TYR	4.0
1	C	94	VAL	3.9
1	C	4	LYS	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	133	LEU	3.8
1	C	172	SER	3.8
1	C	10	ALA	3.8
1	A	171	GLY	3.8
1	C	222	ALA	3.7
1	A	122[A]	HIS	3.6
1	C	92	ALA	3.5
1	C	144	PRO	3.4
1	C	139	TRP	3.3
1	C	171	GLY	3.3
1	C	52	ALA	3.3
1	C	229	GLY	3.3
1	C	102	SER	3.3
1	C	119[A]	ASP	3.2
1	A	92	ALA	3.2
1	C	53	ASP	3.2
1	C	89	ALA	3.1
1	C	152	VAL	3.1
1	C	178	LYS	3.0
1	C	34[A]	SER	3.0
1	C	176	ALA	3.0
1	C	170	TYR	2.9
1	C	117	VAL	2.9
1	C	137	GLY	2.9
1	C	95	GLN	2.9
1	C	230	PHE	2.9
1	C	33	VAL	2.8
1	C	46	ILE	2.8
1	C	205[A]	ASN	2.8
1	C	124	PRO	2.8
1	C	156	LEU	2.8
1	C	83	ASN	2.8
1	C	208	THR	2.8
1	C	142	LEU	2.8
1	C	11[A]	CYS	2.8
1	C	37	ALA	2.7
1	C	91	LYS	2.7
1	C	88	VAL	2.6
1	B	3	GLY	2.6
1	C	7	VAL	2.6
1	C	201	VAL	2.6
1	C	13	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	93	SER	2.5
1	C	35[A]	SER	2.4
1	C	143	SER	2.4
1	C	174	ALA	2.4
1	C	212	PHE	2.4
1	C	43	ALA	2.4
1	C	96	LYS	2.3
1	A	176	ALA	2.3
1	C	45	GLN	2.3
1	C	202	GLU	2.3
1	C	121	GLU	2.3
1	C	232	ILE	2.3
1	C	206	ASN	2.2
1	C	16	ALA	2.2
1	C	42	LEU	2.2
1	C	109	LEU	2.2
1	C	140	ASP	2.2
1	C	167	GLY	2.2
1	C	8	PHE	2.1
1	C	22	THR	2.1
1	C	116	ALA	2.0
1	C	84	SER	2.0
1	C	54[A]	LEU	2.0
1	A	178	LYS	2.0
1	C	224	ILE	2.0
1	C	3	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	REO	C	301	5/5	0.99	0.12	69,69,71,74	0
2	REO	A	301	5/5	1.00	0.13	19,20,30,36	0
2	REO	B	301	5/5	1.00	0.14	18,20,24,26	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around REO A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.