



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 25, 2023 – 08:34 PM EDT

PDB ID : 3AZF
Title : Crystal Structure of Human Nucleosome Core Particle Containing H3K79Q mutation
Authors : Iwasaki, W.; Tachiwana, H.; Kawaguchi, K.; Shibata, T.; Kagawa, W.; Kurumizaka, H.
Deposited on : 2011-05-25
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	
1	E	139	
2	B	106	
2	F	106	
3	C	133	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	133	<p>1% 59% 17% 22%</p>
4	D	129	<p>2% 44% 26% 27%</p>
4	H	129	<p>0% 55% 14% 29%</p>
5	I	146	<p>4% 14% 86% 0%</p>
5	J	146	<p>3% 14% 86% 0%</p>

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12152 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	97	801	504	155	138	4	0	0	0
1	E	99	816	513	158	141	4	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P68431
A	-2	SER	-	expression tag	UNP P68431
A	-1	HIS	-	expression tag	UNP P68431
A	79	GLN	LYS	engineered mutation	UNP P68431
E	-3	GLY	-	expression tag	UNP P68431
E	-2	SER	-	expression tag	UNP P68431
E	-1	HIS	-	expression tag	UNP P68431
E	79	GLN	LYS	engineered mutation	UNP P68431

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	78	619	391	120	107	1	0	0	0
2	F	84	673	424	133	115	1	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P62805
B	-2	SER	-	expression tag	UNP P62805
B	-1	HIS	-	expression tag	UNP P62805
F	-3	GLY	-	expression tag	UNP P62805

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-2	SER	-	expression tag	UNP P62805
F	-1	HIS	-	expression tag	UNP P62805

- Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	108	835	526	165	144	0	0	0
3	G	104	805	508	157	140	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P04908
C	-2	SER	-	expression tag	UNP P04908
C	-1	HIS	-	expression tag	UNP P04908
G	-3	GLY	-	expression tag	UNP P04908
G	-2	SER	-	expression tag	UNP P04908
G	-1	HIS	-	expression tag	UNP P04908

- Molecule 4 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	94	736	462	134	138	2	0	0	0
4	H	91	714	450	128	134	2	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	expression tag	UNP P06899
D	-2	SER	-	expression tag	UNP P06899
D	-1	HIS	-	expression tag	UNP P06899
H	-3	GLY	-	expression tag	UNP P06899
H	-2	SER	-	expression tag	UNP P06899
H	-1	HIS	-	expression tag	UNP P06899

- Molecule 5 is a DNA chain called 146-MER DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	145	Total	C	N	O	P	0	0	0
			2970	1421	538	867	144			
5	J	145	Total	C	N	O	P	0	0	0
			2969	1421	535	869	144			

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		
6	C	1	Total	Cl	0	0
			1	1		
6	E	1	Total	Cl	0	0
			1	1		
6	G	1	Total	Cl	0	0
			1	1		

- Molecule 7 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	1	Total	Mn	0	0
			1	1		
7	I	6	Total	Mn	0	0
			6	6		
7	J	5	Total	Mn	0	0
			5	5		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	22	Total	O	0	0
			22	22		
8	B	19	Total	O	0	0
			19	19		
8	C	29	Total	O	0	0
			29	29		
8	D	13	Total	O	0	0
			13	13		
8	E	40	Total	O	0	0
			40	40		
8	F	28	Total	O	0	0
			28	28		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	14	Total 14	O 14	0	0
8	H	15	Total 15	O 15	0	0
8	I	8	Total 8	O 8	0	0
8	J	10	Total 10	O 10	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3.1

Chain A: 




- Molecule 1: Histone H3.1

Chain E: 



- Molecule 2: Histone H4

Chain B: 




- Molecule 2: Histone H4


Chain F: 





• Molecule 3: Histone H2A type 1-B/E




• Molecule 3: Histone H2A type 1-B/E




• Molecule 4: Histone H2B type 1-J



• Molecule 4: Histone H2B type 1-J



• Molecule 5: 146-MER DNA



G63 T64
T65 G66 A67
G71 A72
A77 G78
C79 T80
G81 A82
A83 G84
A85 T86
G87 C88
C89 T90
T91 T92
T93 G94
A95 T96
G97 G98
A99 G100
C101 A102
G103 T104
T105 T106
C107 C108
A109 A110
A111 T112
A113 C114
A115 C116
T117 T118
T119 T120
G121 G122
T123 A124
G125 A127

T128 C129
T130 G131
A132 A133
G134 G135
T136 G137
C138 A139
T140 A141
T142 T143
G144 A145
DT

● Molecule 5: 146-MER DNA

Chain J: 3% 14% 86%

DA T148
C149 A150
A151 T152
A153 T154
C155 A156
A157 C158
C159 G162
A165 T166
T167 C168
T169 A170
C171 T172
A173 A174
A175 A176
G177 T178
G179 T180
A181 T182
T183 T184
G185 G186
A187 A188
A189 C190
T191 G192
C193 T194
C195 C196
A197 T198
C199 A200
A201 A202
A203 G204
A207 T208
G209

T210 T211
C212 G215
T216 G217
A218 A219
T220 C221
T221 C222
G223 G224
C225 T226
G227 A228
A229 C230
A231 T232
G233 C234
C235 T236
T237 T238
T239 G240
A241 T242
G243 G244
A245 G246
C247 A248
G249 T250
T251 T252
C253 C254
A255 A256
A257 T258
A259 C260
A261 C262
T263 T264
T265 T266
G267 G268
T269 A270

G271 A272
A273 T274
C275 T276
G277 C278
A279 G280
G281 T282
G283 G284
A285 T286
A287 T288
T289 G290
A291 T292

4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.55Å 109.78Å 182.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.80 – 2.70 48.80 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.80-2.70) 99.6 (48.80-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.99 (at 2.69Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.213 , 0.263 0.212 , 0.264	Depositor DCC
R_{free} test set	3000 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtrriage
Anisotropy	0.326	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 60.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.014 for k,h,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12152	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/813	0.63	0/1091
1	E	0.48	0/828	0.66	0/1110
2	B	0.42	0/626	0.65	0/837
2	F	0.46	0/680	0.69	0/908
3	C	0.40	0/845	0.63	0/1139
3	G	0.38	0/815	0.61	0/1100
4	D	0.41	0/747	0.61	0/1004
4	H	0.43	0/725	0.60	0/975
5	I	0.39	0/3332	0.78	0/5141
5	J	0.38	0/3330	0.79	0/5138
All	All	0.41	0/12741	0.72	0/18443

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	834	34	0
1	E	816	0	851	33	0
2	B	619	0	659	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	673	0	722	20	0
3	C	835	0	897	50	0
3	G	805	0	861	26	0
4	D	736	0	758	40	0
4	H	714	0	735	27	0
5	I	2970	0	1640	235	0
5	J	2969	0	1641	198	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
6	E	1	0	0	1	0
6	G	1	0	0	0	0
7	D	1	0	0	0	0
7	I	6	0	0	0	0
7	J	5	0	0	0	0
8	A	22	0	0	2	0
8	B	19	0	0	0	0
8	C	29	0	0	4	0
8	D	13	0	0	0	0
8	E	40	0	0	0	0
8	F	28	0	0	2	0
8	G	14	0	0	2	0
8	H	15	0	0	2	0
8	I	8	0	0	0	0
8	J	10	0	0	0	0
All	All	12152	0	9598	596	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 596 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:J:242:DT:H2''	5:J:243:DG:H5'	1.25	1.16
5:I:9:DC:H2''	5:I:10:DC:H5'	1.29	1.11
5:J:197:DA:H2''	5:J:198:DT:H5'	1.27	1.11
5:J:231:DA:H2''	5:J:232:DT:H5'	1.28	1.10
5:J:190:DC:H2''	5:J:191:DT:H5''	1.32	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/139 (68%)	93 (98%)	2 (2%)	0	100	100
1	E	97/139 (70%)	94 (97%)	2 (2%)	1 (1%)	15	37
2	B	76/106 (72%)	74 (97%)	2 (3%)	0	100	100
2	F	82/106 (77%)	80 (98%)	2 (2%)	0	100	100
3	C	106/133 (80%)	101 (95%)	5 (5%)	0	100	100
3	G	102/133 (77%)	98 (96%)	3 (3%)	1 (1%)	15	37
4	D	92/129 (71%)	89 (97%)	2 (2%)	1 (1%)	14	34
4	H	89/129 (69%)	86 (97%)	2 (2%)	1 (1%)	14	34
All	All	739/1014 (73%)	715 (97%)	20 (3%)	4 (0%)	29	54

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	133	GLU
4	H	104	GLY
4	D	104	GLY
3	G	26	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/113 (75%)	82 (96%)	3 (4%)	36	65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	86/113 (76%)	82 (95%)	4 (5%)	26	54
2	B	63/81 (78%)	61 (97%)	2 (3%)	39	68
2	F	69/81 (85%)	67 (97%)	2 (3%)	42	71
3	C	85/102 (83%)	79 (93%)	6 (7%)	14	34
3	G	83/102 (81%)	78 (94%)	5 (6%)	19	42
4	D	80/107 (75%)	73 (91%)	7 (9%)	10	23
4	H	78/107 (73%)	76 (97%)	2 (3%)	46	75
All	All	629/806 (78%)	598 (95%)	31 (5%)	25	52

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	88	THR
3	G	114	VAL
4	D	112	SER
4	H	88	THR
3	G	73	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	113	HIS
3	G	31	HIS
4	H	63	ASN
3	G	73	ASN
3	C	73	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	97/139 (69%)	-0.17	0 100 100	26, 39, 64, 96	0
1	E	99/139 (71%)	-0.06	1 (1%) 82 83	16, 32, 68, 86	0
2	B	78/106 (73%)	-0.16	1 (1%) 77 78	27, 38, 60, 80	0
2	F	84/106 (79%)	0.02	2 (2%) 59 60	21, 32, 56, 101	0
3	C	108/133 (81%)	0.00	4 (3%) 41 41	21, 38, 75, 125	0
3	G	104/133 (78%)	-0.18	1 (0%) 82 83	26, 43, 72, 98	0
4	D	94/129 (72%)	-0.01	2 (2%) 63 65	22, 40, 75, 105	0
4	H	91/129 (70%)	-0.10	0 100 100	28, 42, 67, 87	0
5	I	145/146 (99%)	0.23	6 (4%) 37 36	43, 99, 141, 165	0
5	J	145/146 (99%)	0.18	4 (2%) 53 54	46, 101, 145, 160	0
All	All	1045/1306 (80%)	-0.00	21 (2%) 65 67	16, 44, 129, 165	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	11	ARG	7.0
3	C	13	LYS	5.0
2	F	102	GLY	4.9
3	C	12	ALA	4.4
1	E	37	LYS	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	MN	J	1001	1/1	0.54	0.11	112,112,112,112	0
7	MN	I	1005	1/1	0.60	0.10	114,114,114,114	0
7	MN	I	1001	1/1	0.61	0.15	125,125,125,125	0
7	MN	I	1002	1/1	0.66	0.09	116,116,116,116	0
7	MN	I	1006	1/1	0.69	0.18	119,119,119,119	0
7	MN	I	1003	1/1	0.85	0.26	85,85,85,85	0
7	MN	I	1004	1/1	0.92	0.15	85,85,85,85	0
7	MN	J	1004	1/1	0.93	0.18	71,71,71,71	0
7	MN	J	1002	1/1	0.95	0.16	87,87,87,87	0
6	CL	E	1001	1/1	0.96	0.13	51,51,51,51	0
6	CL	A	1001	1/1	0.96	0.10	53,53,53,53	0
7	MN	J	1005	1/1	0.96	0.07	121,121,121,121	0
6	CL	G	1001	1/1	0.97	0.15	49,49,49,49	0
6	CL	C	1001	1/1	0.98	0.14	45,45,45,45	0
7	MN	J	1003	1/1	0.98	0.13	72,72,72,72	0
7	MN	D	201	1/1	0.99	0.23	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.