



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 26, 2022 – 11:11 AM EST

PDB ID : 2B1O  
Title : Solution Structure of Ca<sup>2+</sup>-bound DdCAD-1  
Authors : Lin, Z.; Sriskanthadevan, S.; Huang, H.B.; Siu, C.H.; Yang, D.W.  
Deposited on : 2005-09-16

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.27  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	212	86% 11% .

## 2 Ensemble composition and analysis

This entry contains 10 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:31, A:37-A:213 (206)	0.56	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 8, 9, 10
Single-model clusters	7

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3040 atoms, of which 1361 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Calcium-dependent cell adhesion molecule-1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	212	3037	1061	1361	276	332	7	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	
			Total	Ca
2	A	3	3	3

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Calcium-dependent cell adhesion molecule-1




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: Calcium-dependent cell adhesion molecule-1



#### 4.2.2 Score per residue for model 2 (medoid)

- Molecule 1: Calcium-dependent cell adhesion molecule-1







### 4.2.3 Score per residue for model 3

- Molecule 1: Calcium-dependent cell adhesion molecule-1


Chain A: 75% 21% ..



### 4.2.4 Score per residue for model 4

- Molecule 1: Calcium-dependent cell adhesion molecule-1


Chain A: 78% 17% ..



### 4.2.5 Score per residue for model 5

- Molecule 1: Calcium-dependent cell adhesion molecule-1


Chain A: 81% 16% .



### 4.2.6 Score per residue for model 6


- Molecule 1: Calcium-dependent cell adhesion molecule-1

Chain A: 76% 20% ..




#### 4.2.7 Score per residue for model 7

- Molecule 1: Calcium-dependent cell adhesion molecule-1




#### 4.2.8 Score per residue for model 8

- Molecule 1: Calcium-dependent cell adhesion molecule-1




#### 4.2.9 Score per residue for model 9

- Molecule 1: Calcium-dependent cell adhesion molecule-1



#### 4.2.10 Score per residue for model 10

- Molecule 1: Calcium-dependent cell adhesion molecule-1





## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.5
Amber	refinement	7.0

No chemical shift data was provided.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.71±0.01	0±0/1673 ( 0.0± 0.0%)	1.02±0.02	1±1/2275 ( 0.1± 0.1%)
All	All	0.71	0/16730 ( 0.0%)	1.02	13/22750 ( 0.1%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	59	GLU	OE1-CD-OE2	-6.21	115.85	123.30	2	4
1	A	51	ARG	NE-CZ-NH1	6.08	123.34	120.30	10	3
1	A	56	GLU	OE1-CD-OE2	-5.34	116.89	123.30	3	3
1	A	124	VAL	CA-CB-CG1	5.25	118.77	110.90	10	1
1	A	157	ARG	NE-CZ-NH1	5.15	122.87	120.30	1	2

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1633	1332	1577	2±1
All	All	16360	13320	15770	17

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:92:LEU:HD13	1:A:96:PHE:CE1	0.64	2.27	10	1
1:A:92:LEU:HD13	1:A:96:PHE:CE2	0.50	2.41	4	1
1:A:117:MET:HG3	1:A:156:VAL:HG22	0.46	1.86	10	1
1:A:181:GLN:HG2	1:A:182:VAL:H	0.46	1.70	3	2
1:A:3:VAL:HG22	1:A:4:ASP:H	0.46	1.70	4	1
1:A:80:LEU:HD13	1:A:80:LEU:H	0.45	1.71	2	1
1:A:116:GLU:O	1:A:156:VAL:HG23	0.45	2.11	9	1
1:A:191:PHE:CD1	1:A:192:PRO:HD2	0.44	2.47	3	1
1:A:115:TYR:CZ	1:A:156:VAL:HG21	0.44	2.48	1	1
1:A:93:PRO:CG	1:A:96:PHE:CD2	0.43	3.01	8	1
1:A:184:VAL:HG13	1:A:205:PHE:CZ	0.42	2.49	2	1
1:A:156:VAL:HG22	1:A:166:VAL:CG2	0.42	2.44	9	1
1:A:54:ILE:HB	1:A:86:LEU:HD13	0.42	1.91	10	1
1:A:96:PHE:CD2	1:A:139:PRO:CB	0.42	3.03	8	1
1:A:98:TRP:CZ2	1:A:140:ILE:HG21	0.42	2.50	5	1
1:A:41:PHE:CE2	1:A:89:PHE:CE2	0.41	3.09	5	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/212 (97%)	173±3 (84±2%)	25±4 (12±2%)	7±2 (3±1%)	6	36
All	All	2050/2120 (97%)	1730 (84%)	251 (12%)	69 (3%)	6	36

All 25 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	125	ASP	8
1	A	112	ALA	7
1	A	122	TYR	7

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	47	GLY	7
1	A	48	SER	7
1	A	84	ASN	3
1	A	40	LYS	3
1	A	131	ASP	3
1	A	39	ASP	2
1	A	60	ILE	2
1	A	66	GLY	2
1	A	59	GLU	2
1	A	13	GLY	2
1	A	14	LYS	2
1	A	110	SER	2
1	A	71	LEU	1
1	A	85	GLY	1
1	A	202	ASN	1
1	A	74	GLY	1
1	A	3	VAL	1
1	A	86	LEU	1
1	A	212	GLU	1
1	A	192	PRO	1
1	A	193	LYS	1
1	A	57	HIS	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	188/193 (97%)	155±4 (83±2%)	33±4 (17±2%)	<b>4</b> 39
All	All	1880/1930 (97%)	1552 (83%)	328 (17%)	<b>4</b> 39

All 108 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	126	LYS	10
1	A	40	LYS	9
1	A	57	HIS	9

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	80	LEU	8
1	A	186	LYS	8
1	A	14	LYS	7
1	A	21	PHE	7
1	A	100	VAL	7
1	A	163	TYR	7
1	A	9	LYS	7
1	A	160	HIS	7
1	A	38	ASN	6
1	A	213	LYS	6
1	A	41	PHE	6
1	A	81	THR	6
1	A	155	THR	6
1	A	75	SER	5
1	A	133	ASP	5
1	A	211	SER	5
1	A	59	GLU	5
1	A	107	LYS	5
1	A	89	PHE	5
1	A	137	GLN	5
1	A	31	PHE	4
1	A	67	LYS	4
1	A	86	LEU	4
1	A	98	TRP	4
1	A	184	VAL	4
1	A	209	LEU	4
1	A	25	LYS	4
1	A	127	VAL	4
1	A	10	PHE	4
1	A	55	TRP	3
1	A	142	LYS	3
1	A	171	VAL	3
1	A	116	GLU	3
1	A	143	LEU	3
1	A	147	ASP	3
1	A	164	ASP	3
1	A	45	LEU	3
1	A	60	ILE	3
1	A	157	ARG	3
1	A	187	LYS	3
1	A	193	LYS	3
1	A	8	VAL	3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	52	CYS	3
1	A	158	GLN	3
1	A	161	THR	3
1	A	182	VAL	3
1	A	12	PHE	3
1	A	212	GLU	3
1	A	27	GLU	2
1	A	62	THR	2
1	A	188	ASP	2
1	A	202	ASN	2
1	A	42	MET	2
1	A	51	ARG	2
1	A	79	ASP	2
1	A	90	GLN	2
1	A	122	TYR	2
1	A	199	GLN	2
1	A	37	TRP	2
1	A	39	ASP	2
1	A	73	GLN	2
1	A	77	ASN	2
1	A	97	GLN	2
1	A	134	ASP	2
1	A	88	LYS	2
1	A	96	PHE	2
1	A	48	SER	2
1	A	83	ILE	2
1	A	131	ASP	2
1	A	138	LEU	2
1	A	195	MET	2
1	A	103	LYS	2
1	A	148	SER	2
1	A	174	LYS	2
1	A	110	SER	2
1	A	15	ASN	1
1	A	58	ASN	1
1	A	136	VAL	1
1	A	54	ILE	1
1	A	130	LYS	1
1	A	200	ASP	1
1	A	201	ASP	1
1	A	69	GLN	1
1	A	149	GLU	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	84	ASN	1
1	A	49	ASN	1
1	A	101	ASP	1
1	A	115	TYR	1
1	A	29	VAL	1
1	A	82	SER	1
1	A	153	HIS	1
1	A	194	ASN	1
1	A	104	ILE	1
1	A	150	ILE	1
1	A	19	GLU	1
1	A	30	ARG	1
1	A	53	ASN	1
1	A	70	GLU	1
1	A	105	VAL	1
1	A	154	LEU	1
1	A	170	SER	1
1	A	189	GLU	1
1	A	191	PHE	1
1	A	129	CYS	1
1	A	166	VAL	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 7 Chemical shift validation

No chemical shift data were provided