



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 18, 2024 – 11:50 am BST

PDB ID : 7B26
Title : CirpA1 in complex with pseudo-monomeric Properdin lacking TSR2-3
Authors : Lea, S.M.; Johnson, S.; Braunger, K.
Deposited on : 2020-11-26
Resolution : 3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	247	 64% 13% 23%
2	A	134	 61% 18% 21%
3	C	180	 69% 17% 13%
4	D	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MAN	B	506	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Properdin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	191	1498	926	283	268	21	0	1	0

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	229	MET	-	initiating methionine	UNP P27918
B	230	ILE	-	expression tag	UNP P27918
B	231	THR	-	expression tag	UNP P27918
B	232	GLU	-	expression tag	UNP P27918
B	233	GLY	-	expression tag	UNP P27918
B	234	ALA	-	expression tag	UNP P27918
B	235	GLN	-	expression tag	UNP P27918
B	236	ALA	-	expression tag	UNP P27918
B	237	PRO	-	expression tag	UNP P27918
B	238	ARG	-	expression tag	UNP P27918
B	239	LEU	-	expression tag	UNP P27918
B	240	LEU	-	expression tag	UNP P27918
B	241	LEU	-	expression tag	UNP P27918
B	242	PRO	-	expression tag	UNP P27918
B	243	PRO	-	expression tag	UNP P27918
B	244	LEU	-	expression tag	UNP P27918
B	245	LEU	-	expression tag	UNP P27918
B	246	LEU	-	expression tag	UNP P27918
B	247	LEU	-	expression tag	UNP P27918
B	248	LEU	-	expression tag	UNP P27918
B	249	THR	-	expression tag	UNP P27918
B	250	LEU	-	expression tag	UNP P27918
B	251	PRO	-	expression tag	UNP P27918
B	252	ALA	-	expression tag	UNP P27918
B	253	THR	-	expression tag	UNP P27918
B	254	GLY	-	expression tag	UNP P27918
B	255	SER	-	expression tag	UNP P27918

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	470	HIS	-	expression tag	UNP P27918
B	471	HIS	-	expression tag	UNP P27918
B	472	HIS	-	expression tag	UNP P27918
B	473	HIS	-	expression tag	UNP P27918
B	474	HIS	-	expression tag	UNP P27918
B	475	HIS	-	expression tag	UNP P27918

- Molecule 2 is a protein called Properdin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	106	805	495	144	154	12	0	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	ALA	CYS	engineered mutation	UNP P27918

- Molecule 3 is a protein called CirpA1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	156	1261	796	206	253	6	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-glucopyranose-(1-3)-alpha-L-fucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
4	D	2	21	12	9	0	0	0

- Molecule 5 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 11 6 5	0	0
5	B	1	Total C O 11 6 5	0	0
5	B	1	Total C O 11 6 5	0	0
5	B	1	Total C O 11 6 5	0	0
5	B	1	Total C O 11 6 5	0	0
5	B	1	Total C O 11 6 5	0	0
5	B	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0

FUCL
BCC2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	166.31Å 54.51Å 70.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.71 – 3.40 83.16 – 2.84	Depositor EDS
% Data completeness (in resolution range)	73.2 (24.71-3.40) 47.2 (83.16-2.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.86Å)	Xtrriage
Refinement program	PHENIX 1.18rc4_3812	Depositor
R, R_{free}	0.229 , 0.274 0.229 , 0.278	Depositor DCC
R_{free} test set	364 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	61.0	Xtrriage
Anisotropy	0.353	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 3.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3684	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BGC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.30	0/1547	0.58	0/2104
2	A	0.31	0/824	0.56	0/1116
3	C	0.29	0/1291	0.52	0/1749
All	All	0.30	0/3662	0.55	0/4969

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1498	0	1425	30	0
2	A	805	0	748	16	0
3	C	1261	0	1195	27	0
4	D	21	0	19	0	0
5	A	22	0	20	0	0
5	B	77	0	70	0	0
All	All	3684	0	3477	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:418:VAL:HB	1:B:426:GLU:HA	1.57	0.83
3:C:84:GLN:HG2	3:C:85:THR:HG23	1.66	0.77
1:B:374:GLN:HG3	1:B:375:HIS:CD2	2.22	0.74
1:B:329:ARG:HB2	1:B:335:ILE:HD11	1.70	0.72
1:B:418:VAL:HG11	1:B:426:GLU:HB2	1.71	0.71
2:A:33:PHE:CG	2:A:43:CYS:HB3	2.33	0.64
1:B:456:LEU:HD11	2:A:95:GLU:O	1.96	0.64
2:A:89:CYS:HB3	2:A:98:GLN:HB3	1.77	0.64
3:C:139:SER:HB3	3:C:144:CYS:HB3	1.78	0.64
3:C:135:LEU:HD13	3:C:147:PHE:HE1	1.63	0.62
1:B:319:ASP:OD2	1:B:351:ARG:NH1	2.30	0.62
3:C:49:VAL:HG21	3:C:112:ASN:HD22	1.65	0.61
3:C:49:VAL:HG11	3:C:112:ASN:HB3	1.84	0.59
2:A:91:VAL:HG21	2:A:94:SER:O	2.03	0.59
3:C:129:GLY:HA3	3:C:153:VAL:HG22	1.84	0.58
3:C:135:LEU:HD13	3:C:147:PHE:CE1	2.37	0.58
1:B:418:VAL:CB	1:B:426:GLU:HA	2.31	0.57
2:A:60:THR:HG23	2:A:76:ARG:HA	1.86	0.56
1:B:344:GLN:HG3	1:B:370:CYS:SG	2.46	0.56
1:B:335:ILE:HG23	1:B:340:ILE:HB	1.88	0.56
3:C:102:GLU:OE2	3:C:127:SER:OG	2.20	0.55
3:C:129:GLY:HA3	3:C:153:VAL:CG2	2.36	0.55
1:B:329:ARG:CB	1:B:335:ILE:HD11	2.36	0.55
1:B:327:CYS:HB2	1:B:344:GLN:HE21	1.72	0.55
1:B:353:ARG:HD2	1:B:359:ARG:HA	1.88	0.55
2:A:94:SER:OG	2:A:95:GLU:N	2.40	0.55
1:B:329:ARG:NH2	3:C:185:GLU:OE2	2.37	0.55
1:B:402:ALA:HB2	1:B:452:LYS:HD3	1.89	0.54
1:B:463:ASP:OD1	1:B:464:PRO:HD2	2.07	0.54
2:A:71:LEU:HD21	2:A:73:GLN:HE21	1.73	0.54
2:A:95:GLU:HG2	2:A:128:GLU:OE2	2.08	0.53
3:C:148:GLN:OE1	3:C:161:ARG:NE	2.43	0.52
2:A:33:PHE:CD2	2:A:43:CYS:HB3	2.45	0.51
2:A:92:THR:HG23	2:A:93:CYS:H	1.76	0.51
1:B:458:VAL:HG22	2:A:124:LEU:HD21	1.93	0.50
3:C:136:TYR:CZ	3:C:146:LEU:HB3	2.47	0.50
1:B:340:ILE:HD12	3:C:189:LEU:HB3	1.92	0.50
3:C:60:GLU:HG2	3:C:78:LYS:HG2	1.93	0.50
3:C:82:MET:HG3	3:C:87:THR:HB	1.93	0.50
2:A:86:TRP:CH2	2:A:125:GLN:HB2	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:143:LYS:O	3:C:165:VAL:HA	2.13	0.49
1:B:418:VAL:HG12	1:B:427:LYS:H	1.79	0.47
1:B:263:TRP:CE3	1:B:280:GLU:HG3	2.49	0.47
1:B:385:TRP:CG	1:B:403:ARG:NH1	2.83	0.47
3:C:49:VAL:HG11	3:C:112:ASN:CB	2.45	0.46
2:A:111:CYS:HB2	2:A:115:VAL:CG1	2.46	0.45
3:C:56:MET:HA	3:C:61:ILE:HD11	1.97	0.45
1:B:463:ASP:OD1	2:A:120:LEU:HD11	2.15	0.45
3:C:50:TYR:CE2	3:C:137:TYR:HA	2.52	0.45
1:B:329:ARG:HB3	1:B:332:MET:HB3	1.99	0.44
3:C:49:VAL:HG21	3:C:112:ASN:ND2	2.31	0.44
1:B:388:TRP:CE2	1:B:403:ARG:HD3	2.53	0.43
1:B:418:VAL:HG11	1:B:426:GLU:CB	2.45	0.42
2:A:80:TRP:HE1	2:A:112:SER:HB2	1.84	0.42
1:B:374:GLN:HG3	1:B:375:HIS:HD2	1.76	0.42
1:B:333:LYS:HE2	3:C:137:TYR:HD1	1.84	0.42
1:B:321:TRP:HB3	1:B:346:ARG:HD2	2.01	0.41
1:B:369:HIS:HB2	1:B:428:ASN:O	2.21	0.41
3:C:80:LYS:HG2	3:C:88:SER:O	2.20	0.41
3:C:49:VAL:HG11	3:C:112:ASN:HD22	1.85	0.41
3:C:80:LYS:HE3	3:C:88:SER:HB3	2.01	0.41
1:B:319:ASP:CG	1:B:351:ARG:HH12	2.21	0.41
2:A:31:LEU:HD23	2:A:49:GLY:HA2	2.03	0.41
3:C:82:MET:HA	3:C:87:THR:HA	2.02	0.41
3:C:147:PHE:O	3:C:161:ARG:HA	2.21	0.40
1:B:338:GLN:OE1	3:C:159:ILE:HD12	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	186/247 (75%)	181 (97%)	5 (3%)	0	100	100
2	A	104/134 (78%)	95 (91%)	9 (9%)	0	100	100
3	C	154/180 (86%)	151 (98%)	3 (2%)	0	100	100
All	All	444/561 (79%)	427 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	168/212 (79%)	168 (100%)	0	100	100
2	A	88/111 (79%)	88 (100%)	0	100	100
3	C	144/163 (88%)	144 (100%)	0	100	100
All	All	400/486 (82%)	400 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FUC	D	1	4,1	10,10,11	1.88	3 (30%)	14,14,16	0.99	1 (7%)
4	BGC	D	2	4	11,11,12	1.73	3 (27%)	15,15,17	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FUC	D	1	4,1	-	-	0/1/1/1
4	BGC	D	2	4	-	2/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	FUC	O5-C1	4.61	1.51	1.43
4	D	2	BGC	O5-C1	4.29	1.50	1.43
4	D	1	FUC	C2-C3	-2.62	1.48	1.52
4	D	2	BGC	C2-C3	-2.59	1.48	1.52
4	D	2	BGC	O5-C5	2.16	1.47	1.43
4	D	1	FUC	O5-C5	2.14	1.48	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	FUC	C6-C5-C4	-2.03	109.33	113.07

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	2	BGC	C4-C5-C6-O6
4	D	2	BGC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MAN	B	505	1	11,11,12	1.83	4 (36%)	15,15,17	1.29	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MAN	B	501	1	11,11,12	1.91	4 (36%)	15,15,17	1.16	1 (6%)
5	MAN	A	201	2	11,11,12	1.91	4 (36%)	15,15,17	1.12	1 (6%)
5	MAN	B	506	1	11,11,12	1.93	4 (36%)	15,15,17	1.03	1 (6%)
5	MAN	B	503	1	11,11,12	1.88	4 (36%)	15,15,17	1.29	2 (13%)
5	MAN	B	502	1	11,11,12	1.86	4 (36%)	15,15,17	1.16	2 (13%)
5	MAN	A	200	2	11,11,12	1.95	4 (36%)	15,15,17	1.11	1 (6%)
5	MAN	B	504	1	11,11,12	1.92	4 (36%)	15,15,17	1.42	3 (20%)
5	MAN	B	507	1	11,11,12	1.90	4 (36%)	15,15,17	1.26	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MAN	B	505	1	-	2/2/19/22	0/1/1/1
5	MAN	B	501	1	-	1/2/19/22	0/1/1/1
5	MAN	A	201	2	-	2/2/19/22	0/1/1/1
5	MAN	B	506	1	1/1/4/5	0/2/19/22	0/1/1/1
5	MAN	B	503	1	-	0/2/19/22	0/1/1/1
5	MAN	B	502	1	-	2/2/19/22	0/1/1/1
5	MAN	A	200	2	-	0/2/19/22	0/1/1/1
5	MAN	B	504	1	-	2/2/19/22	0/1/1/1
5	MAN	B	507	1	-	0/2/19/22	0/1/1/1

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	504	MAN	O5-C1	4.51	1.50	1.43
5	B	507	MAN	O5-C1	4.47	1.50	1.43
5	A	200	MAN	O5-C1	4.44	1.50	1.43
5	B	506	MAN	O5-C1	4.42	1.50	1.43
5	A	201	MAN	O5-C1	4.41	1.50	1.43
5	B	501	MAN	O5-C1	4.39	1.50	1.43
5	B	503	MAN	O5-C1	4.37	1.50	1.43
5	B	502	MAN	O5-C1	4.32	1.50	1.43
5	B	505	MAN	O5-C1	4.29	1.50	1.43
5	B	506	MAN	C2-C3	-2.90	1.48	1.52
5	B	501	MAN	C2-C3	-2.88	1.48	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	200	MAN	C2-C3	-2.86	1.48	1.52
5	A	201	MAN	C2-C3	-2.74	1.48	1.52
5	B	502	MAN	C2-C3	-2.68	1.48	1.52
5	B	507	MAN	C2-C3	-2.65	1.48	1.52
5	B	503	MAN	C2-C3	-2.65	1.48	1.52
5	A	200	MAN	O5-C5	2.60	1.48	1.43
5	B	504	MAN	C2-C3	-2.57	1.48	1.52
5	B	505	MAN	C2-C3	-2.56	1.48	1.52
5	B	504	MAN	O5-C5	2.50	1.48	1.43
5	A	201	MAN	O5-C5	2.46	1.48	1.43
5	B	507	MAN	O5-C5	2.44	1.48	1.43
5	B	503	MAN	O5-C5	2.44	1.48	1.43
5	B	501	MAN	O5-C5	2.43	1.48	1.43
5	B	506	MAN	O5-C5	2.40	1.48	1.43
5	B	502	MAN	O5-C5	2.33	1.48	1.43
5	B	505	MAN	O5-C5	2.21	1.47	1.43
5	B	506	MAN	O3-C3	2.15	1.48	1.43
5	B	504	MAN	O3-C3	2.12	1.48	1.43
5	B	505	MAN	O3-C3	2.10	1.47	1.43
5	B	502	MAN	O3-C3	2.08	1.47	1.43
5	A	201	MAN	O3-C3	2.07	1.47	1.43
5	A	200	MAN	O3-C3	2.07	1.47	1.43
5	B	503	MAN	O3-C3	2.05	1.47	1.43
5	B	507	MAN	O3-C3	2.04	1.47	1.43
5	B	501	MAN	O3-C3	2.03	1.47	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	504	MAN	C1-C2-C3	3.08	113.45	109.67
5	B	501	MAN	C3-C4-C5	2.67	115.00	110.24
5	A	201	MAN	C3-C4-C5	2.64	114.95	110.24
5	B	505	MAN	C3-C4-C5	2.64	114.94	110.24
5	A	200	MAN	C3-C4-C5	2.60	114.87	110.24
5	B	503	MAN	C3-C4-C5	2.59	114.86	110.24
5	B	505	MAN	C1-C2-C3	2.57	112.83	109.67
5	B	507	MAN	C1-C2-C3	2.50	112.74	109.67
5	B	503	MAN	C1-C2-C3	2.44	112.67	109.67
5	B	502	MAN	C1-C2-C3	2.43	112.66	109.67
5	B	502	MAN	C3-C4-C5	2.41	114.54	110.24
5	B	504	MAN	C3-C4-C5	2.40	114.53	110.24
5	B	506	MAN	C3-C4-C5	2.24	114.23	110.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	504	MAN	O5-C5-C6	2.05	110.42	107.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	506	MAN	C1

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	201	MAN	C4-C5-C6-O6
5	B	505	MAN	C4-C5-C6-O6
5	B	504	MAN	O5-C5-C6-O6
5	A	201	MAN	O5-C5-C6-O6
5	B	502	MAN	C4-C5-C6-O6
5	B	505	MAN	O5-C5-C6-O6
5	B	504	MAN	C4-C5-C6-O6
5	B	502	MAN	O5-C5-C6-O6
5	B	501	MAN	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	191/247 (77%)	-0.45	0 100 100	38, 67, 95, 118	0
2	A	106/134 (79%)	-0.46	0 100 100	36, 53, 141, 158	0
3	C	156/180 (86%)	-0.36	0 100 100	31, 66, 97, 108	0
All	All	453/561 (80%)	-0.42	0 100 100	31, 65, 98, 158	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BGC	D	2	11/12	0.94	0.14	27,33,44,62	0
4	FUC	D	1	10/11	0.97	0.13	27,41,42,42	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MAN	A	201	11/12	0.83	0.48	84,108,117,124	0
5	MAN	B	505	11/12	0.91	0.19	49,61,82,95	0
5	MAN	B	502	11/12	0.94	0.18	33,51,58,61	0
5	MAN	B	506	11/12	0.94	0.17	53,59,72,88	0
5	MAN	B	507	11/12	0.94	0.16	63,67,70,80	0
5	MAN	A	200	11/12	0.94	0.14	47,55,60,63	0
5	MAN	B	504	11/12	0.94	0.09	61,72,84,85	0
5	MAN	B	501	11/12	0.95	0.15	43,49,52,54	0
5	MAN	B	503	11/12	0.95	0.16	59,68,81,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.