



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 26, 2024 – 12:35 PM EDT

PDB ID : 9B44  
Title : Crystal structure of mAb 8-24 Fab, a VRC01-like HIV-1 antibody  
Authors : Kher, G.; Hurlburt, N.; Pancera, M.  
Deposited on : 2024-03-20  
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.3


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div>
1	H	233	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
2	B	216	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 87%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
2	L	216	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 82%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	302	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6640 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MLK8-24 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	215	Total 1656	C 1048	N 286	O 315	S 7	0	0	0
1	H	218	Total 1677	C 1060	N 290	O 320	S 7	0	0	0

- Molecule 2 is a protein called MLK8-24 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	203	Total 1566	C 985	N 258	O 318	S 5	0	0	0
2	L	202	Total 1558	C 979	N 257	O 317	S 5	0	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	H	1	Total C O 4 2 2	0	0
3	L	1	Total C O 4 2 2	0	0
3	L	1	Total C O 4 2 2	0	0
3	L	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	42	Total O 42 42	0	0
4	B	43	Total O 43 43	0	0
4	H	32	Total O 32 32	0	0
4	L	26	Total O 26 26	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MLK8-24 Fab Heavy Chain




- Molecule 1: MLK8-24 Fab Heavy Chain




- Molecule 2: MLK8-24 Fab Light Chain



- Molecule 2: MLK8-24 Fab Light Chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.87Å 77.70Å 266.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.36 – 2.01 44.36 – 2.01	Depositor EDS
% Data completeness (in resolution range)	98.9 (44.36-2.01) 98.9 (44.36-2.01)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 2.01Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158, PDB-REDO	Depositor
R, $R_{free}$	0.227 , 0.281 0.230 , 0.282	Depositor DCC
$R_{free}$ test set	2937 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.4	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6640	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1696	0.65	0/2316
1	H	0.40	0/1717	0.63	0/2343
2	B	0.43	0/1597	0.62	0/2164
2	L	0.41	0/1589	0.60	0/2153
All	All	0.41	0/6599	0.62	0/8976

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1656	0	1590	23	0
1	H	1677	0	1613	13	0
2	B	1566	0	1535	8	0
2	L	1558	0	1524	13	0
3	A	12	0	18	8	0
3	B	12	0	18	2	0
3	H	4	0	6	2	0
3	L	12	0	18	0	0
4	A	42	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	43	0	0	0	0
4	H	32	0	0	1	0
4	L	26	0	0	0	0
All	All	6640	0	6322	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:ASN:HB3	3:A:302:EDO:H12	1.54	0.90
2:B:183:GLU:O	2:B:207:ARG:NH2	2.19	0.76
1:A:35:HIS:HD1	1:A:47:TRP:HE1	1.35	0.74
1:A:100:TYR:HE2	3:B:302:EDO:H22	1.57	0.69
1:A:170:LEU:HD11	3:A:301:EDO:H22	1.73	0.68
1:A:33:PHE:HB2	1:A:100(A):ASN:HD21	1.59	0.67
1:A:112:SER:HB2	3:A:301:EDO:H12	1.77	0.66
1:A:61:GLN:HA	1:A:64:GLN:HG3	1.79	0.64
1:A:100:TYR:CE2	3:B:302:EDO:H22	2.33	0.64
1:A:56:GLY:H	3:A:302:EDO:H11	1.63	0.64
2:L:18:ARG:HG3	2:L:76:SER:HA	1.79	0.63
2:L:32:TYR:HB3	2:L:90:GLN:HE21	1.65	0.62
1:H:201:LYS:HG2	1:H:202:PRO:HD3	1.83	0.61
1:H:198:VAL:HB	1:H:207:VAL:HG23	1.82	0.60
1:H:195:ILE:HG12	1:H:210:ARG:HG3	1.83	0.59
1:A:198:VAL:HB	1:A:207:VAL:HG23	1.84	0.59
2:L:103:LYS:HD2	2:L:104:ARG:N	2.19	0.58
2:L:120:GLN:HE21	2:L:125:THR:HG23	1.69	0.57
1:A:201:LYS:HE3	1:A:201:LYS:HA	1.87	0.57
2:L:13:VAL:HB	2:L:19:ILE:HD11	1.88	0.56
2:L:19:ILE:HD12	2:L:78:VAL:HB	1.90	0.54
1:H:36:TRP:CE2	1:H:80:MET:HB2	2.44	0.53
1:H:95:GLY:HA3	3:H:301:EDO:H11	1.90	0.52
1:A:47:TRP:CE3	2:B:92:GLU:HB3	2.46	0.51
3:A:303:EDO:H21	2:B:94:PHE:H	1.78	0.48
1:H:149:PRO:HD2	1:H:201:LYS:HE3	1.95	0.48
1:A:53:PHE:CD2	1:A:54:ARG:HG3	2.49	0.48
1:A:152:VAL:O	4:A:401:HOH:O	2.20	0.47
1:A:36:TRP:CE2	1:A:80:MET:HB2	2.49	0.46
2:B:141:LYS:HD2	2:B:193:THR:HB	1.98	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:201:LYS:HG2	1:H:202:PRO:CD	2.45	0.46
2:B:187:VAL:HG22	2:B:206:ASN:OD1	2.15	0.46
3:A:303:EDO:C2	2:B:94:PHE:H	2.29	0.46
1:H:97:ASN:OD1	1:H:99:ASP:HB2	2.15	0.46
2:L:27:LEU:HD23	2:L:27:LEU:HA	1.77	0.45
2:L:66:GLY:HA3	2:L:71:PHE:CD2	2.52	0.45
1:A:47:TRP:HZ2	1:A:50:TRP:CD1	2.34	0.45
1:A:60:PRO:O	1:A:64:GLN:HG2	2.17	0.44
2:L:116:PRO:HD3	2:L:128:VAL:HG22	1.99	0.44
1:A:119:PRO:HD2	1:A:205:THR:HG21	1.99	0.44
1:H:152:VAL:HG11	1:H:180:SER:CB	2.47	0.44
1:A:52(A):PRO:HA	1:A:71:ARG:HG2	2.00	0.43
2:B:185:HIS:O	2:B:207:ARG:NE	2.51	0.43
1:A:56:GLY:H	3:A:302:EDO:C1	2.30	0.42
1:H:19:ARG:NH2	4:H:401:HOH:O	2.37	0.42
1:A:62:LYS:HG3	1:A:63:PHE:CD2	2.54	0.42
1:H:94:ARG:HG2	3:H:301:EDO:H22	2.02	0.42
2:L:180:ALA:O	2:L:184:LYS:HG3	2.19	0.42
1:H:154:TRP:CH2	1:H:196:CYS:HB3	2.55	0.41
1:A:189:LEU:HD23	1:A:189:LEU:HA	1.82	0.41
2:L:21:LEU:HD23	2:L:21:LEU:HA	1.81	0.41
1:A:56:GLY:N	3:A:302:EDO:H11	2.33	0.41
2:L:33:LEU:HD13	2:L:71:PHE:CD1	2.56	0.41
1:H:47:TRP:CZ2	1:H:49:GLY:HA2	2.56	0.40
2:L:32:TYR:N	2:L:51:THR:HG1	2.19	0.40
2:B:101:GLU:CD	2:B:169:TYR:HH	2.25	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/233 (91%)	206 (98%)	5 (2%)	0	100	100
1	H	214/233 (92%)	208 (97%)	6 (3%)	0	100	100
2	B	199/216 (92%)	193 (97%)	6 (3%)	0	100	100
2	L	198/216 (92%)	192 (97%)	6 (3%)	0	100	100
All	All	822/898 (92%)	799 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/200 (92%)	178 (97%)	5 (3%)	40	42
1	H	186/200 (93%)	182 (98%)	4 (2%)	47	51
2	B	180/191 (94%)	176 (98%)	4 (2%)	47	51
2	L	179/191 (94%)	174 (97%)	5 (3%)	38	40
All	All	728/782 (93%)	710 (98%)	18 (2%)	42	45

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	ARG
1	A	97	ASN
1	A	100(A)	ASN
1	A	179	SER
1	A	197	ASN
2	B	27	LEU
2	B	27(A)	THR
2	B	33	LEU
2	B	47	LEU
1	H	71	ARG
1	H	96	LYS
1	H	129	LYS
1	H	178	LEU

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
2	L	25	SER
2	L	27(A)	THR
2	L	33	LEU
2	L	103	LYS
2	L	123	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100(A)	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PCA	H	1	1	7,8,9	2.03	1 (14%)	9,10,12	2.44	6 (66%)
1	PCA	A	1	1	7,8,9	1.76	1 (14%)	9,10,12	2.63	6 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	H	1	1	-	0/0/11/13	0/1/1/1
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	1	PCA	CD-N	5.06	1.47	1.34
1	A	1	PCA	CD-N	4.42	1.45	1.34

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CA-N-CD	-3.58	101.33	113.58
1	A	1	PCA	CG-CD-N	3.38	116.67	108.39
1	A	1	PCA	CB-CA-C	-3.29	108.14	112.66
1	H	1	PCA	CA-N-CD	-3.29	102.32	113.58
1	H	1	PCA	CG-CD-N	3.28	116.42	108.39
1	H	1	PCA	OE-CD-CG	-3.17	121.06	126.72
1	A	1	PCA	OE-CD-CG	-2.90	121.55	126.72
1	A	1	PCA	CB-CA-N	2.73	110.76	103.24
1	H	1	PCA	CB-CA-N	2.65	110.52	103.24
1	H	1	PCA	CB-CG-CD	-2.55	100.47	104.41
1	A	1	PCA	CB-CG-CD	-2.51	100.53	104.41
1	H	1	PCA	O-C-CA	-2.21	119.08	124.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	B	302	-	3,3,3	0.27	0	2,2,2	0.60	0
3	EDO	L	301	-	3,3,3	0.54	0	2,2,2	0.07	0
3	EDO	B	301	-	3,3,3	0.40	0	2,2,2	0.50	0
3	EDO	A	302	-	3,3,3	0.35	0	2,2,2	0.15	0
3	EDO	A	301	-	3,3,3	0.39	0	2,2,2	0.44	0
3	EDO	H	301	-	3,3,3	0.42	0	2,2,2	0.50	0
3	EDO	A	303	-	3,3,3	0.49	0	2,2,2	0.38	0
3	EDO	B	300	-	3,3,3	0.49	0	2,2,2	0.33	0
3	EDO	L	302	-	3,3,3	0.43	0	2,2,2	0.43	0
3	EDO	L	303	-	3,3,3	0.38	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	302	-	-	1/1/1/1	-
3	EDO	L	301	-	-	0/1/1/1	-
3	EDO	B	301	-	-	0/1/1/1	-
3	EDO	A	302	-	-	1/1/1/1	-
3	EDO	A	301	-	-	0/1/1/1	-
3	EDO	H	301	-	-	1/1/1/1	-
3	EDO	A	303	-	-	1/1/1/1	-
3	EDO	B	300	-	-	1/1/1/1	-
3	EDO	L	302	-	-	1/1/1/1	-
3	EDO	L	303	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	EDO	O1-C1-C2-O2
3	L	302	EDO	O1-C1-C2-O2
3	A	302	EDO	O1-C1-C2-O2
3	A	303	EDO	O1-C1-C2-O2
3	B	300	EDO	O1-C1-C2-O2
3	H	301	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	EDO	2	0
3	A	302	EDO	4	0
3	A	301	EDO	2	0
3	H	301	EDO	2	0
3	A	303	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	214/233 (91%)	0.33	12 (5%) 31 29	18, 29, 53, 73	0
1	H	217/233 (93%)	0.43	12 (5%) 32 30	21, 31, 49, 81	0
2	B	203/216 (93%)	0.38	8 (3%) 44 42	20, 30, 49, 91	0
2	L	202/216 (93%)	0.47	10 (4%) 35 33	20, 32, 53, 85	0
All	All	836/898 (93%)	0.40	42 (5%) 35 33	18, 31, 52, 91	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	27(A)	THR	8.1
2	B	27(B)	LEU	6.6
2	L	27	LEU	5.7
2	B	32	TYR	5.3
2	L	32	TYR	5.2
2	B	27(A)	THR	4.5
2	L	207	ARG	4.1
1	H	129	LYS	4.0
2	B	27	LEU	3.6
1	A	53	PHE	3.4
1	A	64	GLN	3.4
1	A	100	TYR	3.4
1	H	53	PHE	3.2
2	B	3	VAL	3.1
2	B	150	LEU	3.1
2	L	3	VAL	3.1
1	A	127	SER	3.0
1	A	190	GLY	3.0
2	L	150	LEU	2.9
1	A	213	PRO	2.9
1	H	130	SER	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	H	64	GLN	2.8
1	A	100(A)	ASN	2.7
1	H	75	PHE	2.7
1	H	127	SER	2.7
1	H	65	GLY	2.7
1	A	191	THR	2.7
2	B	25	SER	2.5
2	L	67	SER	2.4
1	A	99	ASP	2.4
2	L	91	TYR	2.4
1	A	75	PHE	2.3
1	A	205	THR	2.3
1	H	186	SER	2.3
2	L	180	ALA	2.1
1	H	134	GLY	2.1
1	A	97	ASN	2.1
1	H	204	ASN	2.1
1	H	100	TYR	2.0
2	L	187	VAL	2.0
2	B	26	SER	2.0
1	H	210	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PCA	A	1	8/9	0.90	0.10	20,20,20,20	0
1	PCA	H	1	8/9	0.90	0.18	20,20,20,20	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	301	4/4	0.83	0.16	29,39,39,56	0
3	EDO	B	301	4/4	0.83	0.13	37,38,39,44	0
3	EDO	H	301	4/4	0.85	0.16	32,47,48,48	0
3	EDO	L	303	4/4	0.85	0.14	42,46,48,50	0
3	EDO	L	302	4/4	0.87	0.13	33,35,38,47	0
3	EDO	L	301	4/4	0.89	0.13	33,38,38,42	0
3	EDO	A	303	4/4	0.91	0.10	28,32,36,37	0
3	EDO	B	302	4/4	0.91	0.12	35,41,42,45	0
3	EDO	B	300	4/4	0.93	0.14	14,31,35,42	0
3	EDO	A	302	4/4	0.94	0.11	15,22,25,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.