



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 04:28 PM EDT

PDB ID : 3B5D  
Title : EmrE multidrug transporter in complex with TPP, C2 crystal form  
Authors : Chang, G.; Chen, Y.J.  
Deposited on : 2007-10-25  
Resolution : 3.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1212 (4.00-3.60)
Clashscore	141614	1288 (4.00-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	110	 91% 9%
1	B	110	 90% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	P4P	A	350	-	X	-	-

## 2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug transporter emrE.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	100	Total C 100 100	0	0	100
1	B	99	Total C 99 99	0	0	99

- Molecule 2 is TETRAPHENYLPHOSPHONIUM (three-letter code: P4P) (formula:  $C_{24}H_{20}P$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C P 25 24 1	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Multidrug transporter emrE

Chain A:  91% 9%



- Molecule 1: Multidrug transporter emrE

Chain B:  90% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	115.10Å 43.70Å 76.40Å 90.00° 108.10° 90.00°	Depositor
Resolution (Å)	20.00 – 3.80 19.99 – 3.80	Depositor EDS
% Data completeness (in resolution range)	83.6 (20.00-3.80) 83.6 (19.99-3.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.71 (at 3.82Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.325 , 0.364 0.465 , 0.478	Depositor DCC
$R_{free}$ test set	335 reflections (10.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	142.5	Xtrriage
Anisotropy	0.764	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.69 , 153.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.77	EDS
Total number of atoms	224	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	213.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P4P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	100	0	0	0	0
1	B	99	0	0	0	0
2	A	25	0	20	0	0
All	All	224	0	20	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	P4P	A	350	-	28,28,28	4.19	23 (82%)	38,38,38	2.50	16 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P4P	A	350	-	-	0/24/24/24	0/4/4/4

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	350	P4P	C6D-C1D	7.16	1.50	1.39
2	A	350	P4P	C5A-C6A	6.96	1.50	1.38
2	A	350	P4P	C5D-C6D	6.87	1.50	1.38
2	A	350	P4P	C2D-C1D	6.72	1.50	1.39
2	A	350	P4P	C3B-C2B	5.90	1.49	1.38
2	A	350	P4P	C2C-C1C	5.25	1.47	1.39
2	A	350	P4P	C2B-C1B	5.08	1.47	1.39
2	A	350	P4P	C6C-C1C	4.72	1.46	1.39
2	A	350	P4P	P-C1D	4.45	1.87	1.79
2	A	350	P4P	C3C-C4C	4.40	1.47	1.38
2	A	350	P4P	C5A-C4A	3.86	1.46	1.38
2	A	350	P4P	C3A-C2A	3.81	1.45	1.38
2	A	350	P4P	P-C1B	3.74	1.86	1.79
2	A	350	P4P	C4C-C5C	3.73	1.46	1.38
2	A	350	P4P	C4D-C5D	3.71	1.46	1.38
2	A	350	P4P	P-C1C	3.57	1.86	1.79
2	A	350	P4P	C3D-C2D	3.44	1.44	1.38
2	A	350	P4P	C5B-C6B	3.37	1.44	1.38
2	A	350	P4P	C2A-C1A	2.60	1.43	1.39
2	A	350	P4P	C3D-C4D	2.54	1.43	1.38
2	A	350	P4P	C6B-C1B	2.51	1.43	1.39
2	A	350	P4P	C3B-C4B	2.50	1.43	1.38
2	A	350	P4P	P-C1A	2.44	1.84	1.79

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	350	P4P	C1D-P-C1C	5.95	121.42	109.43
2	A	350	P4P	C6A-C1A-C2A	5.27	125.49	118.93
2	A	350	P4P	C1D-P-C1A	-4.27	100.82	109.43
2	A	350	P4P	P-C1B-C2B	-4.20	112.62	120.05
2	A	350	P4P	C3A-C2A-C1A	-3.91	115.81	120.29
2	A	350	P4P	C4A-C5A-C6A	-3.68	115.69	120.24
2	A	350	P4P	C3B-C2B-C1B	-3.60	116.16	120.29
2	A	350	P4P	C5A-C4A-C3A	3.17	124.21	119.87
2	A	350	P4P	P-C1B-C6B	3.10	125.53	120.05
2	A	350	P4P	C1B-P-C1A	3.08	115.64	109.43
2	A	350	P4P	P-C1C-C2C	-2.81	115.07	120.05
2	A	350	P4P	P-C1C-C6C	2.73	124.87	120.05
2	A	350	P4P	C3D-C4D-C5D	2.70	123.56	119.87
2	A	350	P4P	C5A-C6A-C1A	-2.65	117.25	120.29
2	A	350	P4P	C2B-C1B-C6B	2.35	121.85	118.93
2	A	350	P4P	C5D-C6D-C1D	-2.04	117.96	120.29




There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.