



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2023 – 07:33 pm BST

PDB ID : 8B74
Title : Tyrosinase from halophilic bacterium *Hahella* sp. CCB MM4
Authors : Rohr, A.K.; Santos, G.A.; Englund, A.N.
Deposited on : 2022-09-28
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6837 atoms, of which 3226 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxidoreductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	430	6654	2166	3226	593	656	13	245	3	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	183	Total	O	0	0
			183	183		

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	51.14Å 55.57Å 156.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.27 – 1.97 78.15 – 1.96	Depositor EDS
% Data completeness (in resolution range)	99.2 (78.27-1.97) 95.4 (78.15-1.96)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.97Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.175 , 0.226 0.185 , 0.234	Depositor DCC
R_{free} test set	1498 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtrriage
Anisotropy	0.577	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6837	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	4/3527 (0.1%)	1.10	17/4810 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	13	ASP	CG-OD1	10.42	1.49	1.25
1	A	400	GLU	CD-OE1	6.28	1.32	1.25
1	A	23	GLU	CD-OE1	-5.61	1.19	1.25
1	A	388	GLU	CD-OE2	5.18	1.31	1.25

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	458	ARG	NE-CZ-NH1	12.66	126.63	120.30
1	A	13	ASP	CB-CG-OD2	-11.32	108.11	118.30
1	A	13	ASP	CB-CG-OD1	7.91	125.42	118.30
1	A	28	ARG	NE-CZ-NH1	7.79	124.19	120.30
1	A	279	ILE	CA-CB-CG1	7.49	125.23	111.00
1	A	28	ARG	NE-CZ-NH2	-6.19	117.21	120.30
1	A	462	HIS	CB-CA-C	-6.00	98.39	110.40
1	A	88	ARG	NE-CZ-NH2	-5.71	117.44	120.30
1	A	75	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	A	356	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	A	458	ARG	NE-CZ-NH2	-5.57	117.52	120.30
1	A	458	ARG	CD-NE-CZ	5.57	131.39	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	380	ARG	CB-CA-C	-5.26	99.87	110.40
1	A	4	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	A	279	ILE	N-CA-CB	5.21	122.77	110.80
1	A	130	ASN	N-CA-CB	5.14	119.84	110.60
1	A	360	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	297	ALA	Peptide
1	A	334	LEU	Peptide
1	A	356	ARG	Sidechain
1	A	428	ARG	Sidechain
1	A	458	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3428	3226	3208	37	1
2	A	183	0	0	7	0
All	All	3611	3226	3208	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:400:GLU:OE2	2:A:501:HOH:O	1.88	0.91
1:A:76[A]:CYS:SG	1:A:78:HIS:CE1	2.80	0.75
1:A:315:VAL:HG22	1:A:462:HIS:CE1	2.23	0.73
1:A:96:GLU:HB3	1:A:272:ALA:HB2	1.71	0.72
1:A:418:ILE:HG22	1:A:420:ILE:HG23	1.73	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:MET:HE3	1:A:251:GLN:NE2	2.11	0.66
1:A:426:LYS:HD2	2:A:601:HOH:O	2.00	0.61
1:A:2:LYS:HB3	1:A:112:THR:HG21	1.84	0.60
1:A:298:PHE:O	1:A:299:PRO:C	2.40	0.59
1:A:76[A]:CYS:HG	1:A:78:HIS:CE1	2.20	0.59
1:A:418:ILE:HG22	1:A:420:ILE:CG2	2.33	0.59
1:A:367:LYS:CB	1:A:400:GLU:HG2	2.32	0.58
1:A:386:ASN:H	1:A:389:ASN:HB2	1.67	0.58
1:A:386:ASN:HD22	1:A:386:ASN:C	2.08	0.57
1:A:420:ILE:HD11	1:A:461:LEU:HD11	1.87	0.56
1:A:387:GLU:OE2	2:A:502:HOH:O	2.18	0.53
1:A:426:LYS:CD	2:A:601:HOH:O	2.54	0.53
1:A:367:LYS:HB3	1:A:400:GLU:HG2	1.91	0.53
1:A:400:GLU:HB3	2:A:501:HOH:O	2.09	0.52
1:A:33:PRO:HA	1:A:38:TYR:CG	2.45	0.51
1:A:177:GLU:CD	1:A:187:ILE:HG22	2.32	0.50
1:A:246:GLN:NE2	2:A:507:HOH:O	2.44	0.49
1:A:367:LYS:HA	1:A:400:GLU:CG	2.44	0.47
1:A:91[A]:LEU:HD22	1:A:113:LEU:CD2	2.43	0.47
1:A:282:LYS:HA	1:A:285:GLU:HG3	1.96	0.47
1:A:103:ARG:HH11	1:A:103:ARG:HG3	1.81	0.46
1:A:279:ILE:HG22	1:A:283:ASP:OD2	2.17	0.45
1:A:384:GLU:O	1:A:389:ASN:ND2	2.50	0.45
1:A:379:ASN:HD21	1:A:434:ASP:H	1.63	0.44
1:A:376:PHE:HB3	1:A:389:ASN:HB3	1.98	0.44
1:A:177:GLU:CD	1:A:187:ILE:CG2	2.86	0.44
1:A:160:ARG:HD2	2:A:529:HOH:O	2.19	0.42
1:A:177:GLU:HG3	1:A:187:ILE:HG22	2.01	0.42
1:A:120:GLU:OE2	1:A:120:GLU:HA	2.20	0.41
1:A:118:TYR:HB2	1:A:169:VAL:HG11	2.02	0.41
1:A:361:ILE:HD11	1:A:375:LEU:HD13	2.01	0.41
1:A:266:ARG:NH1	1:A:278:GLU:OE1	2.54	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:LYS:HZ3	1:A:333:ASP:H[1_545]	1.35	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	425/466 (91%)	413 (97%)	10 (2%)	2 (0%)	29 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	ASP
1	A	335	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	367/391 (94%)	341 (93%)	26 (7%)	14 5

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ASP
1	A	53	THR
1	A	56	HIS
1	A	64	SER
1	A	65	GLN
1	A	81	TRP
1	A	118	TYR
1	A	127	SER
1	A	144	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	182	ASN
1	A	183	SER
1	A	219	ASN
1	A	258	VAL
1	A	279	ILE
1	A	300	LEU
1	A	332	LEU
1	A	337	VAL
1	A	379	ASN
1	A	382	ASN
1	A	384	GLU
1	A	386	ASN
1	A	393[A]	CYS
1	A	393[B]	CYS
1	A	426	LYS
1	A	435	GLU
1	A	458	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	HIS
1	A	78	HIS
1	A	130	ASN
1	A	251	GLN
1	A	363	ASN
1	A	386	ASN
1	A	401	ASN
1	A	455	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/466 (92%)	-0.02	6 (1%) 75 82	21, 32, 61, 90	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	VAL	3.9
1	A	128	PRO	3.1
1	A	412	SER	2.8
1	A	461	LEU	2.3
1	A	409	SER	2.1
1	A	404	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.