



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 7, 2023 – 02:19 PM JST

PDB ID : 5B89  
Title : Crystal structure of a Cysteine Desulfurase from *Thermococcus onnurineus* NA1 in complex with alanine at 1.5 Angstrom resolution  
Authors : Ho, T.-H.; Kang, L.-W.  
Deposited on : 2016-06-13  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	 2% 83% 11% • 5%
1	B	419	 % 83% 13% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ALA	B	401	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine desulfurase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	400	3121	1989	540	581	11	0	0	0
1	B	402	3135	1997	544	583	11	0	0	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP B6YT87
A	-18	HIS	-	expression tag	UNP B6YT87
A	-17	HIS	-	expression tag	UNP B6YT87
A	-16	HIS	-	expression tag	UNP B6YT87
A	-15	HIS	-	expression tag	UNP B6YT87
A	-14	HIS	-	expression tag	UNP B6YT87
A	-13	HIS	-	expression tag	UNP B6YT87
A	-12	SER	-	expression tag	UNP B6YT87
A	-11	SER	-	expression tag	UNP B6YT87
A	-10	GLU	-	expression tag	UNP B6YT87
A	-9	ASN	-	expression tag	UNP B6YT87
A	-8	LEU	-	expression tag	UNP B6YT87
A	-7	TYR	-	expression tag	UNP B6YT87
A	-6	PHE	-	expression tag	UNP B6YT87
A	-5	GLN	-	expression tag	UNP B6YT87
A	-4	GLY	-	expression tag	UNP B6YT87
A	-3	HIS	-	expression tag	UNP B6YT87
A	-2	MET	-	expression tag	UNP B6YT87
A	-1	ALA	-	expression tag	UNP B6YT87
A	0	SER	-	expression tag	UNP B6YT87
B	-19	MET	-	expression tag	UNP B6YT87
B	-18	HIS	-	expression tag	UNP B6YT87
B	-17	HIS	-	expression tag	UNP B6YT87
B	-16	HIS	-	expression tag	UNP B6YT87
B	-15	HIS	-	expression tag	UNP B6YT87

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP B6YT87
B	-13	HIS	-	expression tag	UNP B6YT87
B	-12	SER	-	expression tag	UNP B6YT87
B	-11	SER	-	expression tag	UNP B6YT87
B	-10	GLU	-	expression tag	UNP B6YT87
B	-9	ASN	-	expression tag	UNP B6YT87
B	-8	LEU	-	expression tag	UNP B6YT87
B	-7	TYR	-	expression tag	UNP B6YT87
B	-6	PHE	-	expression tag	UNP B6YT87
B	-5	GLN	-	expression tag	UNP B6YT87
B	-4	GLY	-	expression tag	UNP B6YT87
B	-3	HIS	-	expression tag	UNP B6YT87
B	-2	MET	-	expression tag	UNP B6YT87
B	-1	ALA	-	expression tag	UNP B6YT87
B	0	SER	-	expression tag	UNP B6YT87

- Molecule 2 is ALANINE (three-letter code: ALA) (formula: C<sub>3</sub>H<sub>7</sub>NO<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	6	3	1	2	0	0
2	B	1	6	3	1	2	0	0

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	15	8	1	5	1	0	0
3	B	1	15	8	1	5	1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	254	Total O 254 254	0	0
6	B	291	Total O 291 291	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cysteine desulfurase



- Molecule 1: Cysteine desulfurase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.94Å 78.72Å 171.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.16 – 1.50 49.16 – 1.50	Depositor EDS
% Data completeness (in resolution range)	94.1 (49.16-1.50) 94.1 (49.16-1.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.66 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.167 , 0.194 0.171 , 0.196	Depositor DCC
$R_{free}$ test set	6444 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.3	Xtrriage
Anisotropy	0.039	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6853	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CSS, PLP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.37	6/3180 (0.2%)	1.13	11/4297 (0.3%)
1	B	1.40	9/3195 (0.3%)	1.19	16/4317 (0.4%)
All	All	1.39	15/6375 (0.2%)	1.16	27/8614 (0.3%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	40	TYR	CE1-CZ	-7.06	1.29	1.38
1	B	79	GLU	CD-OE1	6.79	1.33	1.25
1	B	353	GLY	C-O	-6.33	1.13	1.23
1	B	384	GLU	CD-OE1	-6.13	1.19	1.25
1	A	220	GLY	C-O	-6.04	1.14	1.23
1	A	63	GLU	CD-OE2	-5.57	1.19	1.25
1	B	78	GLU	CD-OE2	5.57	1.31	1.25
1	A	225	GLY	C-O	-5.39	1.15	1.23
1	B	230	ARG	CD-NE	-5.35	1.37	1.46
1	B	5	GLU	CD-OE2	-5.34	1.19	1.25
1	B	62	TYR	CE1-CZ	-5.34	1.31	1.38
1	A	77	PHE	C-O	5.21	1.33	1.23
1	B	265	GLU	CD-OE1	5.16	1.31	1.25
1	B	129	GLY	C-O	-5.16	1.15	1.23
1	A	278	ALA	C-O	5.11	1.33	1.23

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	318	ARG	NE-CZ-NH2	-10.81	114.90	120.30
1	B	344	ASP	CB-CG-OD1	9.38	126.75	118.30
1	B	318	ARG	NE-CZ-NH1	8.30	124.45	120.30
1	A	123	ARG	NE-CZ-NH1	-7.85	116.38	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	8	ARG	NE-CZ-NH1	7.60	124.10	120.30
1	B	286	ARG	NE-CZ-NH1	7.57	124.08	120.30
1	A	308	ASP	CB-CG-OD1	7.28	124.85	118.30
1	B	371	ARG	NE-CZ-NH2	6.78	123.69	120.30
1	B	344	ASP	CB-CG-OD2	-6.57	112.39	118.30
1	A	53	ARG	NE-CZ-NH1	6.42	123.51	120.30
1	B	305	GLU	OE1-CD-OE2	-6.33	115.70	123.30
1	B	208	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	A	53	ARG	NE-CZ-NH2	-5.97	117.31	120.30
1	A	205	LEU	CB-CG-CD2	5.97	121.15	111.00
1	A	230	ARG	NE-CZ-NH2	5.90	123.25	120.30
1	B	66	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	B	66	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	A	249	ASP	CB-CG-OD1	5.72	123.44	118.30
1	A	299	LEU	CB-CG-CD2	5.51	120.37	111.00
1	B	147	ASP	CB-CG-OD2	5.48	123.23	118.30
1	A	286	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	B	249	ASP	CB-CG-OD2	-5.44	113.40	118.30
1	A	230	ARG	NE-CZ-NH1	-5.44	117.58	120.30
1	A	123	ARG	NH1-CZ-NH2	5.18	125.09	119.40
1	B	8	ARG	NH1-CZ-NH2	-5.12	113.76	119.40
1	B	294	ARG	NE-CZ-NH2	-5.05	117.77	120.30
1	B	351	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3121	0	3127	40	0
1	B	3135	0	3137	34	0
2	A	6	0	4	2	0
2	B	6	0	4	1	0
3	A	15	0	6	4	0
3	B	15	0	6	2	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	6	0	8	0	0
5	A	4	0	6	3	0
6	A	254	0	0	8	0
6	B	291	0	0	13	2
All	All	6853	0	6298	76	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:LYS:HD3	6:A:664:HOH:O	1.19	1.30
1:A:161:GLN:HE21	1:A:171:HIS:HE1	1.25	0.85
1:B:114:HIS:HD2	1:B:116:SER:H	1.25	0.85
1:A:46:ASN:HD22	1:A:47:VAL:H	1.25	0.84
2:A:400:ALA:N	3:A:401:PLP:C4A	2.41	0.84
1:A:114:HIS:HD2	1:A:116:SER:H	1.26	0.82
1:A:375:HIS:CD2	1:A:376:VAL:H	1.99	0.80
1:A:375:HIS:HD2	1:A:376:VAL:H	1.30	0.78
1:A:53:ARG:NH2	1:B:347:SER:OG	2.18	0.77
1:B:161:GLN:HE21	1:B:171:HIS:HE1	1.32	0.77
1:B:375:HIS:CD2	1:B:376:VAL:H	2.03	0.77
1:B:375:HIS:HD2	1:B:376:VAL:H	1.33	0.74
1:A:178:LYS:HG2	1:A:206:HIS:CE1	2.22	0.74
1:A:114:HIS:CD2	1:A:116:SER:H	2.05	0.73
1:B:114:HIS:CD2	1:B:116:SER:H	2.06	0.73
1:B:1:MET:HE1	1:B:287:ILE:HG12	1.74	0.70
1:A:161:GLN:HE21	1:A:171:HIS:CE1	2.10	0.67
2:B:401:ALA:HB3	6:B:756:HOH:O	1.95	0.67
1:B:1:MET:HE2	1:B:3:ILE:HD11	1.78	0.66
2:A:400:ALA:HB3	6:A:724:HOH:O	1.95	0.65
1:B:48:HIS:HD2	6:B:703:HOH:O	1.81	0.63
1:B:323:HIS:HE1	1:B:326:VAL:O	1.82	0.63
1:B:294:ARG:HG3	6:B:501:HOH:O	1.98	0.63
1:A:49:ARG:HG2	1:A:59:THR:HG21	1.83	0.61
1:B:161:GLN:HE21	1:B:171:HIS:CE1	2.15	0.60
1:B:112:TYR:CZ	1:B:141:GLY:HA2	2.38	0.58
1:A:206:HIS:HE1	6:A:664:HOH:O	1.87	0.58
1:A:171:HIS:HD2	6:A:541:HOH:O	1.86	0.57
1:A:217:GLY:O	1:A:375:HIS:HE1	1.88	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:GLY:O	1:B:375:HIS:HE1	1.88	0.56
1:A:323:HIS:HE1	1:A:326:VAL:O	1.89	0.56
1:B:1:MET:CE	1:B:3:ILE:HD11	2.36	0.55
1:B:174:GLU:HG3	6:B:657:HOH:O	2.07	0.55
1:B:297:HIS:HE1	6:B:753:HOH:O	1.89	0.54
1:B:317:PRO:O	1:B:323:HIS:HD2	1.90	0.54
1:A:193:GLN:HG2	3:A:401:PLP:O3	2.08	0.54
1:A:46:ASN:HD22	1:A:47:VAL:N	2.00	0.53
1:B:206:HIS:HE1	6:B:754:HOH:O	1.92	0.53
1:A:175:GLU:HA	1:A:178:LYS:HE2	1.90	0.52
1:A:57:MET:HE3	6:A:739:HOH:O	2.09	0.51
1:B:48:HIS:CD2	6:B:703:HOH:O	2.60	0.51
1:B:2:ARG:HH11	1:B:2:ARG:HG3	1.75	0.51
1:A:317:PRO:O	1:A:323:HIS:HD2	1.93	0.50
1:A:247:ILE:C	1:A:247:ILE:HD12	2.32	0.50
1:B:114:HIS:HE1	6:B:756:HOH:O	1.95	0.49
1:B:171:HIS:HD2	6:B:632:HOH:O	1.96	0.49
1:A:178:LYS:CG	1:A:206:HIS:CE1	2.95	0.48
1:A:142:ASN:HD22	1:A:322:LYS:NZ	2.10	0.48
1:B:289:LEU:O	1:B:293:GLU:HG3	2.13	0.48
1:B:323:HIS:CE1	1:B:326:VAL:O	2.64	0.48
1:B:354:HIS:HD2	1:B:356:CSS:N	2.11	0.48
1:B:74:GLY:O	1:B:203:LYS:HE3	2.14	0.47
1:A:290:GLY:HA3	5:A:403:EDO:H11	1.97	0.47
1:B:193:GLN:HG2	3:B:400:PLP:O3	2.15	0.47
1:B:193:GLN:NE2	3:B:400:PLP:O3	2.35	0.46
1:B:150:LYS:HE3	6:B:681:HOH:O	2.15	0.46
1:A:193:GLN:NE2	3:A:401:PLP:O3	2.38	0.46
1:A:375:HIS:CD2	1:A:376:VAL:N	2.79	0.46
1:A:29:PRO:HB2	1:A:31:PRO:HD2	1.97	0.45
1:B:294:ARG:NH1	6:B:501:HOH:O	2.44	0.45
1:A:112:TYR:CZ	1:A:141:GLY:HA2	2.52	0.45
1:A:354:HIS:HD2	1:A:356:CSS:N	2.15	0.44
1:A:140:GLU:OE2	1:A:318:ARG:HG2	2.17	0.44
1:B:346:HIS:HD2	6:B:771:HOH:O	2.01	0.44
1:A:174:GLU:O	1:A:178:LYS:HG3	2.18	0.44
1:A:290:GLY:H	5:A:403:EDO:H11	1.85	0.42
1:A:323:HIS:CE1	1:A:326:VAL:O	2.70	0.42
1:A:375:HIS:HD2	1:A:376:VAL:N	2.09	0.42
1:B:99:HIS:HD2	6:B:665:HOH:O	2.03	0.41
1:A:297:HIS:HD2	6:A:645:HOH:O	2.02	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:287:ILE:O	1:B:291:ARG:HD2	2.21	0.41
1:A:48:HIS:HE1	6:A:545:HOH:O	2.03	0.41
1:A:290:GLY:H	5:A:403:EDO:C1	2.34	0.41
1:A:190:ASP:OD2	3:A:401:PLP:N1	2.54	0.40
1:A:114:HIS:HE1	6:A:724:HOH:O	2.04	0.40
1:A:49:ARG:NH2	1:A:63:GLU:OE2	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:605:HOH:O	6:B:746:HOH:O[1_455]	1.84	0.36
6:B:678:HOH:O	6:B:746:HOH:O[1_455]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/419 (95%)	392 (99%)	4 (1%)	1 (0%)	41	18
1	B	399/419 (95%)	393 (98%)	5 (1%)	1 (0%)	41	18
All	All	796/838 (95%)	785 (99%)	9 (1%)	2 (0%)	41	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	47	VAL
1	A	47	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/350 (95%)	329 (99%)	3 (1%)	78	61
1	B	333/350 (95%)	331 (99%)	2 (1%)	86	74
All	All	665/700 (95%)	660 (99%)	5 (1%)	81	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	334	LEU
1	A	397	LEU
1	B	232	GLU
1	B	318	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	46	ASN
1	A	48	HIS
1	A	114	HIS
1	A	142	ASN
1	A	171	HIS
1	A	206	HIS
1	A	295	GLN
1	A	297	HIS
1	A	323	HIS
1	A	354	HIS
1	A	375	HIS
1	B	99	HIS
1	B	114	HIS
1	B	142	ASN
1	B	171	HIS
1	B	206	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	297	HIS
1	B	323	HIS
1	B	346	HIS
1	B	354	HIS
1	B	375	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSS	B	356	1	4,6,7	1.22	0	1,6,8	0.53	0
1	CSS	A	356	1	4,6,7	0.82	0	1,6,8	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	B	356	1	-	0/1/5/7	-
1	CSS	A	356	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.



There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	356	CSS	1	0
1	A	356	CSS	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	402	-	5,5,5	0.88	0	5,5,5	1.42	1 (20%)
2	ALA	B	401	3	5,5,5	1.18	0	6,6,6	2.56	5 (83%)
3	PLP	B	400	1,2	15,15,16	1.40	2 (13%)	20,22,23	1.97	5 (25%)
5	EDO	A	403	-	3,3,3	0.22	0	2,2,2	1.41	0
3	PLP	A	401	1	15,15,16	2.65	5 (33%)	20,22,23	2.31	6 (30%)
2	ALA	A	400	-	5,5,5	0.95	0	6,6,6	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	402	-	-	0/4/4/4	-
2	ALA	B	401	3	-	2/4/4/4	-
3	PLP	B	400	1,2	-	0/6/6/8	0/1/1/1
5	EDO	A	403	-	-	1/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PLP	A	401	1	-	0/6/6/8	0/1/1/1
2	ALA	A	400	-	-	2/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	PLP	C3-C2	6.36	1.47	1.40
3	A	401	PLP	C5-C4	5.13	1.46	1.40
3	A	401	PLP	C3-C4	3.72	1.47	1.40
3	B	400	PLP	C4A-C4	3.27	1.58	1.51
3	A	401	PLP	P-O2P	-3.01	1.43	1.54
3	A	401	PLP	P-O3P	-2.54	1.45	1.54
3	B	400	PLP	P-O3P	-2.27	1.46	1.54

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	PLP	C6-C5-C4	6.73	123.45	118.16
3	B	400	PLP	C5-C6-N1	-4.04	117.09	123.82
3	B	400	PLP	C6-N1-C2	3.85	126.31	119.17
3	B	400	PLP	C6-C5-C4	3.75	121.11	118.16
3	A	401	PLP	C5-C6-N1	-3.67	117.71	123.82
3	A	401	PLP	C6-N1-C2	3.52	125.68	119.17
2	B	401	ALA	OXT-C-CA	3.40	126.36	114.06
3	A	401	PLP	C4-C3-C2	-3.25	115.27	120.07
3	B	400	PLP	C3-C2-N1	-3.22	116.60	120.77
3	A	401	PLP	O3P-P-O2P	3.10	119.49	107.64
3	B	400	PLP	C4A-C4-C5	2.89	123.92	120.94
2	B	401	ALA	C-CA-N	2.84	118.20	107.60
2	B	401	ALA	OXT-C-O	-2.80	117.73	124.09
3	A	401	PLP	O3-C3-C2	2.78	123.56	117.49
4	A	402	GOL	O1-C1-C2	2.50	122.20	110.20
2	B	401	ALA	CB-CA-C	-2.06	105.52	110.29
2	B	401	ALA	CB-CA-N	2.04	116.28	109.81

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	ALA	O-C-CA-CB
5	A	403	EDO	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	400	ALA	O-C-CA-CB
2	A	400	ALA	OXT-C-CA-CB
2	B	401	ALA	OXT-C-CA-CB

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	ALA	1	0
3	B	400	PLP	2	0
5	A	403	EDO	3	0
3	A	401	PLP	4	0
2	A	400	ALA	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	399/419 (95%)	0.15	9 (2%) 60 65	10, 19, 36, 54	0
1	B	401/419 (95%)	-0.08	6 (1%) 73 78	9, 16, 29, 46	0
All	All	800/838 (95%)	0.03	15 (1%) 66 71	9, 18, 34, 54	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	397	LEU	4.6
1	A	0	SER	4.4
1	B	397	LEU	3.8
1	A	-1	ALA	3.8
1	A	333	GLY	3.4
1	A	366	ILE	3.2
1	A	365	GLY	3.1
1	B	92	LEU	2.8
1	B	93	VAL	2.8
1	A	367	ASN	2.7
1	B	-4	GLY	2.7
1	A	-2	MET	2.5
1	B	-3	HIS	2.4
1	B	49	ARG	2.3
1	A	227	LEU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSS	A	356	7/8	0.96	0.07	17,18,25,25	0
1	CSS	B	356	7/8	0.96	0.07	17,18,27,29	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ALA	A	400	6/6	0.87	0.11	25,29,33,33	0
2	ALA	B	401	6/6	0.90	0.09	27,34,37,37	0
4	GOL	A	402	6/6	0.91	0.10	20,22,25,29	0
5	EDO	A	403	4/4	0.92	0.12	25,33,33,34	0
3	PLP	A	401	15/16	0.98	0.09	12,24,32,36	0
3	PLP	B	400	15/16	0.98	0.10	11,21,33,34	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.