



wwPDB EM Validation Summary Report ⓘ

Nov 16, 2024 – 02:23 PM EST

PDB ID : 9B9G
EMDB ID : EMD-44382
Title : Structure of the PI4KA complex bound to Calcineurin
Authors : Shaw, A.L.; Suresh, S.; Yip, C.K.; Burke, J.E.
Deposited on : 2024-04-02
Resolution : 3.50 Å(reported)
Based on initial models : 6NUC, 6BQ1

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2102	61% 19% 19%
1	B	2102	58% 21% 19%
2	D	843	50% 18% 31%
2	F	843	51% 16% 31%
3	E	308	58% 24% 18%
3	G	308	5% 50% 30% 18%
4	H	170	52% 69% 22% 9%
4	J	170	52% 73% 18% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	I	620	
5	K	620	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 48676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4-kinase alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1696	Total	C	N	O	S	0	0
			13625	8815	2266	2447	97		
1	B	1696	Total	C	N	O	S	0	0
			13625	8815	2266	2447	97		

- Molecule 2 is a protein called Tetratricopeptide repeat protein 7B.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	D	578	Total	C	N	O	S	0	0
			4573	2918	793	835	27		
2	F	578	Total	C	N	O	S	0	0
			4573	2918	793	835	27		

- Molecule 3 is a protein called Hyccin.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	254	Total	C	N	O	S	0	0
			2022	1315	325	370	12		
3	G	254	Total	C	N	O	S	0	0
			2022	1315	325	370	12		

- Molecule 4 is a protein called Calcineurin subunit B type 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	H	155	Total	C	N	O	S	1	0
			1236	779	205	246	6		
4	J	155	Total	C	N	O	S	1	0
			1236	779	205	246	6		

- Molecule 5 is a protein called Protein phosphatase 3 catalytic subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	I	353	2866	1849	478	519	20	0	0
5	K	356	2890	1863	485	522	20	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	236	PRO	LEU	engineered mutation	UNP Q08209
I	238	ASN	ASP	engineered mutation	UNP Q08209
I	466	PRO	LEU	engineered mutation	UNP Q08209
K	236	PRO	LEU	engineered mutation	UNP Q08209
K	238	ASN	ASP	engineered mutation	UNP Q08209
K	466	PRO	LEU	engineered mutation	UNP Q08209


- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		AltConf
6	H	4	Total	Ca	0
			4	4	
6	J	4	Total	Ca	0
			4	4	

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphatidylinositol 4-kinase alpha







• Molecule 1: Phosphatidylinositol 4-kinase alpha



GLU	F1737	W1596	THR	M1367	THR	M1010	SER	L767	LYS	L611	THR	THR	Y366	PHE
CYS	F1738	H1597	ILE	I1370	GLY	M1010	ALA	G768	ASP	I612	VAL	VAL	A448	HIS
SER	T1598	T1598	SER	I1371	GLN	S1020	MET	V769	TRR	P613	ALA	A449	F369	TRR
THR	K1740	I1599	LYS	L1371	MET	S1020	LYS	V770	LYS	T616	GLY	GLY	L374	PHE
GLN	D1600	D1600	LYS	E1376	SER	D1024	ASN	I771	ASP	T616	ASN	ASP	Y374	GLU
GLU	A1601	A1601	GLY	E1376	D1168	D1024	ASP	P772	HIS	L620	ASP	ASP	M377	ALA
ALA	K1759	D1602	THR	K1391	THR	K1027	VAL	W773	ILE	L620	ILE	ILE	L462	SER
ASP	L1762	E1605	MET	K1391	M1172	K1027	THR	I774	K637	G621	K637	K637	L462	LYS
GLY	E1605	E1605	SER	K1398	H1185	D1033	T857	L777	S706	H622	L466	L466	L466	LEU
ASN	L1606	L1606	LYS	K1398	H1186	D1033	E860	L777	Q467	I623	Q467	Q467	Q467	LEU
THR	H1608	H1608	THR	R1401	Q1187	D1045	R865	L781	A708	L627	L707	L707	L707	PRO
THR	W1612	W1612	ASN	F1415	Q1187	T1046	I868	L781	W709	L627	S468	S468	S468	ASP
THR	A1613	A1613	THR	K1419	T1190	Y1047	I868	W773	I710	L627	L381	L381	L381	GLY
THR	S1479	S1479	TYR	K1419	M1193	E1048	I868	I774	I710	L627	R382	R382	R382	THR
LEU	K1483	K1483	LEU	TYR	M1193	A1049	P874	L777	N711	L627	T384	T384	T384	ALA
THR	Y1484	Y1484	THR	THR	M1193	A1049	P874	L781	GLU	L627	L384	L384	L384	ALA
ALA	Y1485	Y1485	ALA	ALA	T1197	I1053	R874	L781	SER	L627	L384	L384	L384	ALA
SER	M1486	M1486	SER	ALA	I1201	I1054	W878	L781	THR	L627	L384	L384	L384	ALA
GLN	K1487	K1487	GLN	GLN	I1202	I1064	F887	L781	VAL	L627	L384	L384	L384	ALA
LEU	R1488	R1488	LEU	TYR	S1202	I1064	A888	L781	GLN	L627	L384	L384	L384	ALA
VAL	R1489	R1489	VAL	TYR	D1205	L1084	M889	L781	GLY	L627	L384	L384	L384	ALA
PRO	T1490	T1490	PRO	THR	M1227	L1100	L894	L781	LEU	L627	L384	L384	L384	ALA
ASP	E1501	E1501	ASP	THR	G1240	T1104	M902	L781	PRO	L627	L384	L384	L384	ALA
ASN	R1502	R1502	ASN	THR	V1246	H1109	R903	L781	PRO	L627	L384	L384	L384	ALA
THR	T1505	T1505	THR	THR	P1247	PHE	R903	L781	PRO	L627	L384	L384	L384	ALA
ARG	E1516	E1516	ARG	THR	F1264	GLY	R903	L781	PRO	L627	L384	L384	L384	ALA
ASN	M1529	M1529	ASN	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
LEU	S1530	S1530	LEU	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
ASP	K1531	K1531	ASP	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
ILE	Y1532	Y1532	ILE	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
THR	I1533	I1533	THR	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
VAL	S1534	S1534	VAL	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
GLY	L1535	L1535	GLY	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
ARG	S1536	S1536	ARG	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
GLN	K1541	K1541	GLN	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
GLN	M1545	M1545	GLN	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
THR	G1569	G1569	THR	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
GLN	G1569	G1569	GLN	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
GLY	V1572	V1572	GLY	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
TRP	T1573	T1573	TRP	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
ILE	L1711	L1711	ILE	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
ASN	L1712	L1712	ASN	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
THR	D1713	D1713	THR	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
TYR	Q1714	Q1714	TYR	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
PRO	L1715	L1715	PRO	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
LEU	V1716	V1716	LEU	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
SER	V1586	V1586	SER	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
ASP	P1587	P1587	ASP	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
GLY	K1591	K1591	GLY	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
MET	F1735	F1735	MET	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA
SER	D1736	D1736	SER	THR	F1264	TYR	R903	L781	PRO	L627	L384	L384	L384	ALA




● Molecule 3: Hyccin




● Molecule 3: Hyccin







• Molecule 4: Calcineurin subunit B type 1



• Molecule 4: Calcineurin subunit B type 1



• Molecule 5: Protein phosphatase 3 catalytic subunit alpha



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	235760	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	165000	Depositor
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.526	Depositor
Minimum map value	-0.538	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.038	Depositor
Recommended contour level	0.155	Depositor
Map size (Å)	415.8, 415.8, 415.8	wwPDB
Map dimensions	520, 520, 520	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.7996154, 0.7996154, 0.7996154	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/13924	0.51	7/18834 (0.0%)
1	B	0.25	0/13924	0.51	3/18834 (0.0%)
2	D	0.24	0/4656	0.51	2/6296 (0.0%)
2	F	0.25	0/4656	0.53	2/6296 (0.0%)
3	E	0.25	0/2070	0.49	0/2809
3	G	0.26	0/2070	0.54	1/2809 (0.0%)
4	H	0.25	0/1254	0.52	0/1678
4	J	0.26	0/1254	0.54	0/1678
5	I	0.26	0/2944	0.57	0/3996
5	K	0.26	0/2968	0.57	0/4026
All	All	0.25	0/49720	0.52	15/67256 (0.0%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	LEU	CA-CB-CG	7.81	133.26	115.30
1	B	139	LEU	CA-CB-CG	7.59	132.77	115.30
1	A	101	LEU	CA-CB-CG	6.83	131.01	115.30
2	D	825	LEU	CA-CB-CG	6.38	129.98	115.30
2	D	574	MET	CA-CB-CG	6.29	123.99	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	13625	0	13777	260	0
1	B	13625	0	13777	308	0
2	D	4573	0	4632	103	0
2	F	4573	0	4632	100	0
3	E	2022	0	2027	50	0
3	G	2022	0	2027	63	0
4	H	1236	0	1216	33	0
4	J	1236	0	1216	30	0
5	I	2866	0	2805	91	0
5	K	2890	0	2824	89	0
6	H	4	0	0	0	0
6	J	4	0	0	0	0
All	All	48676	0	48933	1106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 1106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:332:LEU:HD13	2:F:369:LEU:HD22	1.60	0.84
1:B:787:ALA:HB1	1:B:791:LEU:HD12	1.59	0.83
3:G:274:GLU:HG2	3:G:275:PRO:HD3	1.62	0.82
1:A:1865:ILE:HG12	1:A:1949:LEU:HD21	1.60	0.81
3:E:274:GLU:HG2	3:E:275:PRO:HD3	1.63	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1644/2102 (78%)	1552 (94%)	92 (6%)	0	100	100
1	B	1644/2102 (78%)	1542 (94%)	102 (6%)	0	100	100
2	D	566/843 (67%)	549 (97%)	17 (3%)	0	100	100
2	F	566/843 (67%)	546 (96%)	20 (4%)	0	100	100
3	E	244/308 (79%)	236 (97%)	8 (3%)	0	100	100
3	G	244/308 (79%)	231 (95%)	13 (5%)	0	100	100
4	H	154/170 (91%)	154 (100%)	0	0	100	100
4	J	154/170 (91%)	153 (99%)	1 (1%)	0	100	100
5	I	349/620 (56%)	337 (97%)	12 (3%)	0	100	100
5	K	350/620 (56%)	338 (97%)	12 (3%)	0	100	100
All	All	5915/8086 (73%)	5638 (95%)	277 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1508/1841 (82%)	1453 (96%)	55 (4%)	30	59
1	B	1508/1841 (82%)	1438 (95%)	70 (5%)	23	52
2	D	483/712 (68%)	458 (95%)	25 (5%)	19	47
2	F	483/712 (68%)	450 (93%)	33 (7%)	13	39
3	E	229/277 (83%)	222 (97%)	7 (3%)	35	63
3	G	229/277 (83%)	210 (92%)	19 (8%)	9	32
4	H	137/151 (91%)	136 (99%)	1 (1%)	81	89
4	J	137/151 (91%)	136 (99%)	1 (1%)	81	89
5	I	316/547 (58%)	306 (97%)	10 (3%)	34	62
5	K	318/547 (58%)	311 (98%)	7 (2%)	47	70
All	All	5348/7056 (76%)	5120 (96%)	228 (4%)	27	54

5 of 228 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1942	SER
5	K	42	ARG
2	D	574	MET
5	I	347	MET
3	G	159	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	543	GLN
3	G	166	HIS
2	D	738	GLN
2	F	738	GLN
5	I	49	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-44382. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 260




Y Index: 260




Z Index: 260


6.2.2 Raw map



X Index: 260



Y Index: 260




Z Index: 260


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 250




Y Index: 320




Z Index: 281


6.3.2 Raw map



X Index: 251



Y Index: 214




Z Index: 280


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.155. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 722 nm^3 ; this corresponds to an approximate mass of 652 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8.2 Resolution estimates


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.51	4.12	3.58
Unmasked-calculated*	4.51	6.93	4.69

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.51 differs from the reported value 3.5 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-44382 and PDB model 9B9G. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.155 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.155).























9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.155) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8440	 0.3010
A	 0.9470	 0.3710
B	 0.9400	 0.3570
D	 0.8880	 0.3680
E	 0.8760	 0.2990
F	 0.8560	 0.3340
G	 0.8120	 0.2340
H	 0.3890	 0.0750
I	 0.5740	 0.0490
J	 0.3630	 0.0670
K	 0.4960	 0.0350

