



Full wwPDB X-ray Structure Validation Report

May 14, 2020 – 11:00 pm BST

PDB ID : 3BBT
Title : crystal structure of the ErbB4 kinase in complex with lapatinib
Authors : Qiu, C.
Deposited on : 2007-11-11
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	328	 69% 11% 16%
1	D	328	 65% 14% 21%

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 4431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor tyrosine-protein kinase erbB-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	274	Total 2201	C 1420	N 380	O 382	S 19	0	0	0
1	D	259	Total 2091	C 1352	N 361	O 360	S 18	0	0	0

- Molecule 2 is N-{3-CHLORO-4-[(3-FLUOROBENZYL)OXY]PHENYL}-6-[5-({[2-(METHYLSULFONYL)ETHYL]AMINO}METHYL)-2-FURYL]-4-QUINAZOLINAMINE (three-letter code: FMM) (formula: C₂₉H₂₆ClFN₄O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Cl	F	N			O
2	B	1	Total 27	C 21	Cl 1	F 1	N 3	O 1	0	0
2	D	1	Total 28	C 22	Cl 1	F 1	N 3	O 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	44	Total 44	O 44	0	0
3	D	40	Total 40	O 40	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Receptor tyrosine-protein kinase erbB-4

Chain B: 



- Molecule 1: Receptor tyrosine-protein kinase erbB-4

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	102.68Å 102.68Å 185.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.64 – 2.80 29.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.64-2.80) 99.9 (29.64-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.80Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.251 , 0.289 0.245 , 0.281	Depositor DCC
R_{free} test set	1381 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	65.0	Xtrriage
Anisotropy	0.358	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtrriage
Estimated twinning fraction	0.334 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4431	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.35	0/2251	0.56	3/3043 (0.1%)
1	D	0.34	0/2139	0.51	1/2891 (0.0%)
All	All	0.34	0/4390	0.53	4/5934 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	965	ASP	CB-CG-OD2	5.65	123.39	118.30
1	D	818	ASP	CB-CG-OD2	5.51	123.26	118.30
1	B	818	ASP	CB-CG-OD2	5.21	122.99	118.30
1	B	966	ASP	CB-CG-OD2	5.00	122.80	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	720	LYS	Peptide
1	B	721	ILE	Peptide
1	B	733	GLY	Peptide
1	B	972	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2201	0	2264	38	0
1	D	2091	0	2153	31	0
2	B	27	0	14	0	0
2	D	28	0	14	4	0
3	B	44	0	0	0	0
3	D	40	0	0	0	0
All	All	4431	0	4445	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:720:LYS:O	1:B:721:ILE:HG12	1.39	1.21
1:B:720:LYS:C	1:B:721:ILE:HG12	1.55	1.09
1:B:965:ASP:CG	1:B:966:ASP:H	1.81	0.84
1:B:720:LYS:C	1:B:721:ILE:CG1	2.42	0.82
1:B:721:ILE:N	1:B:722:PRO:HD2	1.95	0.81
1:D:816:HIS:O	1:D:817:ARG:HB2	1.80	0.78
1:B:733:GLY:O	1:B:843:LEU:HD21	1.88	0.74
1:B:696:VAL:O	1:B:697:LYS:HG3	1.90	0.72
1:B:727:ILE:HG12	1:B:768:GLN:HG2	1.73	0.71
1:B:816:HIS:O	1:B:817:ARG:HB2	1.89	0.71
1:B:817:ARG:NH1	1:B:840:ALA:HB1	2.04	0.70
1:B:720:LYS:O	1:B:721:ILE:CG1	2.30	0.68
1:D:886:TRP:HD1	1:D:928:MET:CE	2.07	0.68
1:D:727:ILE:HG12	1:D:768:GLN:HG2	1.79	0.65
1:B:965:ASP:OD1	1:B:966:ASP:N	2.30	0.65
1:B:965:ASP:CG	1:B:966:ASP:N	2.50	0.65
1:B:963:GLN:NE2	1:B:969:LYS:H	1.96	0.63
1:D:713:VAL:HG22	1:D:720:LYS:HG2	1.82	0.62
1:B:817:ARG:NH1	1:B:840:ALA:CB	2.63	0.61
1:B:696:VAL:O	1:B:697:LYS:CG	2.50	0.59
1:D:885:ILE:O	1:D:889:MET:HG2	2.05	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:904:ILE:N	1:D:905:PRO:HD2	2.21	0.55
1:B:772:GLN:HE22	1:B:827:LYS:HE2	1.72	0.55
1:D:886:TRP:HD1	1:D:928:MET:HE1	1.71	0.55
1:B:951:ARG:HA	1:B:954:ARG:HH11	1.73	0.54
1:D:886:TRP:HD1	1:D:928:MET:HE3	1.73	0.53
1:B:696:VAL:C	1:B:697:LYS:HG3	2.28	0.53
1:D:815:VAL:HG12	1:D:817:ARG:HG3	1.91	0.52
1:B:726:LYS:HB3	1:B:769:LEU:HB2	1.91	0.51
1:B:723:VAL:HG22	1:B:771:THR:O	2.11	0.51
1:D:718:THR:HG23	1:D:719:VAL:HG13	1.93	0.50
1:B:735:LYS:O	1:B:737:ASN:OD1	2.30	0.49
1:D:769:LEU:HD23	2:D:91:FMM:H282	1.94	0.49
1:D:689:LYS:HB2	1:D:692:GLU:HB2	1.93	0.48
1:B:963:GLN:HE22	1:B:969:LYS:HB2	1.78	0.48
1:B:733:GLY:O	1:B:843:LEU:CD2	2.60	0.48
1:B:864:LEU:HD21	1:B:909:GLU:HG2	1.95	0.48
1:D:816:HIS:ND1	1:D:816:HIS:C	2.67	0.48
1:B:736:ALA:C	1:B:737:ASN:OD1	2.52	0.48
1:D:792:SER:HB3	1:D:960:LEU:HB2	1.95	0.47
1:D:752:HIS:CD2	1:D:754:HIS:H	2.33	0.47
1:D:754:HIS:CE1	1:D:801:GLN:HG2	2.49	0.47
1:D:884:THR:O	1:D:887:GLU:HB2	2.15	0.47
1:D:906:ASP:N	1:D:906:ASP:OD1	2.46	0.47
1:B:750:MET:HE1	1:B:809:LEU:HD23	1.97	0.47
1:B:735:LYS:O	1:B:737:ASN:N	2.48	0.46
1:D:776:HIS:HB2	1:D:826:VAL:HB	1.96	0.46
1:D:886:TRP:CD1	1:D:928:MET:HE1	2.51	0.45
1:D:886:TRP:CD1	1:D:928:MET:CE	2.96	0.44
1:D:837:PHE:CZ	2:D:91:FMM:H32	2.53	0.44
1:B:696:VAL:HG23	1:B:709:LYS:O	2.17	0.44
1:B:701:SER:HB3	1:B:706:THR:HG23	1.99	0.43
1:B:720:LYS:HB2	1:D:720:LYS:H	1.82	0.43
1:D:774:MET:HA	1:D:775:PRO:HD3	1.83	0.43
1:D:777:GLY:HA2	2:D:91:FMM:C16	2.49	0.42
1:D:946:ALA:O	1:D:950:SER:HB2	2.18	0.42
1:D:721:ILE:HA	1:D:722:PRO:HD2	1.78	0.42
1:B:809:LEU:HD22	1:B:814:LEU:HD22	2.01	0.42
1:D:774:MET:H	2:D:91:FMM:H19	1.84	0.42
1:D:894:LYS:HA	1:D:895:PRO:HD3	1.91	0.41
1:B:817:ARG:HH11	1:B:840:ALA:CB	2.34	0.41
1:B:816:HIS:ND1	1:B:816:HIS:C	2.73	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:750:MET:CE	1:B:809:LEU:HD23	2.50	0.41
1:B:963:GLN:HE22	1:B:969:LYS:H	1.68	0.41
1:D:886:TRP:CD1	1:D:928:MET:HE3	2.54	0.41
1:B:688:LEU:HD12	1:B:770:VAL:HG21	2.03	0.40
1:D:762:CYS:HB3	1:D:768:GLN:HB2	2.03	0.40
1:B:734:PRO:O	1:B:767:ILE:HD13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	266/328 (81%)	256 (96%)	6 (2%)	4 (2%)	10	33
1	D	251/328 (76%)	244 (97%)	6 (2%)	1 (0%)	34	66
All	All	517/656 (79%)	500 (97%)	12 (2%)	5 (1%)	15	44

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	722	PRO
1	B	736	ALA
1	B	734	PRO
1	B	765	PRO
1	D	685	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	243/288 (84%)	231 (95%)	12 (5%)	25	57
1	D	231/288 (80%)	227 (98%)	4 (2%)	60	87
All	All	474/576 (82%)	458 (97%)	16 (3%)	37	71

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	696	VAL
1	B	713	VAL
1	B	721	ILE
1	B	742	ASP
1	B	771	THR
1	B	773	LEU
1	B	788	ASP
1	B	913	ARG
1	B	924	VAL
1	B	958	ARG
1	B	963	GLN
1	B	972	SER
1	D	723	VAL
1	D	728	LEU
1	D	835	THR
1	D	901	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	957	GLN
1	B	963	GLN
1	D	752	HIS

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FMM	B	91	-	30,30,44	1.20	2 (6%)	41,41,62	1.92	9 (21%)
2	FMM	D	91	-	31,31,44	1.17	2 (6%)	43,43,62	1.93	10 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMM	B	91	-	-	5/9/9/21	0/4/4/5
2	FMM	D	91	-	-	5/9/9/21	0/4/4/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	91	FMM	C21-C39	-3.75	1.40	1.44
2	B	91	FMM	C21-C39	-3.67	1.40	1.44
2	B	91	FMM	C36-CL3	2.78	1.80	1.73
2	D	91	FMM	C36-CL3	2.68	1.80	1.73

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	91	FMM	C19-N20-C21	5.61	121.40	116.59
2	B	91	FMM	C19-N20-C21	5.50	121.31	116.59
2	D	91	FMM	N18-C19-N20	-4.95	120.94	128.68
2	B	91	FMM	N18-C19-N20	-4.89	121.03	128.68
2	B	91	FMM	C21-C39-C17	4.64	118.80	115.88
2	D	91	FMM	C21-C39-C17	4.03	118.41	115.88
2	B	91	FMM	C19-N18-C17	3.71	120.51	115.40
2	D	91	FMM	C40-C39-C21	-3.68	121.56	124.88
2	D	91	FMM	C19-N18-C17	3.66	120.45	115.40
2	B	91	FMM	C39-C21-N20	-3.26	118.67	121.35
2	D	91	FMM	C39-C21-N20	-3.07	118.82	121.35
2	B	91	FMM	C39-C17-N18	-2.96	119.68	122.83
2	D	91	FMM	C39-C17-N18	-2.67	119.99	122.83
2	D	91	FMM	C32-C33-C35	-2.67	119.82	123.29
2	B	91	FMM	C28-O27-C26	2.58	122.83	117.76
2	B	91	FMM	C32-C33-C35	-2.36	120.22	123.29
2	D	91	FMM	C23-N22-C21	-2.32	122.21	128.26
2	B	91	FMM	O27-C26-C36	2.27	119.21	116.40
2	D	91	FMM	C28-O27-C26	2.12	121.93	117.76

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	91	FMM	C36-C26-O27-C28
2	D	91	FMM	C36-C26-O27-C28
2	D	91	FMM	C29-C28-O27-C26
2	B	91	FMM	C29-C28-O27-C26
2	B	91	FMM	C25-C26-O27-C28
2	B	91	FMM	C24-C23-N22-C21
2	D	91	FMM	C24-C23-N22-C21
2	D	91	FMM	C25-C26-O27-C28
2	B	91	FMM	C38-C23-N22-C21
2	D	91	FMM	C38-C23-N22-C21


There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	91	FMM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)


Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.